

Week Ten

Siva Sundar, EE23B151

December 7, 2024

2nd December

- **Graphs:** consists of the following:

- Set V which contains **vertices** and set A with **arrows**,
- s and t are the **source** and **target** functions respectively.

Note: From every graph we can get a *preorder*. **Hasse Diagram** is a graph that gives a *presentation* of a preorder (P, \leq) . (See page.14)

- **Total order:** They are *posets* (partially ordered sets), with an additional condition: “for all x, y , either $x \leq y$ or $y \leq x$ ”. (They should be *comparable*)
- **Partitions** can be made from preorders. (See page.16)
- Preorder of **upper sets** ($U(X)$ contains q , if $p, q \in X$ and $p \leq q$) on a *discrete preorder* on set X is same as power set $P(X)$.
- **Product Preorder:** Given (P, \leq) and (Q, \leq) , we define $(P \times Q, \leq)$ such that:

$$(p, q) \leq (p', q') \iff p \leq p' \ \& \ q \leq q'$$

4th December

- **Monotone map** is a *structure preserving* function $f : A \rightarrow B$, such that:

$$\forall x, y \in A, \text{ if } x \leq_A y \text{ then } f(x) \leq_B f(y).$$

Cardinality is a function which maps a set to a natural number (which is the number of elements in the set). This function is a monotone map, as:

$$\text{if } X \subseteq Y, \text{ then } n(X) \leq n(Y).$$

If a map $f : X \rightarrow Y$ exists, then there exists a monotone map $g : \text{Prt}(Y) \rightarrow \text{Prt}(X)$. ($\text{Prt}(X)$ gives the set of all partitions on X).

If f and g are monotones, then $f \circ g$ is also monotone.

Let P be a preorder. Monotone maps $P \rightarrow \mathcal{B}$ are in one-to-one correspondence with upper sets of P . (See page.22).

- **Yoneda Lemma:** to know an element is the same as knowing its upper set (the relationships it has with other elements). (see page.20).
- **Pullback map:** Let P and Q be preorders, and $f : P \rightarrow Q$ be a monotone map. Then we can define a monotone map $g : U(Q) \rightarrow U(P)$ which is called the *pullback along f* . ($U(X)$ is the set of all uppersets of X).

7th December

- For a preorder (P, \leq) , and $A \subseteq P$ be a subset, we say $p \in P$ is a **meet** of A if
 - ★ $\forall a \in A$, we have $p \leq a$.
 - ★ $\forall q, q \leq a \forall a \in A$, we have $q \leq p$.

We denote meet ' p ' as: $p \cong \bigwedge A$ or $p \cong \bigwedge_{a \in A} a$. This represents the *greatest lower bound* of the subset A . As the **GLB** is the “greatest among *all* lower bounds”, we can say this is a **Universal property**.

- Similarly, for the preoreder discussed above, we say p is a **join** of A if:
 - ★ $\forall a \in A$, we have $a \leq p$.
 - ★ $\forall q, a \leq q \forall a \in A$, we have $p \leq q$.

We denote join p as: $p \cong \bigvee A$ or $p \cong \bigvee_{a \in A} a$. This represents the *lowest upper bound* of subset A . This is also a universal property.

- Any two things defined by the **same** universal property are automatically **equivalent** in a way known as '*unique up to unique isomorphism*'. For example, we can see that if there exists two meets p and q for a preorder, they will be isomorphic to each other by definition.
- In a *discrete preorder*, there exist **no meets nor joins**.
- In any partial order (where \cong and $=$ are the same), $p \vee p = p \wedge p = p$. (See page 25)
- In a power set $P(X)$, for subsets, say $A, B \in X$, the meet is their intersection, ie, $A \wedge B = A \cap B$ and their join is their union, $A \vee B = A \cup B$.
- For a preorder P , $A \subseteq B \subseteq P$, then we say
 - ★ if meets of A and B exist, then $\bigwedge B \leq \bigwedge A$
 - ★ if joins of A and B exist, then $\bigvee A \leq \bigvee B$
- A monotone map $f : P \rightarrow Q$ has a **generative effect** if there exist elements $a, b \in P$ such that:

$$f(a) \vee f(b) \not\cong f(a \vee b)$$

If the monotone map doesn't have a generative effect, then it will preserve the meets.

- A **Galois connection** between two preorders P and Q is a **pair of monotone maps** $f : P \rightarrow Q$ and $g : Q \rightarrow P$ such that:

$$f(p) \leq q \iff p \leq g(q)$$

We say f is the *left adjoint* and g is the *right adjoint* of the Galois connection.

- If P and Q are **total orders** and $f : P \rightarrow Q$ and $g : Q \rightarrow P$ are drawn with **arrows bending counterclockwise**, then f is **left adjoint** to g *iff* the arrows **do not cross**. (See page 28)
- Galois connections are a kind of relaxed version of isomorphisms. (Page 30)
- Right adjoints **preserve meets**, and Left adjoints **preserve joins** (See *Adjoint Functor Theorem*). Hence, left adjoints will not have generative effects.

- **Closure operator** $j : P \rightarrow P$ on a preorder P is a **monotone** map with:

- ★ $p \leq j(p)$

- ★ $j(j(p)) \cong j(p)$

They can be made by composing left adjoint f with its right adjoint g . The other composite map $g \circ f$ (*interior map*) satisfies: $(g \circ f)(p) \leq p$.