

Week Six

Siva Sundar, EE23B151

August 2024

19th August

Section 19: Terminal Objects

- From any set (even for **null set**), there is **only one map** to a *singleton* set.
- T is an object in a category \mathcal{C} , which is said to be **terminal** only if for any object X in \mathcal{C} :
 - ★ at least one map exists from X to T .
 - ★ that map should be the only map from X to T .

Using these two conditions we can say that: (See page.229)

“There exists multiple terminal objects which are isomorphic to each other.”

- In the category of *endomaps*, we can say that the **singleton set** equipped with an endomap from the **point to itself**, is a terminal ‘*set-with-endomap*’.

The mapping from an endomap X to this terminal object also follows the ‘structure preserving rule’.




Figure 1: Map from X to T

Section 20: Points

- In the start of this section, we see an example which shows how we can use a *terminal object* (defined in the category) to **select an item** from an object (of the same category). Hence, we can define:

“A point of an object X is the map $T \longrightarrow X$ ”

where, T is the terminal object of the category.

- In different categories, the meaning of the word ‘point’ is different from what we think of. For example, in the category of endomaps, the term ‘point’ refers to **fixed point** (See page.232). So, if an endomap does **not** have a fixed point, we say it doesn't have ‘*points*’ (which doesn't mean it doesn't have elements!) (See page.233)

Category	Terminal object	‘Points of X ’ means...
\mathcal{C}	T	map $T \longrightarrow X$
\mathcal{S}	$\boxed{\bullet}$	element of X
\mathcal{S}° endomaps of sets	$\boxed{\bullet \rightarrow \bullet}$	fixed point or equilibrium state

20th August

Product

- A product of A and B (in category \mathcal{C} , also called **factors of \mathbf{P}**) is:
 - ★ an object P in \mathcal{C} .
 - ★ a pair of maps: $P \xrightarrow{p_1} A$ and $P \xrightarrow{p_2} B$ such that, for every other object X in \mathcal{C} , with pair of maps $X \xrightarrow{q_1} A$ and $X \xrightarrow{q_2} B$, there exist **exactly one map** $X \xrightarrow{q} P$: $q_1 = p_1 \circ q$ and $q_2 = p_2 \circ q$.

From this definition, we can say that if there exists two products sharing the same **factors**, the products must be **isomorphic**.

- ‘3D-Space’ can be considered as the **product** of three *linearly independent* axes.
- For Products in ‘categories of endomaps’,





Figure 2: Product in category S^{\curvearrowright}



Internal diagram of $A^{\curvearrowright} \longrightarrow P \longleftarrow B^{\curvearrowright}$

We get another condition from the ‘*structure preserving rule*’:

$$\gamma(a, b) = (\alpha(a), \beta(b))$$

Proof:

As we know, P contains elements of the type (a, b) where $a \in A$ and $b \in B$.

$$\begin{aligned} p_1(a, b) &= a \quad p_2(a, b) = b \\ \Rightarrow (p_1 + p_2)(a, b) &= a + b \\ (Or) \quad (a, b) &= (p_1 + p_2)^{-1}[a + b] \end{aligned} \tag{1}$$

By structure preserving conditions:

$$\begin{aligned} p_1\gamma &= \alpha p_1 \quad p_2\gamma = \beta p_2 \\ \Rightarrow (p_1 + p_2)\gamma &= \alpha p_1 + \beta p_2 \\ (Or) \quad \gamma &= (p_1 + p_2)^{-1}[\alpha p_1 + \beta p_2] \end{aligned}$$

Now, applying γ on the element (a, b) yields:

$$\begin{aligned} \gamma(a, b) &= (p_1 + p_2)^{-1}[\alpha p_1 + \beta p_2](a, b) \\ \Rightarrow (p_1 + p_2)^{-1}[\alpha p_1(a, b) + \beta p_2(a, b)] \\ \Rightarrow (p_1 + p_2)^{-1}[\alpha(a) + \beta(b)] \\ \boxed{\gamma(a, b) = (\alpha(a), \beta(b))} \end{aligned} \tag{used (1)}$$