

REGRESSION

```

/DESCRIPTIVES MEAN STDDEV CORR SIG N
/MISSING LISTWISE
/STATISTICS COEFF OUTS CI(95) R ANOVA COLLIN TOL CHANGE ZPP
/CRITERIA=PIN(.05) POUT(.10)
/NOORIGIN
/DEPENDENT WLB
/METHOD=ENTER SM FA S H WS PE IE SI.

```

Regression

Notes

Output Created		16-JUL-2026 13:35:57
Comments		
Input	Active Dataset	DataSet5
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	496
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any variable used.
Syntax		REGRESSION /DESCRIPTIVES MEAN STDDEV CORR SIG N /MISSING LISTWISE /STATISTICS COEFF OUTS CI(95) R ANOVA COLLIN TOL CHANGE ZPP /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT WLB /METHOD=ENTER SM FA S H WS PE IE SI.
Resources	Processor Time	00:00:00.11
	Elapsed Time	00:00:00.20
	Memory Required	8268 bytes
	Additional Memory	
	Required for Residual Plots	0 bytes

Descriptive Statistics

	Mean	Std. Deviation	N
WLB	3.1382	.45741	496
SM	2.7367	.57330	496
FA	2.8475	.49894	496
S	2.6689	.58444	496
H	2.3700	.62192	496
WS	3.1199	.44966	496
PE	1.5444	.29653	496
IE	1.4311	.20861	496
SI	2.6689	.58444	496

Correlations

		WLB	SM	FA	S	H	WS	PE	IE	SI
Pearson Correlation	WLB	1.000	.552	.417	.282	.144	.258	.107	.541	.282
	SM	.552	1.000	.553	.488	.434	.263	.098	.188	.488
	FA	.417	.553	1.000	.515	.340	.587	.264	.258	.515
	S	.282	.488	.515	1.000	.365	.286	.438	.379	1.000
	H	.144	.434	.340	.365	1.000	.226	.287	-.173	.365
	WS	.258	.263	.587	.286	.226	1.000	.417	.171	.286
	PE	.107	.098	.264	.438	.287	.417	1.000	.245	.438
	IE	.541	.188	.258	.379	-.173	.171	.245	1.000	.379
	SI	.282	.488	.515	1.000	.365	.286	.438	.379	1.000
Sig. (1-tailed)	WLB	.	.000	.000	.000	.001	.000	.009	.000	.000
	SM	.000	.	.000	.000	.000	.000	.014	.000	.000
	FA	.000	.000	.	.000	.000	.000	.000	.000	.000
	S	.000	.000	.000	.	.000	.000	.000	.000	.000
	H	.001	.000	.000	.000	.	.000	.000	.000	.000
	WS	.000	.000	.000	.000	.000	.	.000	.000	.000
	PE	.009	.014	.000	.000	.000	.000	.	.000	.000
	IE	.000	.000	.000	.000	.000	.000	.000	.	.000
	SI	.000	.000	.000	.000	.000	.000	.000	.000	.
N	WLB	496	496	496	496	496	496	496	496	496
	SM	496	496	496	496	496	496	496	496	496
	FA	496	496	496	496	496	496	496	496	496
	S	496	496	496	496	496	496	496	496	496
	H	496	496	496	496	496	496	496	496	496
	WS	496	496	496	496	496	496	496	496	496
	PE	496	496	496	496	496	496	496	496	496
	IE	496	496	496	496	496	496	496	496	496
	SI	496	496	496	496	496	496	496	496	496

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	SI, WS, IE, SM, PE, H, FA ^b	.	Enter

a. Dependent Variable: WLB

b. Tolerance = .000 limit reached.

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Change Statistics				
					R Square Change	F Change	df1	df2	Sig. F Change
1	.734 ^a	.539	.533	.31263	.539	81.664	7	488	.000

a. Predictors: (Constant), SI, WS, IE, SM, PE, H, FA

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	55.870	7	7.981	81.664	.000 ^b
	Residual	47.695	488	.098		
	Total	103.565	495			

a. Dependent Variable: WLB

b. Predictors: (Constant), SI, WS, IE, SM, PE, H, FA

Coefficients^a

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.	95.0% Confidence Interval for B		Correlations			Collinearity Statistics	
	B	Std. Error	Beta			Lower Bound	Upper Bound	Zero-order	Partial	Part	Tolerance	VIF
1	(Constant)	.469	.140	3.358	.001	.194	.743					
	SM	.370	.033	11.215	.000	.305	.435	.552	.453	.345	.553	1.809
	FA	.078	.043	1.829	.068	-.006	.162	.417	.083	.056	.437	2.287
	H	.062	.029	2.152	.032	.005	.119	.144	.097	.066	.613	1.630
	WS	.059	.042	1.410	.159	-.023	.140	.258	.064	.043	.566	1.768
	PE	-.069	.060	-1.152	.250	-.186	.048	.107	-.052	-.035	.632	1.582
	IE	1.162	.081	14.392	.000	1.003	1.321	.541	.546	.442	.696	1.437
	SI	-.169	.034	-2.16	.000	-.235	-.103	.282	-.221	-.154	.507	1.972

a. Dependent Variable: WLB

Excluded Variables^a

Model	Beta In	t	Sig.	Partial Correlation	Collinearity Statistics		
					Tolerance	VIF	Minimum Tolerance
1	S	. ^b	.	.	.000	.	.000

a. Dependent Variable: WLB

b. Predictors in the Model: (Constant), SI, WS, IE, SM, PE, H, FA

Collinearity Diagnostics^a

Model	Dimension	Eigenvalue	Condition Index	Variance Proportions							
				(Constant)	SM	FA	H	WS	PE	IE	SI
1	1	7.833	1.000	.00	.00	.00	.00	.00	.00	.00	.00
	2	.056	11.807	.01	.02	.00	.46	.01	.01	.05	.00
	3	.036	14.775	.00	.26	.02	.09	.01	.25	.00	.02
	4	.028	16.794	.04	.00	.01	.01	.06	.05	.00	.54
	5	.019	20.548	.07	.02	.23	.10	.11	.02	.21	.01
	6	.014	23.700	.02	.67	.06	.15	.00	.52	.05	.14
	7	.008	30.771	.20	.02	.43	.08	.30	.11	.31	.28
	8	.006	35.412	.66	.01	.24	.11	.51	.05	.37	.00

a. Dependent Variable: WLB