

a
pymatgen D of each run
600 K
800 K
1000 K
float64 machine epsilon (a)
52 atoms
416 atoms, 200 ps
Cl excess, 392-416 atoms
R2 limit, 10⁻⁶
R1 limit, 10⁻⁹
Relative difference
identical
b
kinisi D of each run (closest: 0.999 of its threshold)
threshold max(3 s_seed, 2 %)
3 s_seed (when below the floor)
floor 2 %
Relative difference (%)
c
kinisi standard deviation of D (closest: 0.533 of its threshold)
threshold max(3 s_seed, 10 %)
3 s_seed (when below the floor)
floor 10 %
Relative difference (%)
d
Arrhenius fit of each series
|d ln σ(300 K)| / published log half-width (left)
|d E_a| / published E_a error (right)
R3
R2-
R2
|Difference| / uncertainty
MACE-MP-0 52 atoms 50 ps
MatterSim 52 atoms 50 ps
MACE-MP-0 52 atoms 150 ps
MatterSim 52 atoms 150 ps
MACE-MP-0 416 atoms 200 ps
MatterSim 416 atoms 200 ps
Gamma, stage 2 416 atoms 200 ps
x = 0 416 atoms 150 ps
x = 0.25 408 atoms 150 ps
x = 0.50 400 atoms 150 ps
x = 0.75 392 atoms 150 ps
Universal potentials
Fine-tuned
Cl excess, MACE-MP-0
e
Force RMSE in the MACE training logs
± half the last printed digit
* logged over 20 of 22 training configurations
validation set
training set
Gamma, stage 1, epoch 89
Gamma, stage 2, epoch 116
2x2x2, stage 1, epoch 89
2x2x2, stage 2, epoch 119
MACE-MP-0, before fine-tuning
Logged - re-evaluated force RMSE (meV/Å)
f
Thermostat statistics against the July run logs
tolerance
|Difference| / tolerance
Mean T
Mean E
Drift
g
Seed scatter of kinisi D, all runs
3 s_seed = 2 %
s_seed of kinisi D (%)
MACE-MP-0
MatterSim
Gamma, stage 2
2x2x2, stage 1
kinisi posterior std / D (%)