

Type 2 Diabetes Mellitus





Table of contents

01

What Is Type 2 Diabetes?

02

What Causes Type 2 Diabetes?

03

Signs and Symptoms

04

Testing and Diagnosis

05

Treatment

06

Prognosis






01

What Is Type 2 Diabetes?

What Is Type 2 Diabetes?

- A chronic disease affecting how the body uses glucose (blood sugar).
- The body becomes resistant to insulin or does not produce enough insulin.
- Results in high blood sugar levels.



TYPE 2

Body can't use insulin properly



Can develop at
any age



Most cases **can**
be prevented

90-95%

In adults, type 2 diabetes
accounts for approximately
90-95% of all diagnosed
cases of diabetes.

nearly
5,300

Nearly 5,300 youth
diagnosed each year in
2017 and 2018

Risk factors for type 2 diabetes:



Being
overweight



Having a family
history



Being physically
inactive



Being 45 or
older


1.2 Million

People **18 years or older**
diagnosed with diabetes
in 2021

02 What Causes Type 2 Diabetes?



What Causes Type 2 Diabetes?



- **Insulin Resistance**
- **Obesity**
- **Physical Activity**

What Causes Type 2 Diabetes?



- Increasing age
- Family History/Genetics
- Unhealthy Diet

03 Signs and Symptoms





Signs and Symptoms



Frequent Urination




Excessive Thirst



Increased Hunger



Fatigue



Blurred Vision



**Slow Wound
Healing**



04 Testing and Diagnosis





Testing and Diagnosis

Fasting Blood Glucose Test

Measures blood sugar levels after not eating or drinking anything except water for at least 8 hours. It is commonly used to diagnose and monitor diabetes.

Oral Glucose Tolerance Test (OGTT)

Measures how well the body processes glucose. Blood sugar is tested before and after drinking a glucose solution, usually over a 2-hour period. It is often used to diagnose diabetes and gestational diabetes.

HbA1c Test

Measures the average blood sugar level over the past 2–3 months by checking how much glucose is attached to red blood cells. It helps diagnose diabetes and assess long-term blood sugar control.

Random Blood Sugar Test

Measures blood sugar at any time of the day, regardless of when the person last ate. It is used as a quick screening tool for diabetes, especially when symptoms are present.

Testing and Diagnosis



Fasting Blood Glucose Test

Measures blood sugar levels after not eating or drinking anything except water for at least 8 hours. It is commonly used to diagnose and monitor diabetes.



Testing and Diagnosis

Fasting Blood Glucose Test

Measures blood sugar levels after not eating or drinking anything except water for at least 8 hours. It is commonly used to diagnose and monitor diabetes.

Oral Glucose Tolerance Test (OGTT)

Measures how well the body processes glucose. Blood sugar is tested before and after drinking a glucose solution, usually over a 2-hour period. It is often used to diagnose diabetes and gestational diabetes.

HbA1c Test

Measures the average blood sugar level over the past 2–3 months by checking how much glucose is attached to red blood cells. It helps diagnose diabetes and assess long-term blood sugar control.

Random Blood Sugar Test

Measures blood sugar at any time of the day, regardless of when the person last ate. It is used as a quick screening tool for diabetes, especially when symptoms are present.



Testing and Diagnosis

HbA1c Test

Measures the average blood sugar level over the past 2–3 months by checking how much glucose is attached to red blood cells. It helps diagnose diabetes and assess long-term blood sugar control.





Testing and Diagnosis

Fasting Blood Glucose Test

Measures blood sugar levels after not eating or drinking anything except water for at least 8 hours. It is commonly used to diagnose and monitor diabetes.

Oral Glucose Tolerance Test (OGTT)

Measures how well the body processes glucose. Blood sugar is tested before and after drinking a glucose solution, usually over a 2-hour period. It is often used to diagnose diabetes and gestational diabetes.

HbA1c Test

Measures the average blood sugar level over the past 2–3 months by checking how much glucose is attached to red blood cells. It helps diagnose diabetes and assess long-term blood sugar control.

Random Blood Sugar Test

Measures blood sugar at any time of the day, regardless of when the person last ate. It is used as a quick screening tool for diabetes, especially when symptoms are present.

Testing and Diagnosis



Oral Glucose Tolerance Test (OGTT)

Measures how well the body processes glucose. Blood sugar is tested before and after drinking a glucose solution, usually over a 2-hour period. It is often used to diagnose diabetes and gestational diabetes.



Testing and Diagnosis

Fasting Blood Glucose Test

Measures blood sugar levels after not eating or drinking anything except water for at least 8 hours. It is commonly used to diagnose and monitor diabetes.

Oral Glucose Tolerance Test (OGTT)

Measures how well the body processes glucose. Blood sugar is tested before and after drinking a glucose solution, usually over a 2-hour period. It is often used to diagnose diabetes and gestational diabetes.



HbA1c Test

Measures the average blood sugar level over the past 2–3 months by checking how much glucose is attached to red blood cells. It helps diagnose diabetes and assess long-term blood sugar control.

Random Blood Sugar Test

Measures blood sugar at any time of the day, regardless of when the person last ate. It is used as a quick screening tool for diabetes, especially when symptoms are present.




Testing and Diagnosis

Random Blood Sugar Test

Measures blood sugar at any time of the day, regardless of when the person last ate. It is used as a quick screening tool for diabetes, especially when symptoms are present.

Random Blood Sugar Levels



05 Treatment



Treatment

Lifestyle Changes



❖ Healthy Diet



❖ Regular Exercise



❖ Weight Management

Treatment

Medications



❖ Metformin




❖ Other glucose-lowering



❖ Insulin (When needed)

06 Prognosis





Prognosis

- No complete cure currently.
- Can be controlled effectively.
- Early diagnosis improves outcomes.
- Good management reduces complications.

The background is a light purple color. A large, faint stethoscope is centered behind the text. Various medical icons are scattered around: a red and white pill bottle in the top left, a clipboard with a checklist in the top right, a red heart with a white cross in the bottom left, and a green pill bottle with a white cross in the bottom right. There are also several small purple plus signs and dotted lines.

Thank you