### Exercises 1.9

## Exercise 1.9.1

A factory has n suppliers that produce quantities  $x_1 
ldots x_n$  per day. The factory is connected with suppliers by a system of roads, which can be at variable capacities  $c_1 
ldots c_n$ , so that the factory is supplied daily the amount  $x = c_1 x_1 + \dots + c_n x_n$ .

- a. Given that the factory production process starts when the supply reaches the critical daily level b, write a formula for the daily factory revenue.
- b. Formulate the problem as a learning problem.

### Exercise 1.9.2

A number of finantial institutions, each having a wealth  $x_i$ , deposit amounts of money in a fund, at some adjustable rates of deposit  $w_i$ , so the money in the fund is given by  $x = x_1w_1 + \cdots + x_nw_n$ . The fund is set up to function as in the following: as long a sthe fund has less than a certain reserve fund M, the fund manager does not invest. Only the money exceeding the reserve found M is invested. Let  $k = e^{rt}$ , where r and t denote the investment rate of return and time of investment, respectively.

- a. Find the formula for the investment.
- b. Formulate the problem as a learning problem.

### Exercise 1.9.3

- a. Given a continous function  $f:[0,1]\to\mathbb{R}$ , find a linear function L(x)=ax+b with L(0)=f(0) and such that  $\frac{1}{2}\int_0^1(L(x)-f(x))^2dx$  is minimized.
- b. Given a continous function  $f:[0,1]\times[0,1]\to\mathbb{R}$ , find a linear function L(x,y)=ax+by+c with L(0,0)=f(0,0) and such that the error  $\frac{1}{2}\int_{[0,1]^2}(L(x,y)-f(x,y))^2dx$  is minimized

## Exercise 1.9.4

For any compact  $K \subset \mathbb{R}^n$  we associate the symetric matrix  $\rho_{ij} = \int_K x_i x_j dx_1 \dots dx_n$  The invertibility of the matrix  $(\rho_{ij})$  depends both on the shape of K and the dimension n.

- a. Show that if n=2 then  $\det(\rho_{ij})\neq 0$ , for any compact  $K\subset\mathbb{R}^2$ .
- b. Asume  $K = [0,1]^n$ . Show that  $\det(\rho_{ij}) \neq 0$ , for any  $n \geq 1$ .

### SOLUTIONS

#### 1.9.1 (a)

Let  $\mathbf{c} := (c_1, \dots, c_n)$  and  $\mathbf{p} := (x_1, \dots, x_n)$  the roads variable capacities and the produced quantities, respectively. Then  $x = \mathbf{c} \cdot \mathbf{p}$ . Suppose the cost of product per item is k, so if the production starts after the critical daily level b is meet. This is, x - b > 0. It is clear that the revenue  $L_r$  will be given by the formula:

$$L_r(\mathbf{p}; \mathbf{c}, b) = \begin{cases} k(\mathbf{c} \cdot \mathbf{p} - b), & \text{if } \mathbf{c} \cdot \mathbf{p} - b > 0 \\ 0, & \text{otherwise.} \end{cases}$$

### 1.9.1 (b)

The learning problem can be stated like this: "Given a vector of road variable capacities  $\mathbf{c}$  and a daily critical level b, provided that the production of the n factories is expressed by the vector  $\mathbf{p}$ . The goal is find a vector  $\mathbf{p}^*$  and a scalar  $b^*$  such that the ideal revenue  $r(\mathbf{p}) = \mathbf{c} \cdot \mathbf{p}$  is close that provided by the data  $L_r(\mathbf{p}; \mathbf{c}, b)$  (obtained in 1.9.1 (a))". In other words, the pair  $(\mathbf{c}^*, b^*)$  minimizes the distance (in the  $\mathcal{L}_2$  sense) between  $r(\mathbf{p})$  and  $L_r(\mathbf{p}; \mathbf{c}, b)$ . In symbols:

$$(\mathbf{c}^{\star}, b^{\star}) = \underset{\mathbf{c} \in \mathbb{R}^n, \ p \in \mathbb{R}}{\arg \min} \int_{\mathcal{K}} (r(\mathbf{p}) - L_r(\mathbf{p}; \mathbf{c}, b))^2 d\mathbf{p}.$$

### 1.9.2 (a)

If  $\mathbf{w} := (w_1 \dots w_n)$  encodes the adjustable rates of deposit corresponding to each of the n finantial institutions and  $\mathbf{x} := (x_1 \dots x_n)$  the wealth of each of the n institutions, the money in the fund is expressed by  $x = \mathbf{w} \cdot \mathbf{x}$ . It's known that the fund is set to function if the revenue exceds a given capital M and that the investment grows proportional to  $e^{rt}$  (profit per investment), then it is clear that the investment is given by the formula:

$$L_I(\mathbf{x}; \mathbf{w}, M) = \begin{cases} e^{rt}(\mathbf{w} \cdot \mathbf{x} - M), & \text{if } \mathbf{w} \cdot \mathbf{x} > M \\ 0, & \text{otherwise.} \end{cases}$$

#### 1.9.2 (b)

Let  $I(\mathbf{x}) = x = \mathbf{w} \cdot \mathbf{x}$  be the ideal investment,  $L_I(\mathbf{x}; \mathbf{w}, M)$  given in 1.9.2 (b). Then the learning problem can be stated as follows: "Given the vector of wealth of the n institutions  $\mathbf{x}$ , the vector of adjustable rates of deposit  $\mathbf{w}$  and that inversions are placed in the fund if the capital exceeds the quantity M. It is needed to find a tuple  $(\mathbf{w}^*, M)$  that minimizes the distance (in the  $\mathcal{L}_2$  sense) between the functions  $I(\mathbf{x})$  and  $L_I(\mathbf{x}; \mathbf{w}, M)$ "

In mathematical terms this means that:

$$(\mathbf{w}^{\star}, M^{\star}) = \underset{\mathbf{w} \in \mathbb{R}^n, \ M \in \mathbb{R}}{\arg \min} \int_{\mathcal{K}} (I(\mathbf{x}) - L_I(\mathbf{x}; \mathbf{c}, M))^2 d\mathbf{x}.$$

#### 1.9.3 (a)

It follows from the constrain f(0) = L(0) that b = f(0), that said, the task is now to find an  $a^*$  such that  $a^* = \underset{a \in \mathbb{R}}{\arg \min} \|f(x) - L(x)\|_{\mathcal{L}_2([0,1])}^2$ . Now, let  $C(a) := \|f(x) - L(x)\|_{\mathcal{L}_2([0,1])}^2$ , then C(a) has the most explicit form:

$$C(a) = \int_{[0,1]} (f(x) - ax - f(0))^2 dx.$$

The extremizing  $a^*$  can be found by computing critical points and applying the second derivative test to determine the nature of the critical point. Then one has:

 $\frac{d}{da}C(a) = \int_{[0,1]} -2(f(x)-ax-f(0))xdx = a\int_{[0,1]} 2x^2dx + \int_{[0,1]} 2(f(0)-f(x))xdx.$  To find the critical point  $a^*$  we solve the equation  $\frac{d}{da}C(a) = 0$ . Solving for a one finds that such extrema is meet at:

$$a^* = \frac{\displaystyle\int_{[0,1]} (f(x)-f(0))xdx}{\displaystyle\int_{[0,1]} x^2dx}$$
. Now, lets compute the second derivative to determine the nature

of this critical point. A straight-forward calculation for finding the second derivative yields:

 $\frac{d^2}{dx^2}C(a) = 2\int_{[0,1]} x^2 dx. \text{ Because the even function } 2x^2 \text{ satisfies } 2x^2 > 0, \forall x \in [0,1] \text{ and the interval of integration is not symetric, is clear that } \frac{d^2}{dx^2}C(a) = 2\int_{[0,1]} x^2 dx > 0, \forall a \in \mathbb{R}. \text{ From this follows that } a^* \text{ is a minimum. Then, } L(x) = \frac{\int_{[0,1]} (f(t) - f(0))tdt}{\int_{[0,1]} t^2 dt} x + f(0).$ 

# 1.9.3 (b)