

## Exercises 3.15

### Exercise 3.15.1

Let  $p$ ,  $p_i$ ,  $q$ ,  $q_i$  be density functions on  $\mathbb{R}$  and  $\alpha \in \mathbb{R}$ . Show that the cross-entropy satisfies the following properties:

- a.  $S(p_1 + p_2, q) = S(p_1, q) + S(p_2, q)$ ;
- b.  $S(\alpha p, q) = \alpha S(p, q) = S(p, q^\alpha)$ ;
- c.  $S(p, q_1 q_2) = S(p, q_1) + S(p, q_2)$ .

### Exercise 3.15.2

Show that the cross entropy satisfies the following inequality

$$S(p, q) \geq 1 - \int p(x)q(x)dx$$

### Exercise 3.15.3

Let  $p$  a fixed density. Show that the symmetric relative entropy

$$D_{KL}(p||q) + D_{KL}(q||p)$$

reaches its minimum for  $p = q$ , and the minimum is equal to zero.

### Exercise 3.15.4

Consider two exponential densities,  $p_1 = \xi^1 e^{\xi^1 x}$  and  $p_2 = \xi^2 e^{\xi^2 x}$ ,  $x \geq 0$ .

- a. Show that  $D_{KL}(p_1||p_2) = \frac{\xi^2}{\xi^1} - \ln \xi^2 \xi^1 - 1$ .
- b. Verify  $D_{KL}(p_1||p_2) \neq D_{KL}(p_2||p_1)$ .
- c. Show that the triangle inequality doesn't hold for three arbitrary densities.

### Exercise 3.15.5

Let  $X$  be a discrete random variable. Show the inequality

$$H(X) \geq 0.$$

### Exercise 3.15.7

We assume the target variable  $Z$  is  $\mathcal{E}$ -measurable. What is mean squared error function in this case?

### Exercise 3.15.8

Assume that a neural network has an input-output function  $f_{w,b}$  linear in  $w$  and  $b$ . Show that the cost function (3.3.1) reaches its minimum for a unique pair  $(w^*, b^*)$ , which can be computed explicitly.

### Exercise 3.15.9

Show that the Shannon entropy can be retrieved from the Reyni entropy as

$$H(p) = \lim_{\alpha \rightarrow 1} H_\alpha(x).$$

### Exercise 3.15.10

Let  $\phi_\sigma(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{x^2}{2\sigma^2}}$ . Consider the convolution operation  $(f * g)(x) := \int f(t)g(x-t)dt$ .

- Show that  $\phi_\sigma * \phi_\sigma = \phi_{\sigma\sqrt{2}}$ ;
- Find  $\phi_\sigma * \phi_{\sigma'}$  in the case  $\sigma \neq \sigma'$ .

### Exercise 3.15.11

Consider two probability densities,  $p(x)$  and  $q(x)$ . The Cauchy-Schwartz divergence is defined by

$$D_{CS}(p, q) := -\ln\left(\frac{\int p(x)q(x)dx}{\sqrt{\int p(x)^2 dx} \sqrt{\int q(x)^2 dx}}\right).$$

Show the following:

- $D_{CS}(p, q) = 0$  if and only if  $p = q$ ;
- $D_{CS}(p, q) \geq 0$ ;
- $D_{CS}(p, q) = D_{CS}(q, p)$ ;
- $D_{CS}(p, q) = -\ln \int pq dx - \frac{1}{2}H_2(p) - \frac{1}{2}H_2(q)$ , where  $H_2(\cdot)$  denotes the quadratic Reyni entropy.

### Exercise 3.15.12

- Show that for any function  $f \in L^1[0, 1]$  we have the inequality  $\|\tanh(f)\|_1 \leq \|f\|_1$ .
- Show that for any function  $f \in L^2[0, 1]$  we have the inequality  $\|\tanh\|_2 \leq \|f\|_2$ .

### Exercise 3.15.13

Consider two distributions on the sample space  $\mathcal{X} = \{x_1, x_2\}$  given by

$$p = \begin{pmatrix} x_1 & x_2 \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}, \quad q = \begin{pmatrix} x_1 & x_2 \\ \frac{1}{2} & \frac{2}{3} \end{pmatrix}$$

Consider the function  $\phi : \mathcal{X} \rightarrow \mathbb{R}^2$  defined by  $\phi(x_1) = (0, 1)$   $\phi(x_2) = (1, 0)$ . Find the maximum mean discrepancy between  $p$  and  $q$ .

## SOLUTIONS

### 3.15.1 (a)

At vero eos et accusamus et iusto odio dignissimos ducimus qui blanditiis praesentium voluptatum deleniti atque corrupti quos dolores et quas molestias excepturi sint occaecati cupiditate non provident, similique sunt in culpa qui officia deserunt mollitia animi, id est laborum et dolorum fuga.

### x.y.2 (a)

Et harum quidem rerum facilis est et expedita distinctio. Nam libero tempore, cum soluta nobis est eligendi optio cumque nihil impedit quo minus id quod maxime placeat facere possimus, omnis voluptas assumenda est, omnis dolor repellendus.

*Proof.*  $a = a$

□