Best subsets ideal model - main effects only

The LOGISTIC Procedure

Model Information

Data Set WORK.TIMESINCEVAX

Response Variable breakthrough breakthrough

Number of Response Levels

Model binary logit Optimization Technique Fisher's scoring

> Number of Observations Read 3084648 3084648 Number of Observations Used

Response Profile

Total		Ordered
Frequency	breakthrough	Value
	-	
29677	1	1
3054971	0	2

Probability modeled is breakthrough=1.

Model Convergence Status

Convergence criterion (GCONV=1E-8) satisfied.

Model Fit Statistics

		Intercept
	Intercept	and
Criterion	Only	Covariates
AIC	334698.88	9888.961
SC	334711.82	9927.787

-2 Log L 334696.88 9882.961

Testing Global Null Hypothesis: BETA=0

Test	Chi-Square	DF	Pr > ChiSq
Likelihood Ratio	324813.922	2	<.0001
Score	1677433.11	2	<.0001
Wald	2998.5594	2	<.0001

Analysis of Maximum Likelihood Estimates

			Standard	Wald	
Parameter	DF	Estimate	Error	Chi-Square	Pr > ChiSq
Intercept	1	-6.6629	0.0873	5826.4987	<.0001
followup_time	1	-4.3909	0.0810	2940.2357	<.0001
time_since_vax	1	132.9	2.4527	2935.7191	<.0001

Best subsets ideal model - main effects only

The LOGISTIC Procedure

Association of Predicted Probabilities and Observed Responses

Percent	Concordant	99.7	Somers' D	0.995
Percent	Discordant	0.3	Gamma	0.995
Percent	Tied	0.0	Tau-a	0.019
Pairs		90662374367	С	0.997

Odds Ratio Estimates and Wald Confidence Intervals

Odds Ratio	Estimate	95% Confider	nce Limits
followup_time	0.012	0.011	0.015
time since vax	>999.999	>999.999	>999.999

Odds Ratio Estimates and Profile-Likelihood Confidence Intervals

Effect	Unit	Estimate	95% Confid	ence Limits
followup_time	1.0000	0.012	0.011	0.014
time_since_vax	1.0000	>999.999	>999.999	>999.999