



EBM Calculator

Formulas for Effect Estimates

	Outcome	No Outcome	Total
(+)	<i>a</i>	<i>b</i>	<i>a + b</i>
(-)	<i>c</i>	<i>d</i>	<i>c + d</i>
Total	<i>a + c</i>	<i>b + d</i>	<i>a + b + c + d</i>

$z = 1.959964$ for 95% CI

Experimental Event Rate

$$\text{EER} = \frac{a}{a + b}$$

Control Event Rate

$$\text{CER} = \frac{c}{c + d}$$

Absolute Risk Reduction

$$ARR = CER - EER$$

Confidence Interval (Wilson score bounds with error propagation)

$$\text{Lower Bound: } ARR - z \cdot \sqrt{\frac{u_2(1 - u_2)}{r_1} + \frac{w_1(1 - w_1)}{r_2}}$$

$$\text{Upper Bound: } ARR + z \cdot \sqrt{\frac{u_1(1 - u_1)}{r_2} + \frac{w_2(1 - w_2)}{r_1}}$$

where...

$$r_1 = a + b, \quad r_2 = c + d$$

$$u_1 = \frac{2c + z^2 + z \sqrt{\frac{4(c \cdot d)}{r_2} + z^2}}{2r_2 + 2z^2}$$

$$u_2 = \frac{2a + z^2 + z \sqrt{\frac{4(a \cdot b)}{r_1} + z^2}}{2r_1 + 2z^2}$$

$$w_1 = \frac{2c + z^2 - z \sqrt{\frac{4(c \cdot d)}{r_2} + z^2}}{2r_2 + 2z^2}$$

$$w_2 = \frac{2a + z^2 - z \sqrt{\frac{4(a \cdot b)}{r_1} + z^2}}{2r_1 + 2z^2}$$

Risk Ratio

$$RR = \frac{EER}{CER}$$

Confidence Interval (Log-normal, Zhou)

$$\text{Lower Bound: } \exp \left(\ln(RR) - z \cdot \sqrt{\left(\frac{1}{a} - \frac{1}{a+b} \right) + \left(\frac{1}{c} - \frac{1}{c+d} \right)} \right)$$

$$\text{Upper Bound: } \exp \left(\ln(RR) + z \cdot \sqrt{\left(\frac{1}{a} - \frac{1}{a+b} \right) + \left(\frac{1}{c} - \frac{1}{c+d} \right)} \right)$$

Relative Risk Reduction

$$RRR = 1 - RR$$

Confidence Interval (Derived from Log-normal RR)

$$\text{Lower Bound: } 1 - \exp \left(\ln(RR) + z \cdot \sqrt{\left(\frac{1}{a} - \frac{1}{a+b} \right) + \left(\frac{1}{c} - \frac{1}{c+d} \right)} \right)$$

$$\text{Upper Bound: } 1 - \exp \left(\ln(RR) - z \cdot \sqrt{\left(\frac{1}{a} - \frac{1}{a+b} \right) + \left(\frac{1}{c} - \frac{1}{c+d} \right)} \right)$$

Odds Ratio

$$OR = \frac{ad}{cb}$$

Confidence Interval (Log-normal)

$$\text{Lower Bound: } \exp \left(\ln(OR) - z \cdot \sqrt{\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}} \right)$$

$$\text{Upper Bound: } \exp \left(\ln(OR) + z \cdot \sqrt{\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}} \right)$$

Formulas for Diagnostic Tests

	Disease	No Disease	Total
(+)	a	b	$a + b$
(-)	c	d	$c + d$
Total	$a + c$	$b + d$	$a + b + c + d$

$z = 1.959964$ for 95% CI

Sensitivity

$$\text{Sensitivity} = \frac{a}{a + c}$$

Confidence Interval (Wilson Score)

$$\text{Lower Bound: } \frac{(2 \cdot a) + z^2 - z \cdot \sqrt{\left(\frac{4ac}{a+c}\right) + z^2}}{2 \cdot (a + c) + 2z^2}$$

$$\text{Upper Bound: } \frac{(2 \cdot a) + z^2 + z \cdot \sqrt{\left(\frac{4ac}{a+c}\right) + z^2}}{2 \cdot (a + c) + 2z^2}$$

Specificity

$$\text{Specificity} = \frac{d}{b + d}$$

Confidence Interval (Wilson Score)

$$\text{Lower Bound: } \frac{(2 \cdot d) + z^2 - z \cdot \sqrt{\left(\frac{4db}{b+d}\right) + z^2}}{2 \cdot (b + d) + 2z^2}$$

$$\text{Upper Bound: } \frac{(2 \cdot d) + z^2 + z \cdot \sqrt{\left(\frac{4db}{b+d}\right) + z^2}}{2 \cdot (b + d) + 2z^2}$$

Positive Likelihood Ratio

$$LR(+) = \frac{\text{Sensitivity}}{1 - \text{Specificity}} = \frac{a/(a+c)}{b/(b+d)}$$

Confidence Interval (Log-normal, Zhou)

$$\text{Lower Bound: } \exp \left(\ln \left(\frac{(b+d) \cdot a}{(a+c) \cdot b} \right) - z \cdot \sqrt{\left(\frac{c}{a \cdot (a+c)} \right) + \left(\frac{d}{b \cdot (b+d)} \right)} \right)$$

$$\text{Upper Bound: } \exp \left(\ln \left(\frac{(b+d) \cdot a}{(a+c) \cdot b} \right) + z \cdot \sqrt{\left(\frac{c}{a \cdot (a+c)} \right) + \left(\frac{d}{b \cdot (b+d)} \right)} \right)$$

Negative Likelihood Ratio

$$LR(-) = \frac{1 - \text{Sensitivity}}{\text{Specificity}} = \frac{c/(a+c)}{d/(b+d)}$$

Confidence Interval (Log-normal, Zhou)

$$\text{Lower Bound: } \exp \left(\ln \left(\frac{(b+d) \cdot c}{(a+c) \cdot d} \right) - z \cdot \sqrt{\left(\frac{a}{c \cdot (a+c)} \right) + \left(\frac{b}{d \cdot (b+d)} \right)} \right)$$

$$\text{Upper Bound: } \exp \left(\ln \left(\frac{(b+d) \cdot c}{(a+c) \cdot d} \right) + z \cdot \sqrt{\left(\frac{a}{c \cdot (a+c)} \right) + \left(\frac{b}{d \cdot (b+d)} \right)} \right)$$

Positive Predictive Value

$$PPV = \frac{a}{a+b}$$

Confidence Interval (Wilson Score)

$$\text{Lower Bound: } \frac{(2 \cdot a) + z^2 - z \cdot \sqrt{\left(\frac{4ab}{a+b} \right) + z^2}}{2 \cdot (a+b) + 2z^2}$$

$$\text{Upper Bound: } \frac{(2 \cdot a) + z^2 + z \cdot \sqrt{\left(\frac{4ab}{a+b} \right) + z^2}}{2 \cdot (a+b) + 2z^2}$$

Negative Predictive Value

$$\text{NPV} = \frac{d}{c + d}$$

Confidence Interval (Wilson Score)

$$\text{Lower Bound: } \frac{(2 \cdot d) + z^2 - z \cdot \sqrt{\left(\frac{4dc}{c+d}\right) + z^2}}{2 \cdot (c + d) + 2z^2}$$

$$\text{Upper Bound: } \frac{(2 \cdot d) + z^2 + z \cdot \sqrt{\left(\frac{4dc}{c+d}\right) + z^2}}{2 \cdot (c + d) + 2z^2}$$

Formulas for Post-Test Probability

Post-Test Probability

$$\text{PostProb} = \frac{\text{PostOdds}}{\text{PostOdds} + 1}$$

Post-Test Odds

$$\text{PostOdds} = \text{PreOdds} \times \text{LR}$$

Pre-Test Odds

$$\text{PreOdds} = \frac{\text{PreProb}}{1 - \text{PreProb}}$$

Conversions

$$\text{Odds} = \frac{\text{Probability}}{1 - \text{Probability}}$$

$$\text{Probability} = \frac{\text{Odds}}{\text{Odds} + 1}$$

References

1. Altman DG. Confidence intervals for the number needed to treat. *BMJ*. 1998 Nov 7;317(7168):1309–1312.
doi: 10.1136/bmj.317.7168.1309. PMID: 9804726; PMCID: PMC1114210.
<https://pubmed.ncbi.nlm.nih.gov/9804726/>
2. Bender R. Calculating confidence intervals for the number needed to treat. *Control Clin Trials*. 2001 Apr;22(2):102–110.
doi: 10.1016/s0197-2456(00)00134-3. PMID: 11306148.
<https://pubmed.ncbi.nlm.nih.gov/11306148/>
3. Fleiss JL, Levin B, Paik MC. (2003). *Statistical Methods for Rates and Proportions* (3rd ed.). Wiley.
ISBN: 978-0-471-52629-2.
<https://www.wiley.com/en-us/Statistical+Methods+for+Rates+and+Proportions%2C+3rd+Edition-p-9780471526292>
4. Mercaldo ND, Lau KF, Zhou XH. Confidence intervals for predictive values with an emphasis to case-control studies. *Stat Med*. 2007 May 10;26(10):2170–2183.
doi: 10.1002/sim.2677. PMID: 16927452.
<https://pubmed.ncbi.nlm.nih.gov/16927452/>
5. Julious SA. Two-sided confidence intervals for the single proportion: comparison of seven methods by Robert G. Newcombe, *Stat Med*. 2005 Nov 15;24(21):3383–3384.
doi: 10.1002/sim.2164. PMID: 16206245.
<https://pubmed.ncbi.nlm.nih.gov/16206245/>
6. Zhou XH, Obuchowski NA, McClish DK. (2011). *Statistical Methods in Diagnostic Medicine* (2nd ed.). Wiley.
Print ISBN: 9780470183144. Online ISBN: 9780470906514.
<https://onlinelibrary.wiley.com/doi/book/10.1002/9780470906514>