

Introduction to dtametaTMB

This vignette introduces the main functionality of the **dtametaTMB** package for meta-analysis of diagnostic test accuracy (DTA) studies. We demonstrate how to fit three commonly used models:

- the Reitsma bivariate model
- the Rutter and Gatsonis (HSROC) model
- the Hoyer model for multiple thresholds

using real datasets included in the package.

Note: Rendering forest plots via **forest** may take longer in an interactive R session due to the underlying grid graphics. Performance is typically faster when knitting the vignette (e.g. via RMarkdown or Quarto).

How do I load the package?

```
library(dtametaTMB)
```

Which model should I use?

- **Reitsma model:** single threshold per study: use when studies apply similar thresholds.
- **HSROC model:** single threshold per study: use when thresholds vary between studies or in small and extreme data sets
- **Hoyer model:** multiple thresholds per study

Validation

The implementations reproduce benchmark results from the Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy (including parameter estimates, standard errors, and graphical outputs) for standard datasets such as the Anti-CCP and RF examples.

How do I perform a diagnostic test meta-analysis using the bivariate binomial-model of Reitsma?

Unlike the **mada** package, **dtametaTMB** uses the binomial likelihood to model the within study distribution as recommended by The Cochrane Handbook.

For sensitivity in study i , we have the number of diseased individuals testing positive: $y_{Ai} \sim \mathcal{B}(n_{Ai}, \pi_{Ai})$.

Similarly for specificity, we have the number of non-diseased individuals testing negative: $y_{Bi} \sim \mathcal{B}(n_{Bi}, \pi_{Bi})$.

At the study level, we have

$$\begin{pmatrix} \text{logit}(\pi_{Ai}) \\ \text{logit}(\pi_{Bi}) \end{pmatrix} = \begin{pmatrix} \mu_{Ai} \\ \mu_{Bi} \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} \mu_A \\ \mu_B \end{pmatrix}, \Sigma \right), \quad \text{with} \quad \Sigma = \begin{pmatrix} \sigma_A^2 & \sigma_{AB} \\ \sigma_{AB} & \sigma_B^2 \end{pmatrix}.$$

```
data("anticcp")
reitsma <- fitReitsma(data=anticcp,
                      TP=TP,
                      FP=FP,
                      FN=FN,
```

```

      TN=TN,
      study=study)


reitsma
#>
#> Reitsma Bivariate Random-Effects Model
#> -----
#> Number of studies: 37
#> Model fit: Converged (positive definite Hessian)
#>
#> Key parameters (logit scale):
#>   mu_A (sens): 0.653
#>   mu_B (spec): 3.109
summary(reitsma)
#> $estimates
#>           Estimate Std_Error
#> mu_A.sens      0.6534029 0.1274866
#> mu_B.spec      3.1089906 0.1458847
#> sigma2_A.sens  0.5426397 0.1463478
#> sigma2_B.spec  0.5717019 0.1873211
#> sigma_AB      -0.2704148 0.1198520
#>
#> $sensspec
#>           Estimate conflevel  CI_Lower  CI_Upper
#> sens 0.6577769      0.95 0.5995364 0.7116214
#> spec 0.9572621      0.95 0.9439094 0.9675454
#>
#> $RutterGatsonis_recovered
#>           Lambda      Theta      beta sigma2_alpha sigma2_theta
#> Estimate (recovered) 3.730684 -1.203361 0.02608605 0.573133 0.4136981

```

How do I get a summary plot of the Reitsma model?


```
plot(reitsma)
```

Diagnostic Test Accuracy Meta-Analysis



```
plot(reitsma,scale=0.01)
```


Diagnostic Test Accuracy Meta-Analysis



How do I get a coupled forest plot?

Note: Rendering forest plots may take longer in an interactive R session due to the underlying grid graphics. Performance is typically faster when knitting the vignette (e.g. via RMarkdown or Quarto).

```
forest(reitsma)
```



How do I perform a diagnostic test meta-analysis using the Rutter and Gatsonis model?

The number of diseased individuals from study i who test positive is denoted by $y_{i1} \sim \mathcal{B}(n_{i1}, \pi_{i1})$.

Similarly, the number of non-diseased individuals who test positive is $y_{i2} \sim \mathcal{B}(n_{i2}, \pi_{i2})$.

At the study level, we have

$$\text{logit}(\pi_{ij}) = (\theta_i + \alpha_i x_{ij}) \exp(-\beta x_{ij}), \quad \alpha_i \sim \mathcal{N}(\Lambda, \sigma_\alpha^2), \quad \theta_i \sim \mathcal{N}(\Theta, \sigma_\theta^2),$$

where

$$x_{ij} = \begin{cases} -0.5 & \text{for non-diseased individuals,} \\ 0.5 & \text{for diseased individuals.} \end{cases}$$

```
data("RF")
ruttermatsonis <- fitRutterGatsonis(data=RF,
                                     TP=TP,
                                     FP=FP,
                                     FN=FN,
```


```
TN=TN,  
study=study)
```

```
ruttergatsonis  
#>  
#> Rutter & Gatsonis HSROC Model  
#> -----  
#> Number of studies: 50  
#> Model fit: Converged  
#>  
#> Parameters (HSROC scale):  
#>   Lambda: 2.602  
#>   Theta : -0.437  
#>   Beta  : 0.227  
summary(ruttergatsonis)  
#> $estimates  
#>           Estimate Std. Error  
#> Lambda      2.6015985 0.18616691  
#> Theta      -0.4370167 0.14685177  
#> beta        0.2266849 0.16236992  
#> sigma2_alpha 1.3014057 0.30455918  
#> sigma2_theta 0.5422965 0.12366256  
#> logitsens    0.8286883 0.15539440  
#> sens         0.6960775 0.03287424  
#>  
#> $sensspec  
#>      spec conflevel logitsens Std_Error CI_Lower CI_Upper      Sens  
#> 1 0.8669545      0.95 0.8286883 0.1553944 0.5241209 1.133256 0.6960775  
#>   SensCI_Lower SensCI_Upper  
#> 1    0.6281109    0.7564392  
#>  
#> $Reitsma_recovered  
#>           mu_A.sens mu_B.spec sigma2_A.sens sigma2_B.spec  sigma_AB  
#> Estimate (recovered) 0.7712238 1.946381    0.6916646    1.088408 -0.2169451
```

How do I get a summary plot of the Rutter and Gatsonis model?


```
plot(ruttergatsonis)
```

Diagnostic Test Accuracy Meta-Analysis



```
plot(ruttergatsonis,size="se",scale=0.01)
```


Diagnostic Test Accuracy Meta-Analysis



How do I get a coupled forest plot?

Note: Rendering forest plots may take longer in an interactive R session due to the underlying grid graphics. Performance is typically faster when knitting the vignette (e.g. via RMarkdown or Quarto).

```
forest(ruttergatsonis)
```

How do I perform a diagnostic test accuracy meta-analysis with multiple thresholds per study using the Hoyer method?

The Hoyer model assumes that the diagnostic test measurement arises from a latent continuous variable Y_j , with $j = 0$ for the non-diseased individuals and $j = 1$ for the diseased, which follows a parametric survival distribution. The observed test results correspond to interval-censored observations of this latent variable due to the use of thresholds.

Which distributions does dtametaTMB support?

The latent variable Y_j is assumed to follow a location-scale family:

$$Y_j \sim F(\mu_j, \phi_j),$$

where:

- μ_j is a study-specific location parameter
- ϕ_j is a scale parameter

Common choices for F include:

Weibull distribution

$$f(y; \mu_j, \phi_j) = \mu_j \phi_j y^{\phi_j - 1} \exp(-\mu_j y^{\phi_j})$$

Log-normal distribution

$$f(y; \mu_j, \phi_j) = \frac{1}{\phi_j y} \phi \left(\frac{\log(y) - \mu_j}{\phi_j} \right)$$

Log-logistic distribution

$$f(y; \mu_j, \phi_j) = \frac{\mu_j \phi_j y^{\phi_j - 1}}{(1 + \mu_j y^{\phi_j})^2}$$

Random effects structure

To account for between-study heterogeneity and the correlation between diseased and non-diseased groups, we introduce study-specific random effects:

$$\log(y_0) = \beta_0 + \varepsilon_0 + u_{0i}, \quad \log(y_1) = \beta_1 + \varepsilon_1 + u_{1i},$$

with

$$\begin{pmatrix} u_{0i} \\ u_{1i} \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & \rho \sigma_0 \sigma_1 \\ \rho \sigma_0 \sigma_1 & \sigma_1^2 \end{pmatrix} \right).$$

Censoring mechanism

Because only thresholded outcomes (TN, TP, FN, FP) are observed, the latent variable Y_j is subject to censoring.

- Left censoring: $(-\infty, y_k^R]$
- Interval censoring between adjacent thresholds
- Right censoring: $(y_k^L, +\infty)$

where y_k^L and y_k^R denote finite threshold values.

How do I compute sensitivities and specificities?

We can model the ROC curve by predicting sensitivities and specificities (1 - false positive rates) at several threshold values y .

Weibull distribution

$$S_1(y) = \exp(-(y \exp(-\beta_1))^{1/\lambda_1}), \quad \text{with } \beta_1 = -\log(\mu_1), \quad \lambda_1 = 1/\phi_1,$$

$$S_0(y) = \exp(-(y \exp(-\beta_0))^{1/\lambda_0}), \quad \text{with } \beta_0 = -\log(\mu_0), \quad \lambda_0 = 1/\phi_0.$$

Log-normal distribution

$$S_1(y) = 1 - \Phi\left(\frac{1}{\lambda_1}(\log(y) - \beta_1)\right), \quad \text{with } \beta_1 = \mu_1, \quad \lambda_1 = \phi_1,$$

$$S_0(y) = 1 - \Phi\left(\frac{1}{\lambda_0}(\log(y) - \beta_0)\right), \quad \text{with } \beta_0 = \mu_0, \quad \lambda_0 = \phi_0.$$

Log-logistic distribution

$$S_1(y) = 1 / \left(1 + \exp(-\beta_1/\lambda_1) y^{1/\lambda_1}\right), \quad \text{with } \beta_1 = -\phi^{-1} \log(\mu_1), \quad \lambda_1 = 1/\phi_1,$$

$$S_0(y) = 1 / \left(1 + \exp(-\beta_0/\lambda_0) y^{1/\lambda_0}\right), \quad \text{with } \beta_0 = -\phi^{-1} \log(\mu_0), \quad \lambda_0 = 1/\phi_0.$$

How do we interpret test direction? For tests where larger values indicate disease (`testdirection = "greater"`):

- $S_1(y)$ corresponds to **sensitivity** at threshold y
- $S_0(y)$ corresponds to the **false positive rate (FPR) = 1 - specificity**

For tests where smaller values indicate disease (`testdirection = "less"`):

- $S_1(y)$ corresponds to the **false negative rate (FNR) = 1 - sensitivity**
- $S_0(y)$ corresponds to **specificity**

In a nutshell

By modelling the diagnostic marker as a latent continuous outcome, the Hoyer model:

- naturally accommodates **multiple thresholds per study**
- provides a unified framework for **interval-censored diagnostic data**
- preserves the correlation between sensitivity and specificity

The following parameters have to be estimated: β_0 , β_1 , λ_0 , λ_1 , σ_0^2 , σ_1^2 , and ρ , where λ_0 and λ_1 are transformations of ϕ_0 and ϕ_1 .

To obtain initial values, it is necessary to specify a smallest (« $\min y$) and largest (» $\max y$) value across all studies, essentially replacing negative and positive infinity for the left and right censored intervals.

How do I fit the Hoyer model?

```
data("diabetes")
head(diabetes)
#>      study threshold TP   TN   FP  FN   D   H originally_published
#> 1 Study 01         5.9 574 1389  682 262 836 2071                1
#> 2 Study 02         5.0 617 1005  7735  18 635 8740                0
#> 3 Study 02         5.1 607 1617  7123  28 635 8740                0
#> 4 Study 02         5.2 600 2438  6302  35 635 8740                0
#> 5 Study 02         5.3 581 3409  5331  54 635 8740                0
#> 6 Study 02         5.4 563 4422  4318  72 635 8740                0
summary(diabetes$threshold)
#>      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
#>   3.900   5.500   5.800   5.837   6.100   7.600
```

The argument `eval_threshold` specifies the values at which sensitivities and specificities are evaluated after model fitting.

```

hoyer <- fitHoyer(data=diabetes,
                  TP=TP,
                  FP=FP,
                  FN=FN,
                  TN=TN,
                  threshold=threshold,
                  study=study,
                  dist="loglogistic",
                  testdirection="greater",
                  eval_threshold=c(5.0,5.5,6.0,6.5,7.0,7.5),
                  smallest=2,
                  largest=10)

```

```


hoyer
#>
#> Hoyer AFT Model Fit
#> -----
#> Distribution : Loglogistic
#> Number of studies : 38
#> Test direction : greater
#> Optimization : converged
#> -2 negative Log Likelihood : 209322
#>
#> Use summary() for parameter estimates.
summary(hoyer)
#> $sdreport2
#>
#>           Estimate   Std. Error
#> beta0      1.705691148 0.0095630146
#> lambda0    0.039502004 0.0001857265
#> su0        0.058648679 0.0067784870
#> beta1      1.825337107 0.0145652403
#> lambda1    0.056398479 0.0009244722
#> su1        0.088320631 0.0109564611
#> rho        0.863032573 0.0454410883
#> covu0u1    0.004470412 0.0011417200
#> logitSurv1 3.828103143 0.2635035151
#> logitSurv1 2.138160760 0.2593347627
#> logitSurv1 0.595364241 0.2580790576
#> logitSurv1 -0.823870980 0.2591073990
#> logitSurv1 -2.137877556 0.2619046729
#> logitSurv1 -3.361188412 0.2660542694
#> logitSurv0 2.436667140 0.2426039991
#> logitSurv0 0.023873625 0.2420917690
#> logitSurv0 -2.178834277 0.2420878132
#> logitSurv0 -4.205129122 0.2424751080
#> logitSurv0 -6.081185122 0.2431661379
#> logitSurv0 -7.827751473 0.2440943827
#>
#> $sensspec
#>   threshold conflevel      Sens SensCI_Lower SensCI_Upper      Spec
#> 1      5.0      0.95 0.97871219  0.96482617  0.98718904 0.08041904
#> 2      5.5      0.95 0.89455725  0.83615237  0.93379170 0.49403188
#> 3      6.0      0.95 0.64459500  0.52236970  0.75048299 0.89833265
#> 4      6.5      0.95 0.30494258  0.20887630  0.42164362 0.98530044

```


```
#> 5      7.0      0.95 0.10546947 0.06591501 0.16457780 0.99771974
#> 6      7.5      0.95 0.03353069 0.02018063 0.05521454 0.99960164
#>   SpecCI_Lower SpecCI_Upper
#> 1   0.0515556   0.1233403
#> 2   0.3779250   0.6107860
#> 3   0.8461021   0.9342152
#> 4   0.9765667   0.9908097
#> 5   0.9963326   0.9985830
#> 6   0.9993574   0.9997531
```

How do we get a summary ROC plot?

```
plot(hoyer, main="Diagnostic Test Accuracy Meta-Analysis\nLoglogistic Model")
```




```
plot(hoyer,size="sampsize",scale=0.025,main="Diagnostic Test Accuracy Meta-Analysis\nLoglogistic Model")
```



```
plot(hoyer,size="se",scale=0.025,main="Diagnostic Test Accuracy Meta-Analysis\nLoglogistic Model")
```

Diagnostic Test Accuracy Meta-Analysis Loglogistic Model



How do I get a coupled forest plot?

Note: Rendering forest plots may take longer in an interactive R session due to the underlying grid graphics. Performance is typically faster when knitting the vignette (e.g. via RMarkdown or Quarto).

```
forest(hoyer)
```


How does testdirection="less" work?

We use the synthetic anaemia data set. Smaller values in haemoglobin indicate disease.

```
data("anaemia")
summary(anaemia$threshold)
#>      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
#>      7.00   8.00   10.00   10.33   13.00   14.00
```


```
less <- fitHoyer(data=anaemia,
                 TP=TP,
                 FP=FP,
                 FN=FN,
                 TN=TN,
                 threshold=threshold,
                 study=study,
                 dist="loglogistic",
                 testdirection="less",
                 eval_threshold=c(8,9,10,11,12,13),
                 smallest=5,
                 largest=17)
```

```
less
#>
#> Hoyer AFT Model Fit
#> -----
#> Distribution : Loglogistic
#> Number of studies : 15
#> Test direction : less
#> Optimization : converged
#> -2 negative Log Likelihood : 19708.9
#>
#> Use summary() for parameter estimates.
summary(less)
#> $sdreport2
#>           Estimate Std. Error
#> beta0      2.531071186 0.0169254101
#> lambda0    0.058309997 0.0007979991
#> su0        0.065266780 0.0120139561
#> beta1      2.253821209 0.0147254382
#> lambda1    0.060530034 0.0012263705
#> su1        0.056279481 0.0105351710
#> rho       -0.411758265 0.2194120812
#> covu0u1   -0.001512462 0.0010422566
#> logitSurv1 2.880878398 0.2497852789
#> logitSurv1 0.935017349 0.2438804386
#> logitSurv1 -0.805614673 0.2439346922
#> logitSurv1 -2.380207869 0.2483362272
#> logitSurv1 -3.817698831 0.2557856931
#> logitSurv1 -5.140062330 0.2652905618
#> logitSurv0 7.745320971 0.3083465399
#> logitSurv0 5.725375142 0.3001481581
#> logitSurv0 3.918472026 0.2948244780
#> logitSurv0 2.283929334 0.2917357077
#> logitSurv0 0.791708769 0.2903955841
```

```

#> logitSurv0 -0.581001083 0.2904281547
#>
#> $sensspec
#>   threshold conflevel      Sens SensCI_Lower SensCI_Upper      Spec
#> 1         8      0.95 0.05310695    0.0332319    0.08383789 0.9995674
#> 2         9      0.95 0.28190791    0.1957591    0.38769289 0.9967485
#> 3        10      0.95 0.69117423    0.5811538    0.78308198 0.9805157
#> 4        11      0.95 0.91530555    0.8691516    0.94618765 0.9075373
#> 5        12      0.95 0.97849434    0.9649861    0.98686211 0.6881981
#> 6        13      0.95 0.99417678    0.9902443    0.99652965 0.3587023
#>   SpecCI_Lower SpecCI_Upper
#> 1   0.9992086   0.9997636
#> 2   0.9941596   0.9981919
#> 3   0.9657965   0.9889730
#> 4   0.8471134   0.9456136
#> 5   0.5554075   0.7959017
#> 6   0.2404471   0.4970569
plot(less,main="Diagnostic Test Accuracy Meta-Analysis\nLoglogistic Model")

```



How does `fitHoyer` work internally?

Step 1: How do we restructure the data set?

```
res <- restructure(
  data = diabetes,
  TP = TP,
  FP = FP,
  FN = FN,
  TN = TN,
  study = study,
  threshold = threshold,
  testdirection = "greater",
  smallest = 2,
```

```
largest = 10)
```

```
res
```

```
#> $restructured
```

```
#>      study TP   TN   D   H threshold lowerB upperB events1 events0
#> Study 01.1 Study 01 574 1389 836 2071      5.9      NA      5.9      262      1389
#> Study 01.2 Study 01 574 1389 836 2071      5.9      5.9      NA      574      682
#> Study 02.1 Study 02 617 1005 635 8740      5.0      NA      5.0       18      1005
#> Study 02.2 Study 02 607 1617 635 8740      5.1      5.0      5.1       10      612
#> Study 02.3 Study 02 600 2438 635 8740      5.2      5.1      5.2        7      821
#> Study 02.4 Study 02 581 3409 635 8740      5.3      5.2      5.3       19      971
#> Study 02.5 Study 02 563 4422 635 8740      5.4      5.3      5.4       18     1013
#> Study 02.6 Study 02 550 5384 635 8740      5.5      5.4      5.5       13      962
#> Study 02.7 Study 02 522 6267 635 8740      5.6      5.5      5.6       28      883
#> Study 02.8 Study 02 489 6966 635 8740      5.7      5.6      5.7       33      699
#> Study 02.9 Study 02 457 7534 635 8740      5.8      5.7      5.8       32      568
#> Study 02.10 Study 02 429 7927 635 8740      5.9      5.8      5.9       28      393
#> Study 02.11 Study 02 393 8172 635 8740      6.0      5.9      6.0       36      245
#> Study 02.12 Study 02 332 8460 635 8740      6.2      6.0      6.2       61      288
#> Study 02.13 Study 02 236 8670 635 8740      6.6      6.2      6.6       96      210
#> Study 02.14 Study 02 236 8670 635 8740      6.6      6.6      NA      236       70
#> Study 03.1 Study 03  36  998  49 1093      6.2      NA      6.2       13      998
#> Study 03.2 Study 03  36  998  49 1093      6.2      6.2      NA       36       95
#> Study 04.1 Study 04  16  147  41  157      6.0      NA      6.0       25      147
#> Study 04.2 Study 04  16  147  41  157      6.0      6.0      NA       16       10
#> Study 05.1 Study 05 644 1217 795 1503      6.1      NA      6.1      151     1217
#> Study 05.2 Study 05 644 1217 795 1503      6.1      6.1      NA      644      286
#> Study 06.1 Study 06 176 1286 278 2054      5.6      NA      5.6      102     1286
#> Study 06.2 Study 06  69 1867 278 2054      6.5      5.6      6.5      107      581
#> Study 06.3 Study 06  69 1867 278 2054      6.5      6.5      NA       69      187
#> Study 07.1 Study 07  72  258  88  304      6.1      NA      6.1       16      258
#> Study 07.2 Study 07  72  258  88  304      6.1      6.1      NA       72       46
#> Study 08.1 Study 08  89 1382 115 1684      5.3      NA      5.3       26     1382
#> Study 08.2 Study 08  72 1558 115 1684      5.5      5.3      5.5       17      176
#> Study 08.3 Study 08  65 1602 115 1684      5.6      5.5      5.6        7       44
#> Study 08.4 Study 08  65 1602 115 1684      5.6      5.6      NA       65       82
#> Study 09.1 Study 09 207 2704 252 5865      5.8      NA      5.8       45     2704
#> Study 09.2 Study 09 196 3308 252 5865      5.9      5.8      5.9       11      604
#> Study 09.3 Study 09 181 3877 252 5865      6.0      5.9      6.0       15      569
#> Study 09.4 Study 09 181 3877 252 5865      6.0      6.0      NA      181     1988
#> Study 10.1 Study 10  54    1  54  451      3.9      NA      3.9        0        1
#> Study 10.2 Study 10  54   45  54  451      4.7      3.9      4.7        0       44
#> Study 10.3 Study 10  46  363  54  451      5.6      4.7      5.6        8      318
#> Study 10.4 Study 10  43  389  54  451      5.7      5.6      5.7        3       26
#> Study 10.5 Study 10  23  447  54  451      6.2      5.7      6.2       20       58
#> Study 10.6 Study 10  12  451  54  451      6.8      6.2      6.8       11        4
#> Study 10.7 Study 10  12  451  54  451      6.8      6.8      NA       12        0
#> Study 11.1 Study 11 861  611 895 5050      5.0      NA      5.0       34      611
#> Study 11.2 Study 11 837 1000 895 5050      5.1      5.0      5.1       24      389
#> Study 11.3 Study 11 793 1525 895 5050      5.2      5.1      5.2       44      525
#> Study 11.4 Study 11 740 2116 895 5050      5.3      5.2      5.3       53      591
#> Study 11.5 Study 11 687 2762 895 5050      5.4      5.3      5.4       53      646
#> Study 11.6 Study 11 610 3358 895 5050      5.5      5.4      5.5      77      596
#> Study 11.7 Study 11 532 3883 895 5050      5.6      5.5      5.6      78      525
```

#> Study 11.8	Study 11	455	4277	895	5050	5.7	5.6	5.7	77	394
#> Study 11.9	Study 11	376	4585	895	5050	5.8	5.7	5.8	79	308
#> Study 11.10	Study 11	298	4782	895	5050	5.9	5.8	5.9	78	197
#> Study 11.11	Study 11	235	4883	895	5050	6.0	5.9	6.0	63	101
#> Study 11.12	Study 11	136	4984	895	5050	6.2	6.0	6.2	99	101
#> Study 11.13	Study 11	46	5045	895	5050	6.6	6.2	6.6	90	61
#> Study 11.14	Study 11	46	5045	895	5050	6.6	6.6	NA	46	5
#> Study 12.1	Study 12	184	5265	338	5903	5.7	NA	5.7	154	5265
#> Study 12.2	Study 12	184	5265	338	5903	5.7	5.7	NA	184	638
#> Study 13.1	Study 13	72	120	81	333	5.5	NA	5.5	9	120
#> Study 13.2	Study 13	70	144	81	333	5.6	5.5	5.6	2	24
#> Study 13.3	Study 13	66	170	81	333	5.7	5.6	5.7	4	26
#> Study 13.4	Study 13	58	204	81	333	5.8	5.7	5.8	8	34
#> Study 13.5	Study 13	50	234	81	333	5.9	5.8	5.9	8	30
#> Study 13.6	Study 13	40	256	81	333	6.0	5.9	6.0	10	22
#> Study 13.7	Study 13	40	256	81	333	6.0	6.0	NA	40	77
#> Study 14.1	Study 14	285	155	295	1687	4.8	NA	4.8	10	155
#> Study 14.2	Study 14	262	412	295	1687	5.1	4.8	5.1	23	257
#> Study 14.3	Study 14	253	516	295	1687	5.2	5.1	5.2	9	104
#> Study 14.4	Study 14	234	680	295	1687	5.3	5.2	5.3	19	164
#> Study 14.5	Study 14	195	969	295	1687	5.5	5.3	5.5	39	289
#> Study 14.6	Study 14	169	1099	295	1687	5.6	5.5	5.6	26	130
#> Study 14.7	Study 14	138	1211	295	1687	5.7	5.6	5.7	31	112
#> Study 14.8	Study 14	91	1409	295	1687	5.9	5.7	5.9	47	198
#> Study 14.9	Study 14	91	1409	295	1687	5.9	5.9	NA	91	278
#> Study 15.1	Study 15	108	626	114	4816	5.0	NA	5.0	6	626
#> Study 15.2	Study 15	91	2745	114	4816	5.5	5.0	5.5	17	2119
#> Study 15.3	Study 15	74	3660	114	4816	5.7	5.5	5.7	17	915
#> Study 15.4	Study 15	48	4479	114	4816	6.0	5.7	6.0	26	819
#> Study 15.5	Study 15	7	4816	114	4816	6.4	6.0	6.4	41	337
#> Study 15.6	Study 15	7	4816	114	4816	6.4	6.4	NA	7	0
#> Study 16.1	Study 16	184	181	186	5826	5.0	NA	5.0	2	181
#> Study 16.2	Study 16	164	3437	186	5826	5.5	5.0	5.5	20	3256
#> Study 16.3	Study 16	145	4719	186	5826	5.7	5.5	5.7	19	1282
#> Study 16.4	Study 16	84	5593	186	5826	6.0	5.7	6.0	61	874
#> Study 16.5	Study 16	4	5826	186	5826	6.4	6.0	6.4	80	233
#> Study 16.6	Study 16	4	5826	186	5826	6.4	6.4	NA	4	0
#> Study 17.1	Study 17	88	480	92	3692	5.0	NA	5.0	4	480
#> Study 17.2	Study 17	74	2252	92	3692	5.5	5.0	5.5	14	1772
#> Study 17.3	Study 17	61	2954	92	3692	5.7	5.5	5.7	13	702
#> Study 17.4	Study 17	35	3507	92	3692	6.0	5.7	6.0	26	553
#> Study 17.5	Study 17	4	3692	92	3692	6.4	6.0	6.4	31	185
#> Study 17.6	Study 17	4	3692	92	3692	6.4	6.4	NA	4	0
#> Study 18.1	Study 18	21	284	64	320	5.7	NA	5.7	43	284
#> Study 18.2	Study 18	21	284	64	320	5.7	5.7	NA	21	36
#> Study 19.1	Study 19	49	691	57	1132	5.1	NA	5.1	8	691
#> Study 19.2	Study 19	42	814	57	1132	5.2	5.1	5.2	7	123
#> Study 19.3	Study 19	32	994	57	1132	5.3	5.2	5.3	10	180
#> Study 19.4	Study 19	26	1045	57	1132	5.4	5.3	5.4	6	51
#> Study 19.5	Study 19	26	1045	57	1132	5.4	5.4	NA	26	87
#> Study 20.1	Study 20	596	2062	659	6145	5.0	NA	5.0	63	2062
#> Study 20.2	Study 20	392	5015	659	6145	5.5	5.0	5.5	204	2953
#> Study 20.3	Study 20	137	5999	659	6145	6.0	5.5	6.0	255	984

#> Study 20.4	Study 20	137	5999	659	6145	6.0	6.0	NA	137	146
#> Study 21.1	Study 21	187	8562	368	9674	5.5	NA	5.5	181	8562
#> Study 21.2	Study 21	187	8562	368	9674	5.5	5.5	NA	187	1112
#> Study 22.1	Study 22	9	395	17	432	5.8	NA	5.8	8	395
#> Study 22.2	Study 22	9	395	17	432	5.8	5.8	NA	9	37
#> Study 23.1	Study 23	122	215	164	304	5.5	NA	5.5	42	215
#> Study 23.2	Study 23	88	265	164	304	5.7	5.5	5.7	34	50
#> Study 23.3	Study 23	88	265	164	304	5.7	5.7	NA	88	39
#> Study 24.1	Study 24	22	129	44	164	6.1	NA	6.1	22	129
#> Study 24.2	Study 24	22	129	44	164	6.1	6.1	NA	22	35
#> Study 25.1	Study 25	79	217	80	882	6.6	NA	6.6	1	217
#> Study 25.2	Study 25	79	217	80	882	6.6	6.6	NA	79	665
#> Study 26.1	Study 26	114	203	178	223	6.9	NA	6.9	64	203
#> Study 26.2	Study 26	90	219	178	223	7.4	6.9	7.4	24	16
#> Study 26.3	Study 26	73	223	178	223	7.6	7.4	7.6	17	4
#> Study 26.4	Study 26	73	223	178	223	7.6	7.6	NA	73	0
#> Study 27.1	Study 27	135	592	178	688	5.9	NA	5.9	43	592
#> Study 27.2	Study 27	87	674	178	688	6.5	5.9	6.5	48	82
#> Study 27.3	Study 27	87	674	178	688	6.5	6.5	NA	87	14
#> Study 28.1	Study 28	575	980	627	2250	5.5	NA	5.5	52	980
#> Study 28.2	Study 28	486	1773	627	2250	6.1	5.5	6.1	89	793
#> Study 28.3	Study 28	486	1773	627	2250	6.1	6.1	NA	486	477
#> Study 29.1	Study 29	23	109	30	114	6.1	NA	6.1	7	109
#> Study 29.2	Study 29	23	109	30	114	6.1	6.1	NA	23	5
#> Study 30.1	Study 30	424	2112	616	2233	6.0	NA	6.0	192	2112
#> Study 30.2	Study 30	424	2112	616	2233	6.0	6.0	NA	424	121
#> Study 31.1	Study 31	338	4060	367	6436	5.7	NA	5.7	29	4060
#> Study 31.2	Study 31	338	4060	367	6436	5.7	5.7	NA	338	2376
#> Study 32.1	Study 32	57	32	64	44	6.0	NA	6.0	7	32
#> Study 32.2	Study 32	50	40	64	44	6.4	6.0	6.4	7	8
#> Study 32.3	Study 32	47	41	64	44	6.5	6.4	6.5	3	1
#> Study 32.4	Study 32	40	43	64	44	7.0	6.5	7.0	7	2
#> Study 32.5	Study 32	40	43	64	44	7.0	7.0	NA	40	1
#> Study 33.1	Study 33	52	79	65	116	5.6	NA	5.6	13	79
#> Study 33.2	Study 33	52	79	65	116	5.6	5.6	NA	52	37
#> Study 34.1	Study 34	111	83	112	489	5.0	NA	5.0	1	83
#> Study 34.2	Study 34	110	225	112	489	5.5	5.0	5.5	1	142
#> Study 34.3	Study 34	100	381	112	489	6.0	5.5	6.0	10	156
#> Study 34.4	Study 34	78	465	112	489	6.5	6.0	6.5	22	84
#> Study 34.5	Study 34	60	479	112	489	7.0	6.5	7.0	18	14
#> Study 34.6	Study 34	60	479	112	489	7.0	7.0	NA	60	10
#> Study 35.1	Study 35	20	60	21	90	5.9	NA	5.9	1	60
#> Study 35.2	Study 35	19	62	21	90	6.0	5.9	6.0	1	2
#> Study 35.3	Study 35	17	76	21	90	6.1	6.0	6.1	2	14
#> Study 35.4	Study 35	17	79	21	90	6.2	6.1	6.2	0	3
#> Study 35.5	Study 35	17	79	21	90	6.3	6.2	6.3	0	0
#> Study 35.6	Study 35	16	81	21	90	6.4	6.3	6.4	1	2
#> Study 35.7	Study 35	13	81	21	90	6.5	6.4	6.5	3	0
#> Study 35.8	Study 35	12	86	21	90	6.6	6.5	6.6	1	5
#> Study 35.9	Study 35	11	87	21	90	6.7	6.6	6.7	1	1
#> Study 35.10	Study 35	11	87	21	90	6.8	6.7	6.8	0	0
#> Study 35.11	Study 35	11	88	21	90	6.9	6.8	6.9	0	1
#> Study 35.12	Study 35	10	88	21	90	7.0	6.9	7.0	1	0

```

#> Study 35.13 Study 35 10 88 21 90 7.0 7.0 NA 10 2
#> Study 36.1 Study 36 25 243 27 288 5.9 NA 5.9 2 243
#> Study 36.2 Study 36 25 243 27 288 5.9 5.9 NA 25 45
#> Study 37.1 Study 37 12 268 14 293 5.9 NA 5.9 2 268
#> Study 37.2 Study 37 12 268 14 293 5.9 5.9 NA 12 25
#> Study 38.1 Study 38 13 260 19 295 5.9 NA 5.9 6 260
#> Study 38.2 Study 38 13 260 19 295 5.9 5.9 NA 13 35
#> ctype lcutmean
#> Study 01.1 1 1.234050
#> Study 01.2 3 2.038769
#> Study 02.1 1 1.151293
#> Study 02.2 2 1.619339
#> Study 02.3 2 1.638950
#> Study 02.4 2 1.658183
#> Study 02.5 2 1.677053
#> Study 02.6 2 1.695574
#> Study 02.7 2 1.713757
#> Study 02.8 2 1.731616
#> Study 02.9 2 1.749162
#> Study 02.10 2 1.766405
#> Study 02.11 2 1.783356
#> Study 02.12 2 1.808154
#> Study 02.13 2 1.855809
#> Study 02.14 3 2.094827
#> Study 03.1 1 1.258848
#> Study 03.2 3 2.063567
#> Study 04.1 1 1.242453
#> Study 04.2 3 2.047172
#> Study 05.1 1 1.250718
#> Study 05.2 3 2.055437
#> Study 06.1 1 1.207957
#> Study 06.2 2 1.797284
#> Study 06.3 3 2.087194
#> Study 07.1 1 1.250718
#> Study 07.2 3 2.055437
#> Study 08.1 1 1.180427
#> Study 08.2 2 1.686227
#> Study 08.3 2 1.713757
#> Study 08.4 3 2.012676
#> Study 09.1 1 1.225503
#> Study 09.2 2 1.766405
#> Study 09.3 2 1.783356
#> Study 09.4 3 2.047172
#> Study 10.1 1 1.027062
#> Study 10.2 2 1.454270
#> Study 10.3 2 1.635165
#> Study 10.4 2 1.731616
#> Study 10.5 2 1.782508
#> Study 10.6 2 1.870736
#> Study 10.7 3 2.109754
#> Study 11.1 1 1.151293
#> Study 11.2 2 1.619339
#> Study 11.3 2 1.638950

```



```

#> Study 11.4      2 1.658183
#> Study 11.5      2 1.677053
#> Study 11.6      2 1.695574
#> Study 11.7      2 1.713757
#> Study 11.8      2 1.731616
#> Study 11.9      2 1.749162
#> Study 11.10     2 1.766405
#> Study 11.11     2 1.783356
#> Study 11.12     2 1.808154
#> Study 11.13     2 1.855809
#> Study 11.14     3 2.094827
#> Study 12.1      1 1.216807
#> Study 12.2      3 2.021526
#> Study 13.1      1 1.198948
#> Study 13.2      2 1.713757
#> Study 13.3      2 1.731616
#> Study 13.4      2 1.749162
#> Study 13.5      2 1.766405
#> Study 13.6      2 1.783356
#> Study 13.7      3 2.047172
#> Study 14.1      1 1.130882
#> Study 14.2      2 1.598928
#> Study 14.3      2 1.638950
#> Study 14.4      2 1.658183
#> Study 14.5      2 1.686227
#> Study 14.6      2 1.713757
#> Study 14.7      2 1.731616
#> Study 14.8      2 1.757709
#> Study 14.9      3 2.038769
#> Study 15.1      1 1.151293
#> Study 15.2      2 1.657093
#> Study 15.3      2 1.722607
#> Study 15.4      2 1.766113
#> Study 15.5      2 1.824029
#> Study 15.6      3 2.079442
#> Study 16.1      1 1.151293
#> Study 16.2      2 1.657093
#> Study 16.3      2 1.722607
#> Study 16.4      2 1.766113
#> Study 16.5      2 1.824029
#> Study 16.6      3 2.079442
#> Study 17.1      1 1.151293
#> Study 17.2      2 1.657093
#> Study 17.3      2 1.722607
#> Study 17.4      2 1.766113
#> Study 17.5      2 1.824029
#> Study 17.6      3 2.079442
#> Study 18.1      1 1.216807
#> Study 18.2      3 2.021526
#> Study 19.1      1 1.161194
#> Study 19.2      2 1.638950
#> Study 19.3      2 1.658183
#> Study 19.4      2 1.677053

```



```

#> Study 19.5      3 1.994492
#> Study 20.1      1 1.151293
#> Study 20.2      2 1.657093
#> Study 20.3      2 1.748254
#> Study 20.4      3 2.047172
#> Study 21.1      1 1.198948
#> Study 21.2      3 2.003667
#> Study 22.1      1 1.225503
#> Study 22.2      3 2.030222
#> Study 23.1      1 1.198948
#> Study 23.2      2 1.722607
#> Study 23.3      3 2.021526
#> Study 24.1      1 1.250718
#> Study 24.2      3 2.055437
#> Study 25.1      1 1.290108
#> Study 25.2      3 2.094827
#> Study 26.1      1 1.312334
#> Study 26.2      2 1.966501
#> Study 26.3      2 2.014814
#> Study 26.4      3 2.165367
#> Study 27.1      1 1.234050
#> Study 27.2      2 1.823377
#> Study 27.3      3 2.087194
#> Study 28.1      1 1.198948
#> Study 28.2      2 1.756518
#> Study 28.3      3 2.055437
#> Study 29.1      1 1.250718
#> Study 29.2      3 2.055437
#> Study 30.1      1 1.242453
#> Study 30.2      3 2.047172
#> Study 31.1      1 1.216807
#> Study 31.2      3 2.021526
#> Study 32.1      1 1.242453
#> Study 32.2      2 1.824029
#> Study 32.3      2 1.864050
#> Study 32.4      2 1.908856
#> Study 32.5      3 2.124248
#> Study 33.1      1 1.207957
#> Study 33.2      3 2.012676
#> Study 34.1      1 1.151293
#> Study 34.2      2 1.657093
#> Study 34.3      2 1.748254
#> Study 34.4      2 1.831781
#> Study 34.5      2 1.908856
#> Study 34.6      3 2.124248
#> Study 35.1      1 1.234050
#> Study 35.2      2 1.783356
#> Study 35.3      2 1.800024
#> Study 35.4      2 1.816419
#> Study 35.5      2 1.832549
#> Study 35.6      2 1.848424
#> Study 35.7      2 1.864050
#> Study 35.8      2 1.879436

```

```

#> Study 35.9      2 1.894589
#> Study 35.10     2 1.909515
#> Study 35.11     2 1.924222
#> Study 35.12     2 1.938716
#> Study 35.13     3 2.124248
#> Study 36.1      1 1.234050
#> Study 36.2      3 2.038769
#> Study 37.1      1 1.234050
#> Study 37.2      3 2.038769
#> Study 38.1      1 1.234050
#> Study 38.2      3 2.038769
#>
#> $original
#>      study TP   TN   FP  FN threshold   D   H testdirection   sens
#> 1 Study 01 574 1389 682 262      5.9 836 2071      greater 0.68660287
#> 2 Study 02 617 1005 7735 18      5.0 635 8740      greater 0.97165354
#> 3 Study 02 607 1617 7123 28      5.1 635 8740      greater 0.95590551
#> 4 Study 02 600 2438 6302 35      5.2 635 8740      greater 0.94488189
#> 5 Study 02 581 3409 5331 54      5.3 635 8740      greater 0.91496063
#> 6 Study 02 563 4422 4318 72      5.4 635 8740      greater 0.88661417
#> 7 Study 02 550 5384 3356 85      5.5 635 8740      greater 0.86614173
#> 8 Study 02 522 6267 2473 113     5.6 635 8740      greater 0.82204724
#> 9 Study 02 489 6966 1774 146     5.7 635 8740      greater 0.77007874
#> 10 Study 02 457 7534 1206 178     5.8 635 8740      greater 0.71968504
#> 11 Study 02 429 7927 813 206      5.9 635 8740      greater 0.67559055
#> 12 Study 02 393 8172 568 242      6.0 635 8740      greater 0.61889764
#> 13 Study 02 332 8460 280 303      6.2 635 8740      greater 0.52283465
#> 14 Study 02 236 8670 70 399      6.6 635 8740      greater 0.37165354
#> 15 Study 03 36 998 95 13          6.2 49 1093      greater 0.73469388
#> 16 Study 04 16 147 10 25          6.0 41 157       greater 0.39024390
#> 17 Study 05 644 1217 286 151      6.1 795 1503      greater 0.81006289
#> 18 Study 06 176 1286 768 102      5.6 278 2054      greater 0.63309353
#> 19 Study 06 69 1867 187 209      6.5 278 2054      greater 0.24820144
#> 20 Study 07 72 258 46 16          6.1 88 304       greater 0.81818182
#> 21 Study 08 89 1382 302 26        5.3 115 1684      greater 0.77391304
#> 22 Study 08 72 1558 126 43        5.5 115 1684      greater 0.62608696
#> 23 Study 08 65 1602 82 50         5.6 115 1684      greater 0.56521739
#> 24 Study 09 207 2704 3161 45      5.8 252 5865      greater 0.82142857
#> 25 Study 09 196 3308 2557 56      5.9 252 5865      greater 0.77777778
#> 26 Study 09 181 3877 1988 71      6.0 252 5865      greater 0.71825397
#> 27 Study 10 54 1 450 0           3.9 54 451       greater 1.00000000
#> 28 Study 10 54 45 406 0          4.7 54 451       greater 1.00000000
#> 29 Study 10 46 363 88 8           5.6 54 451       greater 0.85185185
#> 30 Study 10 43 389 62 11          5.7 54 451       greater 0.79629630
#> 31 Study 10 23 447 4 31           6.2 54 451       greater 0.42592593
#> 32 Study 10 12 451 0 42           6.8 54 451       greater 0.22222222
#> 33 Study 11 861 611 4439 34       5.0 895 5050      greater 0.96201117
#> 34 Study 11 837 1000 4050 58      5.1 895 5050      greater 0.93519553
#> 35 Study 11 793 1525 3525 102     5.2 895 5050      greater 0.88603352
#> 36 Study 11 740 2116 2934 155     5.3 895 5050      greater 0.82681564
#> 37 Study 11 687 2762 2288 208     5.4 895 5050      greater 0.76759777
#> 38 Study 11 610 3358 1692 285     5.5 895 5050      greater 0.68156425
#> 39 Study 11 532 3883 1167 363     5.6 895 5050      greater 0.59441341

```

#> 40	Study	11	455	4277	773	440	5.7	895	5050	greater	0.50837989
#> 41	Study	11	376	4585	465	519	5.8	895	5050	greater	0.42011173
#> 42	Study	11	298	4782	268	597	5.9	895	5050	greater	0.33296089
#> 43	Study	11	235	4883	167	660	6.0	895	5050	greater	0.26256983
#> 44	Study	11	136	4984	66	759	6.2	895	5050	greater	0.15195531
#> 45	Study	11	46	5045	5	849	6.6	895	5050	greater	0.05139665
#> 46	Study	12	184	5265	638	154	5.7	338	5903	greater	0.54437870
#> 47	Study	13	72	120	213	9	5.5	81	333	greater	0.88888889
#> 48	Study	13	70	144	189	11	5.6	81	333	greater	0.86419753
#> 49	Study	13	66	170	163	15	5.7	81	333	greater	0.81481481
#> 50	Study	13	58	204	129	23	5.8	81	333	greater	0.71604938
#> 51	Study	13	50	234	99	31	5.9	81	333	greater	0.61728395
#> 52	Study	13	40	256	77	41	6.0	81	333	greater	0.49382716
#> 53	Study	14	285	155	1532	10	4.8	295	1687	greater	0.96610169
#> 54	Study	14	262	412	1275	33	5.1	295	1687	greater	0.88813559
#> 55	Study	14	253	516	1171	42	5.2	295	1687	greater	0.85762712
#> 56	Study	14	234	680	1007	61	5.3	295	1687	greater	0.79322034
#> 57	Study	14	195	969	718	100	5.5	295	1687	greater	0.66101695
#> 58	Study	14	169	1099	588	126	5.6	295	1687	greater	0.57288136
#> 59	Study	14	138	1211	476	157	5.7	295	1687	greater	0.46779661
#> 60	Study	14	91	1409	278	204	5.9	295	1687	greater	0.30847458
#> 61	Study	15	108	626	4190	6	5.0	114	4816	greater	0.94736842
#> 62	Study	15	91	2745	2071	23	5.5	114	4816	greater	0.79824561
#> 63	Study	15	74	3660	1156	40	5.7	114	4816	greater	0.64912281
#> 64	Study	15	48	4479	337	66	6.0	114	4816	greater	0.42105263
#> 65	Study	15	7	4816	0	107	6.4	114	4816	greater	0.06140351
#> 66	Study	16	184	181	5645	2	5.0	186	5826	greater	0.98924731
#> 67	Study	16	164	3437	2389	22	5.5	186	5826	greater	0.88172043
#> 68	Study	16	145	4719	1107	41	5.7	186	5826	greater	0.77956989
#> 69	Study	16	84	5593	233	102	6.0	186	5826	greater	0.45161290
#> 70	Study	16	4	5826	0	182	6.4	186	5826	greater	0.02150538
#> 71	Study	17	88	480	3212	4	5.0	92	3692	greater	0.95652174
#> 72	Study	17	74	2252	1440	18	5.5	92	3692	greater	0.80434783
#> 73	Study	17	61	2954	738	31	5.7	92	3692	greater	0.66304348
#> 74	Study	17	35	3507	185	57	6.0	92	3692	greater	0.38043478
#> 75	Study	17	4	3692	0	88	6.4	92	3692	greater	0.04347826
#> 76	Study	18	21	284	36	43	5.7	64	320	greater	0.32812500
#> 77	Study	19	49	691	441	8	5.1	57	1132	greater	0.85964912
#> 78	Study	19	42	814	318	15	5.2	57	1132	greater	0.73684211
#> 79	Study	19	32	994	138	25	5.3	57	1132	greater	0.56140351
#> 80	Study	19	26	1045	87	31	5.4	57	1132	greater	0.45614035
#> 82	Study	20	596	2062	4083	63	5.0	659	6145	greater	0.90440061
#> 81	Study	20	392	5015	1130	267	5.5	659	6145	greater	0.59484067
#> 83	Study	20	137	5999	146	522	6.0	659	6145	greater	0.20789074
#> 84	Study	21	187	8562	1112	181	5.5	368	9674	greater	0.50815217
#> 85	Study	22	9	395	37	8	5.8	17	432	greater	0.52941176
#> 86	Study	23	122	215	89	42	5.5	164	304	greater	0.74390244
#> 87	Study	23	88	265	39	76	5.7	164	304	greater	0.53658537
#> 88	Study	24	22	129	35	22	6.1	44	164	greater	0.50000000
#> 89	Study	25	79	217	665	1	6.6	80	882	greater	0.98750000
#> 90	Study	26	114	203	20	64	6.9	178	223	greater	0.64044944
#> 91	Study	26	90	219	4	88	7.4	178	223	greater	0.50561798
#> 92	Study	26	73	223	0	105	7.6	178	223	greater	0.41011236

```

#> 93 Study 27 135 592 96 43 5.9 178 688 greater 0.75842697
#> 94 Study 27 87 674 14 91 6.5 178 688 greater 0.48876404
#> 95 Study 28 575 980 1270 52 5.5 627 2250 greater 0.91706539
#> 96 Study 28 486 1773 477 141 6.1 627 2250 greater 0.77511962
#> 97 Study 29 23 109 5 7 6.1 30 114 greater 0.76666667
#> 98 Study 30 424 2112 121 192 6.0 616 2233 greater 0.68831169
#> 99 Study 31 338 4060 2376 29 5.7 367 6436 greater 0.92098093
#> 100 Study 32 57 32 12 7 6.0 64 44 greater 0.89062500
#> 101 Study 32 50 40 4 14 6.4 64 44 greater 0.78125000
#> 102 Study 32 47 41 3 17 6.5 64 44 greater 0.73437500
#> 103 Study 32 40 43 1 24 7.0 64 44 greater 0.62500000
#> 104 Study 33 52 79 37 13 5.6 65 116 greater 0.80000000
#> 105 Study 34 111 83 406 1 5.0 112 489 greater 0.99107143
#> 106 Study 34 110 225 264 2 5.5 112 489 greater 0.98214286
#> 107 Study 34 100 381 108 12 6.0 112 489 greater 0.89285714
#> 108 Study 34 78 465 24 34 6.5 112 489 greater 0.69642857
#> 109 Study 34 60 479 10 52 7.0 112 489 greater 0.53571429
#> 110 Study 35 20 60 30 1 5.9 21 90 greater 0.95238095
#> 111 Study 35 19 62 28 2 6.0 21 90 greater 0.90476190
#> 112 Study 35 17 76 14 4 6.1 21 90 greater 0.80952381
#> 113 Study 35 17 79 11 4 6.2 21 90 greater 0.80952381
#> 114 Study 35 17 79 11 4 6.3 21 90 greater 0.80952381
#> 115 Study 35 16 81 9 5 6.4 21 90 greater 0.76190476
#> 116 Study 35 13 81 9 8 6.5 21 90 greater 0.61904762
#> 117 Study 35 12 86 4 9 6.6 21 90 greater 0.57142857
#> 118 Study 35 11 87 3 10 6.7 21 90 greater 0.52380952
#> 119 Study 35 11 87 3 10 6.8 21 90 greater 0.52380952
#> 120 Study 35 11 88 2 10 6.9 21 90 greater 0.52380952
#> 121 Study 35 10 88 2 11 7.0 21 90 greater 0.47619048
#> 122 Study 36 25 243 45 2 5.9 27 288 greater 0.92592593
#> 123 Study 37 12 268 25 2 5.9 14 293 greater 0.85714286
#> 124 Study 38 13 260 35 6 5.9 19 295 greater 0.68421053
#> spec fpr
#> 1 0.670690488 0.329309512
#> 2 0.114988558 0.885011442
#> 3 0.185011442 0.814988558
#> 4 0.278947368 0.721052632
#> 5 0.390045767 0.609954233
#> 6 0.505949657 0.494050343
#> 7 0.616018307 0.383981693
#> 8 0.717048055 0.282951945
#> 9 0.797025172 0.202974828
#> 10 0.862013730 0.137986270
#> 11 0.906979405 0.093020595
#> 12 0.935011442 0.064988558
#> 13 0.967963387 0.032036613
#> 14 0.991990847 0.008009153
#> 15 0.913083257 0.086916743
#> 16 0.936305732 0.063694268
#> 17 0.809713906 0.190286094
#> 18 0.626095424 0.373904576
#> 19 0.908958130 0.091041870
#> 20 0.848684211 0.151315789

```

```

#> 21 0.820665083 0.179334917
#> 22 0.925178147 0.074821853
#> 23 0.951306413 0.048693587
#> 24 0.461040068 0.538959932
#> 25 0.564023870 0.435976130
#> 26 0.661040068 0.338959932
#> 27 0.002217295 0.997782705
#> 28 0.099778271 0.900221729
#> 29 0.804878049 0.195121951
#> 30 0.862527716 0.137472284
#> 31 0.991130820 0.008869180
#> 32 1.000000000 0.000000000
#> 33 0.120990099 0.879009901
#> 34 0.198019802 0.801980198
#> 35 0.301980198 0.698019802
#> 36 0.419009901 0.580990099
#> 37 0.546930693 0.453069307
#> 38 0.664950495 0.335049505
#> 39 0.768910891 0.231089109
#> 40 0.846930693 0.153069307
#> 41 0.907920792 0.092079208
#> 42 0.946930693 0.053069307
#> 43 0.966930693 0.033069307
#> 44 0.986930693 0.013069307
#> 45 0.999009901 0.000990099
#> 46 0.891919363 0.108080637
#> 47 0.360360360 0.639639640
#> 48 0.432432432 0.567567568
#> 49 0.510510511 0.489489489
#> 50 0.612612613 0.387387387
#> 51 0.702702703 0.297297297
#> 52 0.768768769 0.231231231
#> 53 0.091879075 0.908120925
#> 54 0.244220510 0.755779490
#> 55 0.305868405 0.694131595
#> 56 0.403082395 0.596917605
#> 57 0.574392413 0.425607587
#> 58 0.651452282 0.348547718
#> 59 0.717842324 0.282157676
#> 60 0.835210433 0.164789567
#> 61 0.129983389 0.870016611
#> 62 0.569975083 0.430024917
#> 63 0.759966777 0.240033223
#> 64 0.930024917 0.069975083
#> 65 1.000000000 0.000000000
#> 66 0.031067628 0.968932372
#> 67 0.589941641 0.410058359
#> 68 0.809989701 0.190010299
#> 69 0.960006866 0.039993134
#> 70 1.000000000 0.000000000
#> 71 0.130010834 0.869989166
#> 72 0.609967497 0.390032503
#> 73 0.800108342 0.199891658

```

```

#> 74 0.949891658 0.050108342
#> 75 1.000000000 0.000000000
#> 76 0.887500000 0.112500000
#> 77 0.610424028 0.389575972
#> 78 0.719081272 0.280918728
#> 79 0.878091873 0.121908127
#> 80 0.923144876 0.076855124
#> 82 0.335557364 0.664442636
#> 81 0.816110659 0.183889341
#> 83 0.976240846 0.023759154
#> 84 0.885052719 0.114947281
#> 85 0.914351852 0.085648148
#> 86 0.707236842 0.292763158
#> 87 0.871710526 0.128289474
#> 88 0.786585366 0.213414634
#> 89 0.246031746 0.753968254
#> 90 0.910313901 0.089686099
#> 91 0.982062780 0.017937220
#> 92 1.000000000 0.000000000
#> 93 0.860465116 0.139534884
#> 94 0.979651163 0.020348837
#> 95 0.435555556 0.564444444
#> 96 0.788000000 0.212000000
#> 97 0.956140351 0.043859649
#> 98 0.945812808 0.054187192
#> 99 0.630826600 0.369173400
#> 100 0.727272727 0.272727273
#> 101 0.909090909 0.090909091
#> 102 0.931818182 0.068181818
#> 103 0.977272727 0.022727273
#> 104 0.681034483 0.318965517
#> 105 0.169734151 0.830265849
#> 106 0.460122699 0.539877301
#> 107 0.779141104 0.220858896
#> 108 0.950920245 0.049079755
#> 109 0.979550102 0.020449898
#> 110 0.666666667 0.333333333
#> 111 0.688888889 0.311111111
#> 112 0.844444444 0.155555556
#> 113 0.877777778 0.122222222
#> 114 0.877777778 0.122222222
#> 115 0.900000000 0.100000000
#> 116 0.900000000 0.100000000
#> 117 0.955555556 0.044444444
#> 118 0.966666667 0.033333333
#> 119 0.966666667 0.033333333
#> 120 0.977777778 0.022222222
#> 121 0.977777778 0.022222222
#> 122 0.843750000 0.156250000
#> 123 0.914675768 0.085324232
#> 124 0.881355932 0.118644068

```


Step 2: How do we get initial values?

```
init <- getInitParms(res$restructured,
                     dist="loglogistic")
init
#>               beta0_init lambda0_init beta1_init lambda1_init su0_init
#> Initial value   1.681153   0.04964348   1.815414   0.08091965 0.1552818
#>               su1_init coru0u1_init distcode
#> Initial value 0.1284618   0.3948626       3
```

Step 3: How do we fit the model?

```
hoyerAFT <- fitHoyerAFT(data=res,
                        init=init,
                        threshold=c(5.0,5.5,6.0,6.5,7.0,7.5))
```

```
hoyerAFT
#>
#> Hoyer AFT Model Fit
#> -----
#> Distribution : Loglogistic
#> Number of studies : 38
#> Test direction : greater
#> Optimization : converged
#> -2 negative Log Likelihood : 209322
#>
#> Use summary() for parameter estimates.
summary(hoyerAFT)
#> $sdreport2
#>           Estimate   Std. Error
#> beta0       1.705691148 0.0095630146
#> lambda0     0.039502004 0.0001857265
#> su0         0.058648679 0.0067784870
#> beta1       1.825337107 0.0145652403
#> lambda1     0.056398479 0.0009244722
#> su1         0.088320631 0.0109564611
#> rho         0.863032573 0.0454410883
#> covu0u1     0.004470412 0.0011417200
#> logitSurv1  3.828103143 0.2635035151
#> logitSurv1  2.138160760 0.2593347627
#> logitSurv1  0.595364241 0.2580790576
#> logitSurv1 -0.823870980 0.2591073990
#> logitSurv1 -2.137877556 0.2619046729
#> logitSurv1 -3.361188412 0.2660542694
#> logitSurv0  2.436667140 0.2426039991
#> logitSurv0  0.023873625 0.2420917690
#> logitSurv0 -2.178834277 0.2420878132
#> logitSurv0 -4.205129122 0.2424751080
#> logitSurv0 -6.081185122 0.2431661379
#> logitSurv0 -7.827751473 0.2440943827
#>
#> $sensspec
#>   threshold conflevel      Sens SensCI_Lower SensCI_Upper      Spec
#> 1         5.0       0.95 0.97871219   0.96482617   0.98718904 0.08041904
```

```

#> 2      5.5      0.95 0.89455725 0.83615237 0.93379170 0.49403188
#> 3      6.0      0.95 0.64459500 0.52236970 0.75048299 0.89833265
#> 4      6.5      0.95 0.30494258 0.20887630 0.42164362 0.98530044
#> 5      7.0      0.95 0.10546947 0.06591501 0.16457780 0.99771974
#> 6      7.5      0.95 0.03353069 0.02018063 0.05521454 0.99960164
#> SpecCI_Lower SpecCI_Upper
#> 1      0.0515556 0.1233403
#> 2      0.3779250 0.6107860
#> 3      0.8461021 0.9342152
#> 4      0.9765667 0.9908097
#> 5      0.9963326 0.9985830
#> 6      0.9993574 0.9997531

```

References

- Reitsma, J. B., et al. (2005). Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of Clinical Epidemiology*, 58(10), 982–990.
- Rutter, C. M., & Gatsonis, C. A. (2001). A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Statistics in Medicine*, 20(19), 2865–2884.
- Harbord, R. M., Deeks, J. J., Egger, M., Whiting, P., & Sterne, J. A. C. (2007). A unification of models for meta-analysis of diagnostic accuracy studies. *Biostatistics*, 8(2), 239–251.
- Riley, R. D., Ensor, J., Jackson, D., & Burke, D. L. (2018). Deriving percentage study weights in multi-parameter meta-analysis models. *Statistical Methods in Medical Research*, 27(10), 2885–2905.
- Hoyer, A., Hirt, S., Kuss, O. (2018). Meta-analysis of full ROC curves using bivariate time-to-event models for interval-censored data. *Research Synthesis Methods*, 9(1).
- Deeks, J. J., Bossuyt, P. M., Leeflang, M. M., & Takwoingi, Y. (editors) (2023). *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy*. Version 2.0 (updated July 2023). Cochrane. Available from <https://training.cochrane.org/handbook-diagnostic-test-accuracy/current>