

DTMC

1. Basic Knowledge

2. Introduction

2.1. Random Variables and Random Process

2.2. Definition

2.3. State Transition Matrix

2.4. Classification of States

2.4.1. Accessible

2.4.2. Communicate

2.4.3. Irreducible

2.4.4. Transient and Recurrent and Absorbing

2.4.4. Periodicity

2.4.5. Ergodicity

3. Stationary and Limiting Distributions

3.1. Limiting Distributions (Steady state distribution)

3.2. Stationary distribution

3.3. Relations and Properties

3.4. Theorem

1. Basic Knowledge

Σ Markov Chains

DTMC

- State Transition Matrix

① States: $\Pi_k = (\pi_1, \pi_2, \pi_3)$

② Transition Matrix:

$$\begin{matrix} & \begin{matrix} P_1 & P_2 & P_3 \end{matrix} \\ \begin{matrix} P_1 \\ P_2 \\ P_3 \end{matrix} & \begin{bmatrix} P_{11} & P_{12} & P_{13} \\ P_{21} & P_{22} & P_{23} \\ P_{31} & P_{32} & P_{33} \end{bmatrix} \end{matrix} \rightarrow \text{end state}$$

Start State

③ Equation:

$$\pi_{k+1} = \pi_k \cdot P$$

- Stationary Distribution

two Definitions:

① $\lim_{n \rightarrow \infty} (P)^n =$

② $\pi \cdot P = \pi \quad \pi \rightarrow \text{Steady State Probability}$

Properties:

① $\lim_{n \rightarrow \infty} P^n$: each row are equal

② $\pi_j = \lim_{n \rightarrow \infty} P_{ij}^n$

- Property of DTMC

① Accessibility: $j \leftarrow i$

State j is accessible from state i if $P_{ij}^{(n)} > 0$ for some $n \geq 0$

② Communicability: $j \leftrightarrow i$

State j is communicate if $j \leftarrow i$ & $i \leftarrow j$

③ Irreducibility: (property of a chain)

A Markov Chain is irreducible

\Leftrightarrow all states belong to one class (all states communicate with each other)

Lemma:

there exists some $n > 0$, for all i and j , $P_{ij}^{(n)} > 0$

\Rightarrow all states communicate \Rightarrow the Markov Chain irreducible

④ Periodicity: (property of a class)

Period of j : the greatest common divisor (GCD) of the set of integers, such that $P_{jj}^{(n)} > 0$

Aperiodic: A state has period 1.

Periodicity is a class property \rightarrow All nodes in the same class has the same property

Lemma:

A MC with Periodical state

\Rightarrow there is no steady state

Lemma 2:

A Markov chain Periodical + reducibility

\Rightarrow there is no steady state

① reducibility: $3 \nrightarrow 1$



If start at 3 $\pi = (0, 0, 1)$

If start at 1 π will not be $(0, 0, 1)$

② Periodical:



$$P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

π will change in sequence: (start from $(1, 0)$)

$(1, 0) \rightarrow (0, 1) \rightarrow (1, 0) \rightarrow \dots$

- Transient and Recurrent State (for state i)

① Transient:

there is a probability that the process will

move to state j and never return to state i

class property

② Recurrent:

The process returns to state i with probability 1

\star (State i is no transient

\rightarrow must be recurrent)

class property

③ Absorbing: if $P_{ii} = 1$

(that means, after arriving state i , we can never

leave $\rightarrow i$)

\star if i is Absorbing $\rightarrow i$ is recurrent

- Classes

Two states are in the same class

$\hat{=}$ two states communicate with each other

\Leftrightarrow A way to solve reducibility

$$M = (1-p)B + p \cdot A$$

where: $A \rightarrow$ original transition matrix


$$B \rightarrow \frac{1}{n} \begin{bmatrix} 1 & 1 & 1 \\ \vdots & \vdots & \vdots \\ 1 & 1 & 1 \end{bmatrix}_{n \times n}$$

that is:

① create virtual connection to all other nodes

② weight of new connections lower than original ones

e.g.



$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0.5 & 0 & 0.5 \end{bmatrix}$$

take $p=0.3$, $n=3$

$$\therefore M = 0.7 \cdot A + 0.3/3 \cdot \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

- Ergodicity

If All states in a MC: recurrent, aperiodic and communicate with each other

\Leftrightarrow ergodicity

★ If a chain is ergodic

\Rightarrow the steady state exists

2. Introduction

2.1. Random Variables and Random Process

- A **random variable** is a variable whose value is unknown or a function that assigns values to each of an experiment's outcomes.
- A **random process** is a **collection of random variables** usually indexed by time.

2.2. Definition

Consider the random process $X_n, n = 0, 1, 2, \dots$, where $R_{X_i} = S \subset 0, 1, 2, \dots$. We say that this process is a **Markov chain** if:

$$\begin{aligned} P(X_{m+1} = j | X_m = i, X_{m-1} = i_{m-1}, \dots, X_0 = i_0) \\ = P(X_{m+1} = j | X_m = i) \end{aligned}$$

for all $m, j, i, i_0, i_1, \dots, i_{m-1}$. If the number of states is finite, e.g., $S = 0, 1, 2, \dots, r$, we call it a **finite Markov chain**.

- $P(X_{m+1} = j | X_m = i)$ is called **transition probabilities**

2.3.State Transition Matrix

We often list the transition probabilities in a matrix. The matrix is called the **state transition matrix** or **transition probability matrix** and is usually shown by **P**.

Assuming the states are **1, 2, ..., r**, then the state transition matrix is given by

$$P = \begin{bmatrix} p_{11} & p_{12} & \dots & p_{1r} \\ p_{21} & p_{22} & \dots & p_{2r} \\ \vdots & \vdots & \vdots & \vdots \\ p_{r1} & p_{r2} & \dots & p_{rr} \end{bmatrix}.$$

Note that $p_{ij} \geq 0$ and for all i , we have

$$\begin{aligned} \sum_{k=1}^r p_{ik} &= \sum_{k=1}^r P(X_{m+1} = k | X_m = i) \\ &= 1. \end{aligned}$$

2.4. Classification of States

2.4.1. Accessible

We say that state j is accessible from state i , written as $i \rightarrow j$, if $p_{ij}^{(n)} > 0$ for some n .

We assume every state is accessible from itself since $p_{ii}^{(0)} = 1$.

2.4.2. Communicate

Two states i and j are said to **communicate**, written as $i \leftrightarrow j$, if they are accessible from each other. In other words,

$$i \leftrightarrow j \text{ means } i \rightarrow j \text{ and } j \rightarrow i.$$

- Communication is an **equivalence relation**

2.4.3. Irreducible

A Markov chain is said to be **irreducible** if all states communicate with each other.

2.4.4. Transient and Recurrent and Absorbing

For any state i , we define

$$f_{ii} = P(X_n = i, \text{ for some } n \geq 1 | X_0 = i).$$

State i is **recurrent** if $f_{ii} = 1$, and it is **transient** if $f_{ii} < 1$.

- if a state is not recurrent, then it will be transient

A state i is called **absorbing** if there are no outgoing transitions from the state. that is $p_{ii} = 1$

- an absorbing state is a recurrent state

2.4.4. Periodicity

The **period** of a state i is the largest integer d satisfying the following property:

$p_{ii}^{(n)} = 0$, whenever n is not divisible by d . The period of i is shown by $d(i)$. If $p_{ii}^{(n)} = 0$, for all $n > 0$, then we let $d(i) = \infty$.

- If $d(i) > 1$, we say that state i is **periodic**
- If $d(i) = 1$, we say that state i is **aperiodic**.

or we can say:

The **period** of state j is the **greatest common divisor (GCD)** of the set of integer, such that $P_{jj}^n > 0$

2.4.5. Ergodicity

If all states in a chain are recurrent, aperiodic, and communicate with each other, the chain is said to be ergodic

3. Stationary and Limiting Distributions

3.1. Limiting Distributions (Steady state distribution)

The probability distribution $\pi = [\pi_0, \pi_1, \pi_2, \dots]$ is called the **limiting distribution** of the Markov chain X_n iff

$$\pi_j = \lim_{n \rightarrow \infty} P(X_n = j | X_0 = i)$$

for all $i, j \in S$, and we have

$$\sum_{j \in S} \pi_j = 1$$

when a limiting distribution exists, it does not depend on the initial state ($X_0 = i$), so we can write

$$\pi_j = \lim_{n \rightarrow \infty} P(X_n = j), \text{ for all } j \in S.$$

3.2. Stationary distribution

$$\pi = \pi P, \sum_{j \in S} \pi_j = 1$$

3.3. Relations and Properties

- The limiting distribution of a Markov chain is a stationary distribution of the Markov chain.
- Limiting distribution is Unique (because it need suitable for all initial state)
- Stationary Distribution may not be unique (linear equation groups' solution)
- Not All Markov Chains Have a Stationary Distribution (linear equation group's solution)
- Not All Markov Chains Have Limiting Distribution
- if a limiting distribution exists, it is the unique stationary distribution

3.4. Theorem

- If a chain is **irreducible** and **aperiodic**, it will has a unique stationary distribution, and this stationary distribution will be its limiting distribution