Can satellite altimetry and gravimetry data be used to evaluate Arctic-Ocean satellite sea surface salinity? – insight from ECCO state estimation

Tony Lee, Adam Wang, Severine Fournier, Rom Kwok
NASA Jet Propulsion Laboratory, California Institute of Technology

Special thanks to Ou Wang of JPL for providing ECCO-v4 model-data differences in the Arctic Ocean and Ichiro Fukumori of JPL for discussion

Motivation, rationales, and approach

Motivation & rationales:

- Changes of Arctic Ocean freshwater are important to ocean, climate, & BGC.
- Sea surface salinity (SSS) data are important to study Arctic freshwater changes, ocean-ice interaction, and interaction with the North Atlantic Ocean.
- Large uncertainties of L-band satellite SSS; but how bad is the S/N radio?
- Lack of in-situ salinity data to validate satellite SSS.
- SSH from altimetry & OBP from gravimetry can be used to infer columnintegrated freshwater changes in the Arctic Ocean (e.g., Morrison et al. 2012)


Questions addressed: To what extent Arctic SSS & SSH-OBP variations are coherent? Can SSS-OBP data can be used to evaluate satellite SSS?

Significance: validated satellite SSS are complementary to SSH-OBP data to study Arctic freshwater changes due to better sampling (SMOS+Aquarius+SMAP).


Approach: Examine relationships of SSS & SSH-OBP in ECCO-V4 for a proof-of-concept study as a 1st step.

Uncertainty characteristics of L-band Satellite SSS: Tropics & subtropics ✓ high-latitudes ★

Zonally averaged STD of ΔSSS for (Aquarius V4.0 - Argo-SIO) & (Argo-SIO - Argo-UH) (Argo-SIO & Argo-UH are two different "ground truth" datasets)




SSS signals in the Arctic Ocean are large! So how bad is the S/N ratio for L-band satellite SSS?




Timmermans et al. (2014) and NOAA Arctic Report Card 2013 Update. Gradient estimates provided by Mike Steele of APL, University of Washington

Comparison of time-mean salinity profile in the Arctic Ocean between Ice-Tethered Profiler data & co-located ECCO estimates


ECCO-v4 reproduces the structure <200 m well, which dominate the steric changes



Local correlation of SSS & 0-200 halosteric height: they are coherent over much of the Arctic Ocean




Local correlation of SSS & SSH: poor correlation around the Arctic shelves (because of barotropic coastal Kelvin waves, unrelated to SSS)



Without Seasonal Cycle

With Seasonal Cycle


Local correlation of SSS & SSH-OBP: much strong correlation around the shelves (because barotropic wave signals have been removed)




With Seasonal Cycle

Without Seasonal Cycle


Linear regression slope for SSS vs. SSH-OBP (in psu/cm): good "calibration" sensitivity around the shelves ~ 0.3 psu/cm



SSS & SSH-OBP relationships in Chukchi Shelf and Siberian Shelf



Standard deviation of Arctic SSS from Aquarius & SMAP: large differences in magnitudes!



Ongoing work: use altimetry-gravimetry data to evaluate satellite SSS and improve Arctic Ocean SSS retrievals.

Summary

- ECCO ocean-sea ice state estimation product illustrates the strong relationship of SSS and SSH-OBP variations around Arctic shelves.
- This relationship could be used to evaluate satellite SSS in the Arctic Ocean using altimetry-gravimetry data and improve SSS retrievals.