

# **1 Genotypic variation in a foundation tree 2 results in heritable ecological network 3 structure**

**4 Matthew K. Lau<sup>1,2</sup>, Louis J. Lamit<sup>1,3,4</sup>, Rikke Reese Næsborg<sup>1,5</sup>, Stuart R.  
5 Borrett<sup>6</sup>, Matthew A. Bowker<sup>7</sup>, and Thomas G. Whitham<sup>1,8</sup>**

**6 <sup>1</sup>Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011,  
7 USA**

**8 <sup>2</sup>Harvard Forest, Harvard University, 324 N Main St, Petersham, MA 01366, USA**

**9 <sup>3</sup>Department of Biology, State University of New York College of Environmental  
10 Sciences, Syracuse University, 107 College Place Syracuse, NY 13244, USA**

**11 <sup>4</sup>Department of Environmental Forest Biology and Forestry, Syracuse, NY 13210, USA**

**12 <sup>5</sup>Santa Barbara Botanic Garden, 1212 Mission Canyon Road, Santa Barbara, CA 93105**

**13 <sup>6</sup>Department of Biology and Marine Biology, University of North Carolina Wilmington,  
14 601 South College Road, Wilmington, NC 28403, USA**

**15 <sup>7</sup>Duke Network Analysis Center, Duke University, Durham, NC 27708, USA**

**16 <sup>8</sup>School of Forestry, 200 E. Pine Knoll Dr., Northern Arizona University, Flagstaff, AZ  
17 86011, USA**

**18 <sup>9</sup>Center for Adaptable Western Landscapes, Northern Arizona University, Flagstaff, AZ  
19 86011, USA**

## **20 ABSTRACT**

Elucidating the genetic basis to ecological network structure is fundamental to understanding evolution in complex ecosystems. Although previous work has demonstrated that genetic variation can influence food webs and trophic chains, we are unaware of a study that quantified the contribution of phenotypic variation to heritable variation in network structure. To examine this, in a 20+ year old common garden we observed epiphytic lichens associated with narrowleaf cottonwood (*Populus angustifolia*), a riparian ecosystem foundation species. We constructed and conducted genetic analyses of signed, weighted, directed lichen interaction networks on individual trees. We found three primary results. First, genotype identity significantly predicted lichen network similarity; i.e., replicates of the same genotype supported more similar lichen networks than different genotypes. Second, broad sense heritability estimates showed that plant genotype explained network similarity ( $H^2 = 0.41$ ), degree ( $H^2 = 0.32$ ) and centralization ( $H^2 = 0.33$ ). Third, of several tree phenotypic traits examined, bark roughness was both heritable ( $H^2 = 0.32$ ) and significantly predicted by lichen network similarity ( $R^2 = 0.26$ ). These results support a mechanistic, genetic pathway from variation in a heritable tree trait to ecological network structure and demonstrate that evolution can act at the community level to influence not only abundances of organisms but also interactions at the scale of entire networks. Given that network structure can influence system-wide stability and resilience, our findings have important implications for how evolution acts in ecosystems.

Keywords: networks | heritability | community | genetics | lichen | cottonwood | *Populus* | common garden

21 **INTRODUCTION**

22 Evolution occurs in the context of complex ecological networks. Community genetics  
23 studies have shown that genetic variation in foundation species, which have large effects  
24 on ecosystems by modulating and stabilizing local conditions (Ellison et al., 2005),  
25 plays a significant role in defining distinct communities of interacting organisms: such  
26 as endophytes, pathogens, lichens, arthropods, and soil microbes (Busby et al., 2015;  
27 Barbour et al., 2009; Lamit et al., 2015a). Multiple studies have now demonstrated that  
28 genetic variation influences numerous functional traits (e.g., phytochemical, phenologi-  
29 cal, morphological) that in combination result in a multivariate functional trait phenotype  
30 (Holeski et al., 2012) in which individual plant genotypes support different communities  
31 and ecosystem processes (Bailey et al., 2009; Whitham et al., 2012). Recently, the  
32 importance of genetic variation in structuring ecological systems was reviewed, and  
33 not only were many instances of strong genetic effects found in many ecosystems but  
34 the effect of intraspecific variation was at times greater than inter-specific variation  
35 (Des Roches et al., 2018). There is now evidence to support that selection, acting on  
36 this heritable variation, tends to occur among groups of species (Wade, 2007) and that  
37 genetic variation and phylogenetic relatedness contribute to variation in community  
38 assembly (Crutsinger, 2016) and species interactions (Whitham et al., 2006; Bailey et al.,  
39 2009; Moya-Laraño, 2011), which shape the structure of ecological interaction networks  
40 (Rezende et al., 2007; Guimarães et al., 2007; Gómez et al., 2009).

41 Empirical and theoretical work in network ecology and evolutionary biology point  
42 to the need for examinations of the genetic basis of ecological network structure. Anal-  
43 yses of ecological networks have demonstrated that indirect effects can lead to self-  
44 organization, producing sign-changing, amplifying and/or dampening effects (Fath and  
45 Patten, 1998; Newman, 2006; Sole and Bascompte, 2006), and other studies have demon-  
46 strated that indirect effects of interactions among species can lead to network structures  
47 that amplify or dampen the effects of selection, such as the formation of star-like struc-  
48 tures in which there is a “central” species or group of species that interact with other  
49 peripheral species can amplify selection events (Lieberman et al., 2005). Also, work by  
50 Toju et al. (2014, 2016, 2017) observed consistent patterns of centralized interactions of  
51 species modules (i.e., groups of species that interact more strongly within their group  
52 than with other species) focused around hubs of plant-fungal interactions. In other words,  
53 a small number of plant and fungal symbionts tended to have disproportionate numbers  
54 of interactions with other species and likely are the drivers in determining community  
55 assembly, structure and dynamics. Interspecific indirect genetic effects (IIGE) theory  
56 (*sensu* Shuster et al. (2006)) in evolutionary biology also point to the importance of  
57 studying the genetics of interaction network structure. Genetically based differences in  
58 network structure among individuals can be acted upon by natural selection when there  
59 are fitness consequences of different networks of IIGEs, leading to community evolution  
60 per Whitham et al. (2020) and, by extension, network evolution. For example, although  
61 the analysis was of abundances rather than interaction networks, Gehring et al. (2014,  
62 2017) found that the mycorrhizal communities on the roots of drought tolerant and intol-  
63 erant trees are dominated by different orders of ectomycorrhizal fungal mutualists that  
64 also differ in the benefits they provide that enhance tree performance. Because drought  
65 tolerant genotypes are three times more likely to survive record droughts, selection acts

66 both on the tree and its fungal community and with increased drought the community  
67 phenotype has changed over time. Also, in an antagonistic interaction context, Busby  
68 et al. (2015) found that with the addition of a damaging leaf pathogen to cottonwoods  
69 in a common garden, the impacts of these strong interactors results in a different and  
70 diminished community of arthropods relative to control trees. This effect was dependent  
71 on the hybrid cross-type of the tree and was, therefore, determined by tree genetics.  
72 These examples collectively support the possibility that selection acting on the tree may  
73 alter the network structure of associated communities in which different networks of  
74 communities are most likely to survive pathogen outbreaks. Regardless of whether the  
75 IIGE is unilateral (i.e., tree affects the community) or reciprocal (i.e., the community  
76 also affects the relative fitness of the tree), selection at the level of the tree population or  
77 its community, or both, can change network structure and alter community dynamics  
78 (Whitham et al., 2020).

79 In this context, the “genetic similarity rule” of community genetics provides a useful  
80 framework we can apply to interaction networks at the nexus of ecological and evolution-  
81 ary dynamics. In a study combining experimental common gardens and landscape-  
82 scale observations of interactions between *Populus* spp. (cottonwoods) and arthropods,  
83 Bangert et al. (2006) observed that individual genotypes that are more genetically similar  
84 will tend to have similar phytochemical traits and thus tend to have similar interactions  
85 with other species. Although this is likely to have consequences for interactions and  
86 network structure, studies in the network ecology literature generally do not include a  
87 genetic component (Lau et al., 2017) and community genetics studies have primarily  
88 focused on community composition in terms of the abundance of species (Des Roches  
89 et al., 2018). Some studies have examined the effects of genetic variation on trophic  
90 chains in plant-associated communities (including *Populus*, *Solidago*, *Oenothera*, *Salix*)  
91 (Bailey et al., 2005; Johnson, 2008; Smith et al., 2011, 2015; Barbour et al., 2016) and  
92 generally found that increasing genotypic diversity leads to increased trophic complexity.  
93 Only two other studies, that we are aware of, have explicitly examined the effect of  
94 genotypic variation on the structure of interaction networks between tree individuals and  
95 associated herbivores (Lau et al., 2016a; Keith et al., 2017) and both found that genotypic  
96 diversity generates increased network modularity (i.e., compartmentalization). However,  
97 both of these studies were examining networks at the scale of forest stands, rather  
98 than networks associated with individual trees; therefore, neither was able to observe  
99 replicated networks in order to statistically test for genetic effects on network structure  
100 and quantify the genetic component (i.e., heritable variation) in network structure.

101 Here, we investigate how genetic variation in a foundation tree species determines  
102 the structure of a network of interactions among a community of tree associated lichens.  
103 Using a long-term (20+ years), common garden experiment with clonally replicated  
104 *Populus angustifolia* individuals of known genetic identity (Martinsen et al., 2001).  
105 We focused on a community of 9 epiphytic lichen species, as previous research has  
106 demonstrated significant compositional responses of epiphytes to genotypic variation  
107 (Winfrey et al., 2011; Zytynska et al., 2011). Applying a probability-theory based  
108 network modeling approach (Araújo et al., 2011), we constructed a set of interaction  
109 network models for the lichens associated with individual trees. Using these models,  
110 we then examined the genetic basis of the structure of these ecological networks via  
111 several network metrics that measure different aspects of network structure at the scale

of individual species (i.e., nodes) or the entire network observed on each tree genotype. Given the potential importance of focal or “central” nodes (e.g., species) for determining network dynamics (Lieberman et al., 2005), we focused on network metrics that measure centrality for individual species and centralization for whole networks. Both of these metrics measure how much a species is connected in the network relative to other species. As there is a preponderance of evidence that in natural systems evolution occurs in communities comprised of networks of interacting species (Lau et al., 2016a; Keith et al., 2017; Thompson, 2013; Bascompte et al., 2006), we set out to test two hypotheses. First, per the community similarity rule (Bangert et al., 2006) and IIGE theory (Whitham et al., 2020), we hypothesize that trees of the same genotype (i.e., clones) will support more similar lichen interaction networks relative to less related genotypes. In other words, epiphytic lichen network structure is heritable, which can be calculated via comparisons of within and among group variation in network structure. Second, heritability of lichen network structure is the result of underlying phenotypic covariation in tree traits important to interactions between trees and lichens and among lichens. Evidence that such trait covariance generates variation in interactions among community members provides an intermediate genetics-based mechanism for the underlying factors determining lichen distribution and abundance. In combination, evaluating these two hypotheses is fundamental to understanding variation and dynamics of network structure and evolution.

## MATERIALS AND METHODS

### Study System

The study was conducted along the Weber River, UT (USA), which is a cottonwood (*Populus* spp.) dominated riparian ecosystem. Although two native species, *Populus angustifolia* (James) and *Populus fremontii* (S. Watson), occur here and are known to hybridize, in order to focus on intra-specific genetic variation we only sampled pure or advanced generation back-crosses of *P. angustifolia*. Bark lichens have been intensively sampled in this system and provide an ideal community in which to observe and model interaction networks, as their sessile nature permits accurate identification of individuals and their highly localized, direct contact interactions and slow population turnover rates facilitate the assessment of interactions among lichen species on individual trees (Lamit et al., 2015b).

A long-term, common garden experiment was used to isolate the effect of tree genotype from the effect of the localized microenvironment associated with each individual and spatial autocorrelation. Established in 1992, asexually propagated clones of genotyped *P. angustifolia* individuals were obtained from wild collections and planted in fully randomized design at the Ogden Nature Center, Ogden, UT. From the population of established individuals in the common garden, we chose a total of ten genotypes, replicated between 3 and 8 times each, for sampling. We selected tree genotypes that generally had lichens present in order to permit the construction of interaction networks.

### Bark Lichens and Trait Observations

On each tree, presence or absence of each lichen species was assessed in 50 total 1 cm<sup>2</sup> cells arrayed in a checkerboard pattern. Given the small size and sessile nature of lichens,

155 we were able to rapidly assess lichen interactions by quantifying thalli of different  
156 species occurring in close proximity. Sampling was restricted to the northern aspect  
157 of the trunk to maximize the abundance of lichens and control for the effect of trunk  
158 aspect. Two adjacent  $100\text{ cm}^2$  quadrats centered at 50 cm and 95 cm from ground level  
159 were sampled (Fig 1). The observed lichen community included: *Athallia holocarpa*,  
160 *Candelariella subdeflexa*, *Myriolecis hagenii*, *Rinodina freyi*, *Physcia adscendens*,  
161 *Physciella melanachra*, *Physcia undulata*, *Xanthomendoza galericulata*, *Xanthomendoza*  
162 *montana*. Several other species were not observed in the present study but are known to  
163 occur in this region: *Melanohalea elegantula*, *Melanohalea subolivacea*, *Phaeophyscia*  
164 *ciliata* and *Phaeophyscia orbicularis*.

165 The cell size and checkerboard sampling pattern was chosen to isolate the individuals  
166 in each cell. *Xanthomendoza galericulata* thallus size in this common garden, we had  
167 observed a median thallus size of  $0.12 \pm 0.001\text{ cm}^2$  (1 S.E.) (Supporting Information,  
168 Fig. 1). Based on the median thallus size, we expected thalli observed in each cell to  
169 generally be spatially independent of thalli present in other cells but exposed to similar  
170 micro-environmental conditions created by the bark and the location of the sampling area  
171 on an individual tree. Therefore, we were confident in treating the cell-wise observations  
172 in quadrats as independent with respect to lichen-lichen interactions. We quantified  
173 the texture of the bark in the quadrat is the percent of  $1\text{ cm}^2$  cells with rough bark. In  
174 addition to bark roughness, we also examined several bark chemistry traits by taking  
175 bark samples immediately adjacent to each quadrat. We used previously collected  
176 phytochemical data from Lamit et al. (2011), including the concentration of condensed  
177 tannins, carbon and nitrogen. Additionally, we quantified bark pH for each tree. Bark  
178 samples were collected in June 2011 by excavating adjacent to the quadrat down to a  
179 depth of 2 mm. Bark pieces were air dried for storage. In May of 2019, samples were  
180 processed for pH measurements by crushing with a mortar and pestle until all pieces  
181 were  $\approx 0.5\text{ cm}$  in diameter, creating equivalent surface areas among samples. 0.5 g of  
182 crushed bark was placed in a 15 ml Falcon collection tube with 5 ml of deionized water.  
183 Tubes were capped and let sit for 24 hrs prior to pH measurement. **pH instrument?**

#### 184 **Lichen Network Modeling**

185 For each tree, repeated observations of lichens were made in order to construct replicated  
186 interaction networks for each genotype. We conducted a modified sampling procedure  
187 originally developed by Lamit et al. (2015b) with the addition that we quantified the  
188 presence of lichens in the  $1\text{ cm}^2$  cells on individual trees of *P. angustifolia*. Unipartite  
189 networks were generated using the conditional probabilities of each species pair, i.e., the  
190 probability of observing one species given an observation of another species  $P(S_i|S_j)$ ,  
191 based on the method developed by Araújo et al. (2011). To calculate conditional  
192 probabilities, we quantified the individual probabilities of species occurrences  $P(S_i)$  and  
193 the joint probability of co-occurrences  $P(S_i, S_j)$  using the frequencies of each species  
194 and their co-occurrences. We were then able to calculate the conditional probabilities of  
195 each species pair as  $P(S_i|S_j) = \frac{P(S_i, S_j)}{P(S_j)}$ , based on the axioms of probability. This yielded  
196 a matrix that could possibly be asymmetric, i.e.,  $P(S_i|S_j)$  does not have to be equal to  
197  $P(S_j|S_i)$ . Another important property of this matrix is that the diagonal,  $P(S_i|S_i)$ , was  
198 equal to one for all species present and zero for species that were not observed in any  
199 cell.

200 We then applied an analytical procedure to remove non-significant links between  
 201 species. This procedure determines if the joint probability of a species pair (i.e.,  $P(S_i, S_j)$ )  
 202 is different from zero (Fig. 2). Here, a confidence interval  $CI_{95\%}$  is calculated as as  
 203  $CI_{95\%} = E[S_iS_j] * Z_{95\%} * \sqrt{V(S_iS_j)}$ , where the expected frequency of co-occurrences  
 204  $E(S_iS_j)$  is the total number of cells surveyed ( $N$ ) times the independent probabilities of  
 205 each species  $P(S_i) * P(S_j)$ ,  $Z_{95\%}$  is the Z-score for 95% from a Z-distribution and the  
 206 expected variance of  $E(S_iS_j)$  is the total number of cells times the expected probability  
 207 of  $S_iS_j$  and its compliment (i.e.,  $V(S_iS_j) = N * E[P(S_i, S_j)] * (1 - E[P(S_i, S_j)])$ ). If  
 208 the observed number of co-occurrence falls outside of the confidence interval, the  
 209 joint probability  $P(S_i, S_j)$  is determined to be equal to the product of the individual  
 210 probabilities (i.e.,  $P(S_i)P(S_j)$ ), and the conditional probability reduces to the individual  
 211 probability of that species  $P(S_i)$ . Therefore, unless the co-occurrence of a species pair  
 212 falls outside the confidence interval, the probability that the observation of one species  
 213 given the other is no different than simply observing that species alone. This enables us  
 214 to remove links from a given network by re-scaling the resulting conditional probabilities  
 215 through subtraction of the individual probabilities from the conditional probabilities (i.e.,  
 216 how different the conditional probability is from the independent probability), which  
 217 makes any species with a non-significant conditional probability zero.

218 The resulting matrix ( $\mathbf{D} = D_{ij}$ ) can be interpreted as one species' impact on another  
 219 with zero being no effect and values less than or greater than zero being negative and  
 220 positive effects, respectively. We will refer to  $\mathbf{D}$  as a signed, weighted interaction  
 221 matrix. As such,  $\mathbf{D}$  has the properties that it can be asymmetric (i.e.,  $D_{ij}$  does not  
 222 necessarily equal  $D_{ji}$ ) and it scales between -1 and 1, and, therefore, does not have the  
 223 mathematical properties of a probabilistic network (Poisot et al., 2016). Also, as the  
 224 method does not track individuals within species and interactions such as competitive  
 225 exclusion or facilitation within species would result in the same species being observed.  
 226 Therefore, the results of intra-specific interactions always results in the same species  
 227 being observed and a resulting  $D_{ii} = 0$ . In the context of these networks, asymmetry and  
 228 positive/negative valued connections are distinct quantities. In-coming and out-going  
 229 connections can be interpreted as “influenced by” and “influenced”, respectively; while  
 230 positive and negative should be seen as one species increasing or decreasing, respectively,  
 231 the probability of another species' occurrence.

## 232 Analyses, Software and Data

233 To quantify the structural variation of lichen networks we calculated several metrics at  
 234 both the level of node and whole networks. Although there are many other metrics, for  
 235 the sake of simplicity we focus on a subset that represent the primary interesting features  
 236 of network structure (see Lau et al. (2017)). We calculated the number of interactions or  
 237 “links” in each network (degree), which provides a measure of the size of the network  
 238 (Lau et al., 2016a; Borrett and Lau, 2014). We also calculated the centralization of each  
 239 network using Freeman's centrality, which measures the evenness of the distribution of  
 240 interactions among the species in the network, using the sna package (Butts, 2019).  
 241 In a network with low centralization species have similar strengths and numbers of  
 242 interactions. A network with high centralization tends to have one or small number  
 243 of species that interact with other species. We used a related function to calculate  
 244 the centrality of each species (i.e., node level centrality) in each network as well. To

245 calculate separate metrics for positive and negative links, as the networks contained not  
246 only positive and negative connections but also directional connections (both in-coming  
247 and out-going), we calculated the same network metrics for all combinations of these  
248 types of connections using recently developed methods for signed, weighted and directed  
249 networks (Everett and Borgatti, 2014) using the `signnet` package (Schoch, 2020).

250 We used a combination of parametric and non-parametric, permutation based frequent-  
251tist statistical analyses to test for the effects of genetic variation on lichen communities  
252 and their interaction networks. To assess the effect of genotype on traits as univariate re-  
253 sponse variables (including the metrics of network structure), we used additive, random  
254 effects models with Restricted Maximum Likelihood (REML) conducted in R via the  
255 `lme4` and `RLRsim` packages (Bates et al., 2015; Scheipl et al., 2008). To conform to  
256 test assumptions, traits were root transformed with the exception of condensed tannin  
257 concentration and carbon-nitrogen ratio, which were rank and  $\log_{10}$  transformed, respec-  
258 tively. Differences in node level centrality among species was tested using ANOVA and  
259 Tukey-HSD multiple comparison tests. Correlations among trait variables and network  
260 metrics were quantified and tested using linear correlations of Pearson's  $r$ . For multivari-  
261 ate response variables, such as lichen community composition and network structure,  
262 we used distance based multivariate statistical approaches. To quantify the similarity of  
263 lichen networks among individual trees, we calculated the pairwise Euclidean distance  
264 of the **D** interaction matrices among all trees (Newman, 2010). To test for the effects of  
265 genotype and other predictor variables on network similarity we conducted Permuta-  
266 tional Analysis of Variance (PERMANOVA) with `vegan` (Oksanen et al., 2019). For  
267 visualization of multivariate patterns, we used Non-metric Multi-Dimensional Scaling  
268 (NMDS) (Goslee and Urban, 2007) to produce dimensionally reduced ordinations of  
269 these multi-variate responses and fitted vectors for continuous predictor variables to  
270 the ordinated values (Oksanen et al., 2019). Using random initial configurations with a  
271 maximum of 500 iterations and a change in stress threshold of less than  $10^{-12}$ . This was  
272 repeated for one to four dimension configurations, and the configuration with the lowest  
273 dimensionality and unexplained variation less than 10% was selected. For all tests  
274 where genotype was used as a predictor, we quantified the heritability of the response  
275 variable. Because the trees in the garden were clonal replicates of each genotype, we  
276 calculated broad-sense heritability, which is the genotypic variance divided by the total  
277 phenotypic variance (Conner and Hartl, 2004), which can be interpreted as a measure of  
278 the phenotypic variance due to genotypic variation. All analyses were conducted using  
279 R version 4.0.2 (R Core Team, 2020). Code and data for the project are openly available  
280 as a reproducible workflow using `drake` (Landau, 2018), which is archived via Zenodo  
281 [zenodo.com/doi/XXXXXX](https://zenodo.com/doi/XXXXXX).

## 282 RESULTS

283 In support of our first hypotheses, we found that tree genotype influenced lichen net-  
284 work structure and that multiple lichen network metrics were heritable. Tree genotype  
285 significantly predicted the structural similarity of lichen networks and, overall, network-  
286 level metrics responded significantly to tree genotype, including network degree and  
287 centralization including both in-coming and out-going links or when separated into  
288 in-coming only or out-going only (Table 1, Fig. 4). Metrics including only positive links

also showed a significant effect of tree genotype, including positive degree and positive in-going centralization. Metrics calculated with negative links were not significant, including degree (negative) and both in-coming (negative) and out-going centralization (negative).

	Response	df	RLRT	H2	p-value
Lichen Network Similarity		9	3.5821	0.41	0.0537
Degree		9	3.5175	0.32	0.0255
Degree (positive)		9	3.6925	0.32	0.0229
Degree (negative)		9	0.0327	0.03	0.3859
Centralization		9	4.0444	0.33	0.0184
Centralization In-Degree		9	4.4812	0.35	0.0142
Centralization In-Degree (positive)		9	3.9852	0.33	0.0190
Centralization In-Degree (negative)		9	0.3304	0.11	0.2508
Centralization Out-Degree		9	3.8615	0.32	0.0205
Centralization Out-Degree (positive)		9	3.5585	0.31	0.0248
Centralization Out-Degree (negative)		9	0.0862	0.05	0.3446

**Table 1.** Genotypic effects on the associated lichen network structure. *RLRT* is the statistic from the restricted likelihood ratio tests.

The genetic response of network centralization was driven by variation in *Athallia holocarpa*. Centrality varied significantly among species ( $F_{8,324} = 7.99$ ,  $R^2 = 0.16$ ,  $p\text{-value} < 0.0001$ ). *Athallia holocarpa* was the main species to exhibit a significant response to tree genotype in terms of positive centrality for both the in-coming ( $RLRT = 3.61$ ,  $H^2 = 0.32$ ,  $p\text{-value} = 0.0240$ ) and out-going ( $RLRT = 3.13$ ,  $H^2 = 0.30$ ,  $p\text{-value} = 0.0327$ ) perspectives, but not for either negative centrality metrics in-coming ( $RLRT = 0$ ,  $H^2 = 0$ ,  $p\text{-value} = 1$ ) or out-going ( $RLRT = 0$ ,  $H^2 = 0$ ,  $p\text{-value} = 0.4543$ ). None of the other species' centralities showed a genotypic response (Supporting Information, Fig. 2) with the exception of *X. montana* ( $RLRT = 2.92$ ,  $H^2 = 0.32$ ,  $p\text{-value} = 0.0375$ ); however, the centrality of *X. montana* was much lower overall relative to *A. holocarpa* and the variation in *X. montana* centrality was restricted to two genotypes (Fig. 5).

In support of our second hypothesis, analysis of trait covariation revealed that genotype indirectly influenced lichen network centralization via genetically based variation in bark roughness. The percent cover of rough bark ( $RLRT = 4.8526$ ,  $H^2 = 0.3221$ ,  $p\text{-value} = 0.0113$ ) and condensed tannins ( $RLRT = 3.0522$ ,  $H^2 = 0.3205$ ,  $p\text{-value} = 0.0343$ ) both displayed significant responses to tree genotype. None of the other bark traits, pH ( $RLRT = 0.00$ ,  $H^2 = 0.00$ ,  $p\text{-value} = 1.0000$ ) or carbon-nitrogen ratio ( $RLRT = 0.0000$ ,  $H^2 = 0.0000$ ,  $p\text{-value} = 1.0000$ ), showed a significant response to tree genotype and none other than bark roughness was correlated with network similarity (Table 2); therefore, we focused our subsequent analyses on the indirect effect of genotype on lichen network structure via bark roughness. We found that bark roughness was significantly correlated with network similarity and other lichen network metrics, including negative correlations with overall network degree ( $df = 35$ ,  $t = -2.13$ ,  $r = -0.34$ ,  $p\text{-value} = 0.04$ ) and centralization ( $df = 35$ ,  $t = -2.52$ ,  $r = -0.39$ ,  $p\text{-value} = 0.02$ ). In other words, trees with more similar levels of bark roughness tended to have lichen interaction networks

318 with similar structure. To quantify the genetic bases of this effect of bark roughness on  
 319 network structure, we used the residual values from regressions of network degree and  
 320 centralization in tests of the effect of tree genotype and found no significant effect of tree  
 321 genotype for either degree ( $RLRT = 0.00$ ,  $H^2 = 0.00$ ,  $p\text{-value} = 1.0000$ ) or centralization  
 322 ( $RLRT = 0.00$ ,  $H^2 = 0.00$ ,  $p\text{-value} = 1.0000$ ), suggesting that the observed relationship  
 323 between bark roughness and lichen network structure was largely genetically based  
 324 (Fig. 6).

	df	SS	R2	Pseudo-F	p-value
Bark Roughness	1	20850.09	0.26	12.9234	0.0101
Condensed Tannins	1	5993.66	0.07	3.7150	0.0813
pH	1	1273.19	0.02	0.7892	0.3712
Carbon:Nitrogen Ratio	1	3896.18	0.05	2.4150	0.1890
Residual	32	51627.33	0.64		
Total	36	80993.59	1.00		

**Table 2.** PERMANOVA Pseudo-F Table of lichen network similarity response to bark traits.

## 325 **DISCUSSION**

326 Subsection changes proposed by Tom

- 327     • Current

- 328         – Implications of Ecological Network Heritability
- 329         – Evolution and Genetically Based Network Structure
- 330         – Conclusion

- 331     • TGW

- 332         – Importance of Network Heritability
- 333         – Networks and Levels of Selection in Community Evolution
- 334         – Conclusions

335 **START TGW**

336 DISCUSSION (omitted the intro para and moved to conclusions)

337 Importance of Network Heritability

338 Although previous studies have examined aspects of networks, such as trophic 340  
 339 complexity (Barbour et al., 2016) and forest stand-level interaction network structure  
 340 341 (Lau et al., 2016b; Keith et al., 2017), this is the first study that we are aware  
 341 of to 342 examine the heritability of network structure with replicated networks at  
 342 the genotype 343 scale. Previous work in the evolution of ecological networks have

343 primarily focused on 344 macro-evolutionary dynamics (Rezende et al., 2007; Weber et  
344 al., 2017; Valverde et al., 345 2018; Harmon et al., 2019) or have been simulation based  
345 individual-level models that 346 integrate intraspecific variation to the species level  
346 (Maliet et al., 2020), even though 347 recent syntheses have pointed to the importance  
347 of processes operating across scales of 348 organization (Guimarāes, 2020). There are  
348 two important functional ramifications of 349 genetically based variation in network  
349 structure. First, Second, Furthermore,

350 Networks and Levels of Selection in Community Evolution

### 351 **START MKL**

352 Per general systems theory, ecosystems are generally:

- 353 • self-organizing: an effect is its own cause (autocatalytic, cycles)
- 354 • holarchic: spatially and temporally nested, hierarchical sub-systems
- 355 • open: energy and matter passes across the system boundary

356 The holarchic nature provides a means for superior systems, i.e. the foundation  
357 species, to influence the sub-systems, i.e. the lichen networks, by altering the success  
358 and survival of individuals and/or their traits, both of which can alter the structure of  
359 interaction networks.

360 Evolutionarily, frequencies of traits (or genotypes) can increase or decrease in their  
361 moment of central tendency, spread, and/or kurtosis.

### 362 **END MKL**

363 Multilevel selection theory argues that selection can occur simultaneously at individ-  
364 ual, group and community levels and the demonstration of evolution at any scale requires  
365 demonstrating three key elements (review by Whitham et al. 2020). The community net-  
366 works in our study are based upon the interactions of the lichen community in which the  
367 differential survival and performance of individual tree genotypes will simultaneously  
368 result in selection occurring on the lichen community it supports. (We could introduce  
369 holobiont theory here). First, for selection to occur at the community level, there must  
370 be variation in the structure (composition, 426 abundance, species interactions, diversity,  
371 networks) of communities across the tree land427 scape. Second, these differences  
372 must be genetically based and heritable in which 428 community structure is passed  
373 from one tree generation to the next. For example, numerous 429 studies show that  
374 related individuals tend to support the same communities of insects 430 and microbes,  
375 and ecosystem processes of biodiversity, nutrient cycling and stabil431 ity, whereas  
376 unrelated individuals support more different communities and ecosystem 432 processes.  
377 Importantly, the current 433 study shows that networks are also heritable traits that  
378 greatly increases its utility as a 434 community phenotype that selection can act upon.  
379 Third, selection must act on these 435 differences to favor some communities over others  
380 leading to change over time

381 (i.e., 436 community evolution). Since our findings show genetic variation in lichen  
382 networks and show that lichen networks are heritable, future studies should evaluate  
383 how networks might change over time in response 438 to an invasive species, climate  
384 change, or some other agent of selection. For example, in this study, when a tree dies  
385 due to drought, its community network of interacting lichens is selected against, while

386 those networks of other trees that survive persist. Thus, if drought related mortality is  
387 non-random as shown by Sthultz et al. (2009), network structure can evolve via selection  
388 acting on whole communities.

389           439 In the present study, intraspecific, genotypic diversity in the heritable  
390 bark traits of individual trees appear to be a major causal factor resulting in the creation  
391 of lichen meta-communities ....

392           Conclusions – I think this intro to the Discussion is a better conclusion section could  
393 be merged with the your old conclusions section. I suggest that it be more focused  
394 with key numbered points. We found support for both of our hypotheses. First, tree  
395 genotype influenced the network 323 structure of lichen communities associated with  
396 narrowleaf cottonwoods in a riparian 324 forest ecosystem. Network similarity and  
397 metrics of network structure tended to be more 325 similar on trees of the same genotype.  
398 Generally, this genetic effect was manifested 326 in positive interactions and largely  
399 driven by *A. holocarpa*. (Not sure what the broader point of this sentence on *A. holocarpa*  
400 is as it appears to weaken a community perspective?? Need to clarify. Second, the  
401 genetically 327 based trait, bark roughness, was observed to affect network variation,  
402 largely via shifts 328 in positive in-coming and out-going interactions. Chemistry traits,  
403 whether genetically 329 based (e.g., tannin concentration) or not, were not significantly  
404 correlated with lichen

405           330 network structure. Bark roughness has been demonstrated previously to be  
406 under strong 331 genetic control (Bdeir et al., 2017), and bark roughness has also been  
407 shown to be an 332 important tree trait influencing bark lichens (Lamit et al., 2015b);  
408 however this is the 333 first demonstration of a link from genetics to lichen network  
409 structure. As such, these 334 results have important implications for the influence of  
410 genetically based variation in 335 ecosystems with networks of interacting species.

411           **END TGW**

412           We found support for both of our hypotheses. First, tree genotype influenced the  
413 network structure of lichen communities associated with narrowleaf cottonwoods in a  
414 riparian forest ecosystem. Network similarity and metrics of network structure tended  
415 to be more similar on trees of the same genotype. Generally, this genetic effect was  
416 manifested in positive interactions and largely driven by *A. holocarpa*. Second, the  
417 genetically based trait, bark roughness, was observed to affect network variation, largely  
418 via shifts in positive in-coming and out-going interactions. Chemistry traits, whether  
419 genetically based (e.g., tannin concentration) or not, were not significantly correlated  
420 with lichen network structure. Bark roughness has been demonstrated previously to be  
421 under strong genetic control (Bdeir et al., 2017), and bark roughness has also been shown  
422 to be an important tree trait influencing bark lichens (Lamit et al., 2015b); however this  
423 is the first demonstration of a link from genetics to lichen network structure. As such,  
424 these results have important implications for the influence of genetically based variation  
425 in ecosystems with networks of interacting species.

426           **Implications of Ecological Network Heritability**

427           Significant heritability of lichen interaction network structure is in line with the genetic  
428 similarity rule, networks observed on trees of the same genotype tended to be structurally  
429 similar. Although previous studies have examined aspects of networks, such as trophic  
430 complexity (Barbour et al., 2016) and forest stand-level interaction network structure

(Lau et al., 2016b; Keith et al., 2017), this is the first study that we are aware of to examine the heritability of network structure with replicated networks at the genotype scale. Previous work in the evolution of ecological networks have primarily focused on macro-evolutionary dynamics (Rezende et al., 2007; Weber et al., 2017; Valverde et al., 2018; Harmon et al., 2019) or have been simulation based individual-level models that integrate intraspecific variation to the species level (Maliet et al., 2020), even though recent syntheses have pointed to the importance of processes operating across scales of organization (Guimarães, 2020). There are two important functional ramifications of genetically based variation in network structure.

First, heritability of network structure suggests that some amount of interaction network complexity is determined and therefore could be predicted by genetic identity. Variation in space and time create variation in ecological networks that influences evolutionary dynamics via shifts in ecological dynamics, such as population demographics (Guimarães, 2020). Given that ecosystems are comprised of hundreds and thousands of species, each having a multitude of interactions, the potential to find traction for making predictions in the context of ecological, let alone evolutionary, dynamics seems daunting. The promise of predictability lies in the presence of asymmetries in ecosystems, such as hierarchy created by foundation species via differences in body size and/or life-history strategies (Ellison et al., 2005). The second is that heritability (i.e., genetic determination) means that there is structure in the spatial or temporal variation that is created by individuals of foundation species whose traits are in part determined by underlying trait differences. Although this variation is inherently a function of both genetic and environmental effects (Conner and Hartl, 2004), the community and network-level effects are also a function of the scale of the interaction (Shuster et al., 2006).

Second, even if the composition of the communities is the same among individuals and genotypes, interactions may not be. We didn't observe compositional differences using the same data from which the lichen networks were derived. If we only had our composition dataset from this study, we would have concluded no response of the lichen community to tree genotype, even though the underlying interactions among lichen species does vary among genotypes. As such differences in network structure could occur without observable differences in species richness or community composition, which have been the primary focus of almost all previous community genetics studies (Des Roches et al., 2018). Community composition of lichens has previously been observed to be different among tree genotypes in the same experimental garden (Lamit et al., 2011, 2015b). The different results observed in the present study is likely a result of differences in lichen quantification and the tree genotypes observed leading to overall higher abundances of observed lichens to assure the possibility of observing lichen interactions. The previous study used a visual percent cover estimation, unlike the current study, which observed lichens at the scale of 1 cm<sup>2</sup> cells, which could over-estimate cover depending on the frequency at which actual thallus size was less than 1 cm<sup>2</sup>, as well as both the northern and southern aspects of each tree. These differences do not negate the findings of either study. The present study's finding of differences in network structure without significant compositional differences points to the importance of quantifying how network structure changes in response to genetic variation in order to fully understand evolutionary dynamics in complex communities. Having not observed a compositional effect of tree genotype without measuring the

477 network structure could lead to the conclusion of no genetic effect on the community,  
478 even though differences in network structure are leading to altered, local evolutionary  
479 dynamics. It is possible that these underlying differences in interactions among lichens  
480 could lead to differences in community composition at a future point in time via their  
481 effects on species abundances (Shuster et al., 2006); however, this is not needed for  
482 evolutionary dynamics to occur via selection that leads to shifts in trait distributions  
483 without shifting species abundance distributions, which is possible under stabilizing,  
484 disruptive and directional selection (Conner and Hartl, 2004), so long as the relative  
485 abundances of each species is imperceptibly changed. Thus, it is imperative that further  
486 community genetics research assess or at least be aware of the potential effects of  
487 variation in interactions and not just observe species abundances, otherwise community  
488 level genetic effects may be underestimated, especially when cumulative interaction  
489 effects are taken into account (Borrett et al., 2007, 2010).

490 Furthermore, the demonstration of the heritability of interaction networks, without  
491 significant differences in community composition, provides clear empirical evidence  
492 that variation in network structure points to the need to expand IIGEs to encompass  
493 the structure of interaction networks. Although IIGE theory provides a quantitative  
494 framework within which to approach evolutionary theory at higher levels of biological  
495 organization (from populations to communities and ecosystems), this theory has focused  
496 on modeling the strong effects of foundation species (Shuster et al., 2006; Whitham  
497 et al., 2012, 2020) and has not yet integrated developments from the ecological or  
498 evolutionary network theory literature. Thus, it has not developed a way to examine  
499 complex interactions among species; however, previous studies have demonstrated  
500 this network context is likely to be important, as altering the structure of interaction  
501 networks provides a means for genetic effects to be dampened or magnified within  
502 the system of interacting species (Smith et al., 2011; Keith et al., 2017). Although  
503 such a synthesis necessitates a much greater effort than can be afforded in this paper,  
504 it is possible to point to several productive pathways forward. In terms of interaction  
505 networks, foundation species are relatively central within the system of interactions, that  
506 is their direct and/or indirect effects are greater than other species. So, when the more  
507 centralized (foundation) species have genetically based interactions, genetic effects will  
508 tend to be propagated and possibly magnified in the community. Here, we found that  
509 even though more abundant or more centralized (i.e., “important”) species were present  
510 in the community, their effects were not the main component responding to genetic  
511 effects. Considering the impact of network structure would be a productive path forward  
512 for the theoretical development and application of the IIGE concept.

### 513 **Evolution and Genetically Based Network Structure**

514 The demonstration of evolution at any scale of biological organization requires demon-  
515 strating three key elements. First, there must be variation in the structure (composition,  
516 abundance, species interactions, diversity, networks) of communities across the land-  
517 scape. Second, these differences must be genetically based and heritable in which  
518 community structure is passed from one generation to the next. For example, numerous  
519 studies show that related individuals tend to support the same communities of insects  
520 and microbes, and ecosystem processes of biodiversity, nutrient cycling and stabil-  
521 ity, whereas unrelated individuals support more different communities and ecosystem

processes (Des Roches et al., 2018; Whitham et al., 2020). Importantly, the current study shows that networks are also heritable traits that greatly increases its utility as a community phenotype that selection can act upon. Third, selection must act on these differences to favor some communities over others leading to change over time (i.e., community evolution). Since our findings show that networks are heritable, another metric of community evolution is showing how networks change over time in response to an invasive species, climate change, or some other agent of selection.

Intra-specific, genotypic diversity could be creating lichen meta-communities on individual trees that form interaction modules with different dynamics. When communities are comprised of individuals whose habitat is primarily determined by another organism, these communities inherently form modules within the larger ecosystem, as they tend to interact more with each other than with other individuals (Lau et al., 2017). Our study demonstrates that the environmental differences determined by the genetic variation within a single species can not only impact community composition, as repeatedly demonstrated in other community genetics studies (Whitham et al., 2006; Des Roches et al., 2018), but also shape the structure of interactions among individuals. Some network structures are likely to be more stable, either in response to disturbance or via self-organized dynamics. For example, centralized networks, although more efficient, are theorized to be more susceptible to targeted “attacks” in the terminology of defense networks. As mentioned previously, one class of networks that are theorized to have amplifying effects on networks have centralized “star” shapes with one or a few species at the center and radiating interactions out from the central core (Lieberman et al., 2005). This is structurally what we have observed with the networks that tend to occur on some of the genotypes in our study, i.e., the more centralized networks. It is likely that these networks could function as hot-spots of evolutionary dynamics resulting from the amplifying effect the centralized network structure found on that tree genotype, as multiple studies have found significant impacts of the removal of foundation species in different systems (Keith et al., 2017; Des Roches et al., 2018).

Ecological network studies have focused on asymmetry and the quantification of its structure in communities. The impacts of asymmetry on evolution from community dynamics have primarily produced qualitative discussion (Bascompte et al., 2006; Díaz-Castelazo et al., 2010; Guimarães et al., 2011; Thompson, 2013). More specific predictions can be found in applications of evolutionary game theory, and although developed at the population scale, such theory can apply to communities (Lieberman et al., 2005). One seemingly useful direction is the classification of networks into two general categories, rooted and cyclic, in which rooted networks have interactions in which evolutionary effects emanate from one or multiple origins but these effects do not have feedbacks to the origin, whereas cyclic networks contain feedbacks to one or more origins. This is equivalent to “unidirectional” and “reciprocal” genetic effects in the context of IIGE theory (Whitham et al., 2020). As we do not have an estimate of the effect of the lichen on the fitness of the tree they occur on, we can not determine whether the lichen networks in this system are cyclic or not. In terrestrial ecosystems, lichens play important ecological roles, such as substrate stabilization (Root et al., 2011) and nitrogen fixation (Nelson et al., 2018). Some epiphytic lichens can have demonstrable effects on the availability of nutrients for the trees that they are associated with (Norby and Sigal, 1989). Although none of the lichens the present study’s system is known to

568 fix nitrogen, it is possible that they might add micro-nutrients or provide some other  
569 unobserved benefit to their host trees. Elucidating the presence of and quantifying  
570 such feedbacks would allow for the determination of the cyclic nature and potential  
571 evolutionary dynamics.

572 Since lichen individuals are multi-species complexes, there is also the potential for  
573 evolutionary dynamics to shift within the context of the lichen symbiosis. There is  
574 substantial evidence that lichens have served as the “cradle of symbiotrophic fungal  
575 diversification” (Arnold et al., 2009) and recent research has shown significant net-  
576 work structure of endolichenic fungi and lichens collected from across North America  
577 (Chagnon et al., 2016). Analysis of the structure of ecological networks has generally  
578 supported the conclusion that nestedness, or the degree to which species tend to inter-  
579 act with similar subsets of the community, tends to promote stability in mutualistic,  
580 primarily bipartite (i.e., two-mode), networks and modularity contributes to the stabi-  
581 lization of antagonistic networks (Elias et al., 2013; Grilli et al., 2016). Although there  
582 is growing evidence that the nestedness of mutualistic networks is not necessarily the  
583 result of selection for systems-level properties that promote stability but could be either  
584 product of asymptotic abundance distributions leading to uneven interaction frequencies  
585 (Staniczenko et al., 2013) and/or a by-product of selection and divergence creating  
586 network “spandrels” in ecosystems (Valverde et al., 2018), this does not preclude the  
587 functional consequences of network structure but rather the developmental or evolu-  
588 tionary processes that have produced the structure. In the present study, we did not  
589 examine nestedness or modularity of the lichen networks as we could not find metrics for  
590 analyzing networks that are not only weighted and directed but also signed. Hopefully  
591 future network theoretic developments will make the appropriate metrics available to  
592 conduct these analyses.

## 593 Conclusion

594 In the face of the high degree of complexity and potential context dependency of  
595 ecological processes, the current study points to the utility of considering the spatial and  
596 temporal scales of interactions, as discussed in previous studies (Bangert et al., 2006;  
597 Zook et al., 2010; Zytnyska et al., 2012). In the present research, we found that the  
598 assembly of ecological networks can have a measurable genetic basis depending on  
599 the spatial scale of interactions, due in part, to asymmetries in size and longevity of  
600 organisms. The importance of the scale of network organization to create hierarchical  
601 structure (Guimarães, 2020) and the potential for foundation species to create this  
602 structure in the vast majority of ecosystems (Ellison et al., 2005; Whitham et al., 2006)  
603 suggests that future work would be aided by determining these modules that include  
604 species with large differences in body-size and longevity. As heritable variation is the  
605 raw material for natural selection to act upon, a genetic basis for interaction network  
606 structure indicates that conserving genetic variation is important to consider in efforts  
607 to restore or preserve complex species interactions and their associated ecosystem  
608 functions (Whitham et al., 2012; Evans et al., 2013; Whitham et al., 2020). Going  
609 forward, future work could extend the many previous community genetics studies that  
610 have focused on sessile organisms, such as galling insects (Bailey et al., 2005; Whitham  
611 et al., 2006; Crutsinger et al., 2014; Smith et al., 2011; Keith et al., 2017), to quantify  
612 the frequency of these interactions in the context of the larger community. Network

613 modeling and analysis will provide useful tools for the identification of species within  
614 network modules that are most important to study in systems where little is known about  
615 the natural history of organisms in an ecosystem is lacking. Such investigations will  
616 bring us closer to understanding the evolutionary drivers of Darwin's entangled bank  
617 and the interconnectedness of species in complex communities (Darwin, 1859; Dátilo  
618 et al., 2016).

## 619 Acknowledgments

620 This work was supported by the National Science Foundation grant (DEB-0425908) and  
621 Integrative Graduate Research Traineeship (IGERT) fellowships for M.L. and L.L. The  
622 Ogden Nature Center staff helped to maintain the common gardens. Lichen sampling  
623 was supported by Todd Wojtowicz, Luke Evans and David Solance Smith.

## 624 REFERENCES

- 625 Araújo, M. B., Rozenfeld, A., Rahbek, C., and Marquet, P. A. (2011). Using species  
626 co-occurrence networks to assess the impacts of climate change. *Ecography*, 34:897–  
627 908.
- 628 Arnold, A. E., Miadlikowska, J., Higgins, K. L., Sarvate, S. D., Gugger, P., Way, A.,  
629 Hofstetter, V., Kauff, F., and Lutzoni, F. (2009). A phylogenetic estimation of trophic  
630 transition networks for ascomycetous Fungi: Are lichens cradles of symbioticrophic  
631 Fungal diversification? *Systematic Biology*, 58(3):283–297.
- 632 Bailey, J. K., Schweitzer, J. A., Ubeda, F., Koricheva, J., LeRoy, C. J., Madritch, M. D.,  
633 Rehill, B. J., Bangert, R. K., Fischer, D. G., Allan, G. J., and Whitham, T. G. (2009).  
634 From genes to ecosystems: a synthesis of the effects of plant genetic factors across  
635 levels of organization. *Philosophical transactions of the Royal Society of London.  
Series B, Biological sciences*, 364(1523):1607–16.
- 636 Bailey, J. K., Wooley, S. C., Lindroth, R. L., and Whitham, T. G. (2005). Importance  
637 of species interactions to community heritability: a genetic basis to trophic-level  
638 interactions. *Ecology Letters*, 0(0):051122062725008.
- 639 Bangert, R. K., Turek, R. J., Rehill, B., Wimp, G. M., Schweitzer, J. A., Allan, G. J.,  
640 Bailey, J. K., Martinsen, G. D., Keim, P., Lindroth, R. L., and Whitham, T. G. (2006).  
641 A genetic similarity rule determines arthropod community structure. *Molecular  
ecology*, 15:1379–1391.
- 642 Barbour, M. A., Fortuna, M. A., Bascompte, J., Nicholson, J. R., Julkunen-Tiitto, R.,  
643 Jules, E. S., and Crutsinger, G. M. (2016). Genetic specificity of a plant-insect food  
644 web: Implications for linking genetic variation to network complexity. *Proceedings of  
the National Academy of Sciences of the United States of America*, 113(8):2128–2133.
- 645 Barbour, R. C., O'Reilly-Wapstra, J. M., Little, D. W. D., Jordan, G. J., Steane, D. A.,  
646 Humphreys, J. R., Bailey, J. K., Whitham, T. G., Potts, B. M., De Little, D. W., Jordan,  
647 G. J., Steane, D. A., Humphreys, J. R., Bailey, J. K., Whitham, T. G., and Potts, B. M.  
648 (2009). A geographic mosaic of genetic variation within a foundation tree species  
649 and its community-level consequences. *Ecology*, 90(7):1762–1772.
- 650 Bascompte, J., Jordano, P., and Olesen, J. M. (2006). Asymmetric Coevolutionary  
651 Networks Facilitate Biodiversity Maintenance \r10.1126/science.1123412. *Science*,  
652 312:431–433.


- 656 Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects  
657 models using lme4. *Journal of Statistical Software*, 67(1):1–48.
- 658 Bdeir, R., Muchero, W., Yordanov, Y., Tuskan, G. A., Busov, V., and Gailing, O. (2017).  
659 Quantitative trait locus mapping of Populus bark features and stem diameter. *BMC  
660 Plant Biology*.
- 661 Borrett, S. R., Fath, B. D., and Patten, B. C. (2007). Functional integration of ecological  
662 networks through pathway proliferation. *Journal of Theoretical Biology*, 245(1):98–  
663 111.
- 664 Borrett, S. R. and Lau, M. K. (2014). enaR: An R package for Ecosystem Network  
665 Analysis. *Methods in Ecology and Evolution*, 5(11):1206–1213.
- 666 Borrett, S. R., Whipple, S. J., and Patten, B. C. (2010). Rapid development of indirect  
667 effects in ecological networks. *Oikos*, 119(7):1136–1148.
- 668 Busby, P. E., Lamit, L. J., Keith, A. R., Newcombe, G., Gehring, C. A., Whitham, T. G.,  
669 and Dirzo, R. (2015). Genetics-based interactions among plants, pathogens, and  
670 herbivores define arthropod community structure. *Ecology*, 96(7):1974–1984.
- 671 Butts, C. T. (2019). *sna: Tools for Social Network Analysis*. R package version 2.5.
- 672 Chagnon, P. L., U'Ren, J. M., Miadlikowska, J., Lutzoni, F., and Elizabeth Arnold,  
673 A. (2016). Interaction type influences ecological network structure more than local  
674 abiotic conditions: evidence from endophytic and endolichenic fungi at a continental  
675 scale. *Oecologia*, 180(1):181–191.
- 676 Conner, K. and Hartl, D. L. (2004). *A Primer of Ecological Genetics: a textbook*.  
677 Sinauer Associates, Inc., Sunderland, Massachusetts, 1st edition.
- 678 Crutsinger, G. M. (2016). A community genetics perspective: Opportunities for the  
679 coming decade. *New Phytologist*.
- 680 Crutsinger, G. M., Rudman, S. M., Rodriguez-Cabal, M. A., McKown, A. D., Sato,  
681 T., MacDonald, A. M., Heavyside, J., Gerald, A., Hart, E. M., LeRoy, C. J., and  
682 El-Sabaawi, R. W. (2014). Testing a ‘genes-to-ecosystems’ approach to understanding  
683 aquatic-terrestrial linkages. *Molecular Ecology*, 23(23):5888–5903.
- 684 Darwin, C. (1859). *On the Origin of Species*. Murray, London.
- 685 Dátillo, W., Lara-Rodríguez, N., Jordano, P., Guimarães, P. R., Thompson, J. N., Marquis,  
686 R. J., Medeiros, L. P., Ortiz-Pulido, R., Marcos-García, M. A., and Rico-Gray,  
687 V. (2016). Unravelling Darwin’s entangled bank: Architecture and robustness of  
688 mutualistic networks with multiple interaction types. *Proceedings of the Royal Society  
689 B: Biological Sciences*.
- 690 Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison,  
691 M. T., Schweitzer, J. A., and Palkovacs, E. P. (2018). The ecological importance of  
692 intraspecific variation. *Nature Ecology and Evolution*, 2(1):57–64.
- 693 Díaz-Castelazo, C., Guimarães, P. R., Jordano, P., Thompson, J. N., Marquis, R. J., and  
694 Rico-Gray, V. (2010). Changes of a mutualistic network over time: Reanalysis over a  
695 10-year period. *Ecology*.
- 696 Elias, M., Fontaine, C., and Frank Van Veen, F. J. (2013). Evolutionary history and  
697 ecological processes shape a local multilevel antagonistic network. *Current Biology*,  
698 23(14):1355–1359.
- 699 Ellison, A. M., Bank, M. S., Clinton, B. D., Colburn, E. A., Elliott, K., Ford, C. R.,  
700 Foster, D. R., Kloeppe, B. D., Knoepp, J. D., Lovett, G. M., Mohan, J., Orwig,  
701 D. A., Rodenhouse, N. L., Sobczak, W. V., Stinson, K. A., Stone, J. K., Swan, C. M.,

- 702 Thompson, J., Von Holle, B., and Webster, J. R. (2005). Loss of foundation species:  
703 consequences for the structure and dynamics of forested ecosystems. *Frontiers in*  
704 *Ecology and the Environment*, 3(9):479–486.
- 705 Evans, D. M., Pocock, M. J. O., and Memmott, J. (2013). The robustness of a network  
706 of ecological networks to habitat loss. *Ecology letters*, 16:844–52.
- 707 Everett, M. G. and Borgatti, S. P. (2014). Networks containing negative ties. *Social*  
708 *Networks*, 38(1):111–120.
- 709 Fath, B. D. and Patten, B. C. (1998). Network synergism: Emergence of positive  
710 relations in ecological systems. *Ecological Modelling*, 107(2):127 – 143.
- 711 Gehring, C., Flores-Rentería, D., Sthultz, C. M., Leonard, T. M., Flores-Rentería,  
712 L., Whipple, A. V., and Whitham, T. G. (2014). Plant genetics and interspecific  
713 competitive interactions determine ectomycorrhizal fungal community responses to  
714 climate change. *Molecular Ecology*, 23(6):1379–1391.
- 715 Gehring, C. A., Sthultz, C. M., Flores-Rentería, L., Whipple, A. V., and Whitham, T. G.  
716 (2017). Tree genetics defines fungal partner communities that may confer drought  
717 tolerance. *Proceedings of the National Academy of Sciences*.
- 718 Gómez, J. M., Abdelaziz, M., Camacho, J. P. M., Muñoz-Pajares, A. J., and Perfectti, F.  
719 (2009). Local adaptation and maladaptation to pollinators in a generalist geographic  
720 mosaic. *Ecology Letters*, 12(7):672–682.
- 721 Goslee, S. C. and Urban, D. L. (2007). The ecodist package for dissimilarity-based  
722 analysis of ecological data. *Journal of Statistical Software*, 22:1–19.
- 723 Grilli, J., Rogers, T., and Allesina, S. (2016). Modularity and stability in ecological  
724 communities. *Nature Communications*, 7.
- 725 Guimarães, P. R. (2020). The Structure of Ecological Networks Across Levels of  
726 Organization. *Annual Review of Ecology, Evolution, and Systematics*, 51(1).
- 727 Guimarães, P. R., Jordano, P., and Thompson, J. N. (2011). Evolution and coevolution  
728 in mutualistic networks. *Ecology letters*, 14(9):877–85.
- 729 Guimarães, P. R., Rico-Gray, V., Oliveira, P. S., Izzo, T. J., dos Reis, S. F., and Thompson,  
730 J. N. (2007). Interaction Intimacy Affects Structure and Coevolutionary Dynamics in  
731 Mutualistic Networks. *Current Biology*, 17(20):1797–1803.
- 732 Harmon, L. J., Andreazzi, C. S., Débarre, F., Drury, J., Goldberg, E. E., Martins,  
733 A. B., Melián, C. J., Narwani, A., Nuismer, S. L., Pennell, M. W., Rudman, S. M.,  
734 Seehausen, O., Silvestro, D., Weber, M., and Matthews, B. (2019). Detecting the  
735 macroevolutionary signal of species interactions. *Journal of Evolutionary Biology*,  
736 32(8):769–782.
- 737 Holeski, L. M., Hillstrom, M. L., Whitham, T. G., and Lindroth, R. L. (2012). Relative  
738 importance of genetic, ontogenetic, induction, and seasonal variation in producing a  
739 multivariate defense phenotype in a foundation tree species. *Oecologia*, 170:695–707.
- 740 Johnson, M. T. J. (2008). Bottom-up effects of plant genotype on aphids, ants, and  
741 predators. *Ecology*.
- 742 Keith, A. R., Bailey, J. K., Lau, M. K., and Whitham, T. G. (2017). Genetics-based inter-  
743 actions of foundation species affect community diversity, stability and network struc-  
744 ture. *Proceedings of the Royal Society B: Biological Sciences*, 284(1854):20162703.
- 745 Lamit, L., Bowker, M., Holeski, L., Næsborg, R. R., Wooley, S., Zinkgraf, M., Lindroth,  
746 R., Whitham, T., and Gehring, C. (2011). Genetically-based trait variation within a  
747 foundation tree species influences a dominant bark lichen. *Fungal Ecology*, 4(1):103–


- 748 109.
- 749 Lamit, L. J., Busby, P. E., Lau, M. K., Compson, Z. G., Wojtowicz, T., Keith, A. R.,  
750 Zinkgraf, M. S., Schweitzer, J. A., Shuster, S. M., Gehring, C. A., and Whitham, T. G.  
751 (2015a). Tree genotype mediates covariance among communities from microbes to  
752 lichens and arthropods. *Journal of Ecology*, 103(4):840–850.
- 753 Lamit, L. J., Lau, M. K., Næsborg, R. R., Wojtowicz, T., Whitham, T. G., and Gehring,  
754 C. A. (2015b). Genotype variation in bark texture drives lichen community assembly  
755 across multiple environments. *Ecology*, 96(4):960–971.
- 756 Landau, W. M. (2018). The drake r package: a pipeline toolkit for reproducibility and  
757 high-performance computing. *Journal of Open Source Software*, 3(21).
- 758 Lau, M. K., Borrett, S. R., Baiser, B., Gotelli, N. J., and Ellison, A. M. (2017). Ecological  
759 network metrics: opportunities for synthesis. *Ecosphere*, 8(8):e01900.
- 760 Lau, M. K., Keith, A. R., Borrett, S. R., Shuster, S. M., and Whitham, T. G. (2016a).  
761 Genotypic variation in foundation species generates network structure that may drive  
762 community dynamics and evolution. *Ecology*, 97(3):15–0600.
- 763 Lau, M. K., Keith, A. R., Borrett, S. R., Shuster, S. M., and Whitham, T. G. (2016b).  
764 Genotypic variation in foundation species generates network structure that may drive  
765 community dynamics and evolution. *Ecology*.
- 766 Lieberman, E., Hauert, C., and Nowak, M. A. (2005). Evolutionary dynamics on graphs.  
767 *Nature*, 433(7023):312–316.
- 768 Lieberman, E., Hauert, C., and Nowak, M. A. (2005). Evolutionary dynamics on graphs.  
769 *Nature*, 433(7023):312–316.
- 770 Maliet, O., Loeuille, N., and Morlon, H. (2020). An individual-based model for  
771 the eco-evolutionary emergence of bipartite interaction networks. *Ecology Letters*,  
772 23(11):1623–1634.
- 773 Martinsen, G. D., Whitham, T. G., Turek, R. J., and Keim, P. (2001). Hybrid populations  
774 selectively filter gene introgression between species. *Evolution*, 55(7):1325–1335.
- 775 Moya-Laraño, J. (2011). Genetic variation, predator-prey interactions and food web  
776 structure. *Philosophical transactions of the Royal Society of London. Series B,  
777 Biological sciences*, 366(1569):1425–37.
- 778 Nelson, P. R., McCune, B., Wheeler, T., Geiser, L. H., and Crisafulli, C. M. (2018).  
779 Lichen community development along a volcanic disturbance gradient at Mount St.  
780 Helens. In *Ecological Responses at Mount St. Helens: Revisited 35 years after the  
781 1980 Eruption*, pages 185–198. Springer New York.
- 782 Newman, M. (2010). *Networks: An Introduction*. Oxford University Press.
- 783 Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings  
784 of the National Academy of Sciences of the United States of America*, 103(23):8577–  
785 82.
- 786 Norby, R. J. and Sigal, L. L. (1989). Nitrogen fixation in the lichen *Lobaria pulmonaria*  
787 in elevated atmospheric carbon dioxide. *Oecologia*, 79(4):566–568.
- 788 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,  
789 P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoces, E., and  
790 Wagner, H. (2019). *vegan: Community Ecology Package*. R package version 2.5-6.
- 791 Poisot, T., Cirtwill, A. R., Cazelles, K., Gravel, D., Fortin, M., and Stouffer, D. B.  
792 (2016). The structure of probabilistic networks. *Methods in Ecology and Evolution*,  
793 7(3):303–312.

- 794 R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R  
795 Foundation for Statistical Computing, Vienna, Austria.
- 796 Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P., and Bascompte, J. (2007).  
797 Non-random coextinctions in phylogenetically structured mutualistic networks. *Nature*,  
798 448(7156):925–8.
- 799 Root, H. T., Miller, J. E. D., and McCune, B. (2011). Biotic soil crust lichen diversity  
800 and conservation in shrub-steppe habitats of Oregon and Washington. *The Bryologist*,  
801 114(4):796.
- 802 Scheipl, F., Greven, S., and Kuechenhoff, H. (2008). Size and power of tests for a zero  
803 random effect variance or polynomial regression in additive and linear mixed models.  
804 *Computational Statistics & Data Analysis*, 52(7):3283–3299.
- 805 Schoch, D. (2020). *signnet: An R package to analyze signed networks*.
- 806 Shuster, S. M., Lonsdorf, E. V., Wimp, G. M., Bailey, J. K., and Whitham, T. G. (2006).  
807 COMMUNITY HERITABILITY MEASURES THE EVOLUTIONARY CONSE-  
808 QUENCES OF INDIRECT GENETIC EFFECTS ON COMMUNITY STRUCTURE.  
809 *Evolution*, 60(5):991.
- 810 Smith, D. S., Bailey, J. K., Shuster, S. M., and Whitham, T. G. (2011). A geographic  
811 mosaic of trophic interactions and selection: trees, aphids and birds. *Journal of*  
812 *evolutionary biology*, 24(2):422–9.
- 813 Smith, D. S., Lamit, L. J., Lau, M. K., Gehring, C. A., Shuster, S. M., and Whitham,  
814 T. G. (2015). Introduced elk alter traits of a native plant and its plant-associated  
815 arthropod community. *Acta Oecologica*, 67:8–16.
- 816 Sole, R. and Bascompte, J. (2006). *Self-Organization in Complex Ecosystems*. Princeton  
817 University Press, Princeton, N.J.
- 818 Staniczenko, P. P., Kopp, J. C., and Allesina, S. (2013). The ghost of nestedness in  
819 ecological networks. *Nature Communications*, 4(1):1–6.
- 820 Thompson, J. N. (2013). *Relentless Evolution*. University of Chicago Press.
- 821 Toju, H., Guimarães, P. R., Olesen, J. M., and Thompson, J. N. (2014). Assembly of  
822 complex plant-fungus networks. *Nature Communications*.
- 823 Toju, H., Yamamichi, M., Guimarães, P. R., Olesen, J. M., Mougi, A., Yoshida, T., and  
824 Thompson, J. N. (2017). Species-rich networks and eco-evolutionary synthesis at the  
825 metacommunity level.
- 826 Toju, H., Yamamoto, S., Tanabe, A. S., Hayakawa, T., and Ishii, H. S. (2016). Network  
827 modules and hubs in plant-root fungal biomes. *Journal of the Royal Society Interface*.
- 828 Valverde, S., Piñero, J., Corominas-Murtra, B., Montoya, J., Joppa, L., and Solé, R.  
829 (2018). The architecture of mutualistic networks as an evolutionary spandrel. *Nature*  
830 *Ecology and Evolution*, 2(1):94–99.
- 831 Wade, M. J. (2007). The co-evolutionary genetics of ecological communities.
- 832 Weber, M. G., Wagner, C. E., Best, R. J., Harmon, L. J., and Matthews, B. (2017).  
833 Evolution in a Community Context: On Integrating Ecological Interactions and  
834 Macroevolution.
- 835 Whitham, T. G., Allan, G. J., Cooper, H. F., and Shuster, S. M. (2020). Intraspecific Ge-  
836 netic Variation and Species Interactions Contribute to Community Evolution. *Annual*  
837 *Review of Ecology, Evolution, and Systematics*, 51(1).
- 838 Whitham, T. G., Bailey, J. K., Schweitzer, J. A., Shuster, S. M., Bangert, R. K., LeRoy,  
839 C. J., Lonsdorf, E. V., Allan, G. J., DiFazio, S. P., Potts, B. M., Fischer, D. G.,


- 840 Gehring, C. A., Lindroth, R. L., Marks, J. C., Hart, S. C., Wimp, G. M., and Wooley,  
841 S. C. (2006). A framework for community and ecosystem genetics: from genes to  
842 ecosystems. *Nature reviews. Genetics*, 7:510–523.
- 843 Whitham, T. G., Gehring, C. A., Lamit, L. J., Wojtowicz, T., Evans, L. M., Keith, A. R.,  
844 and Smith, D. S. (2012). Community specificity: Life and afterlife effects of genes.
- 845 Winfree, R., Gross, B. J., and Kremen, C. (2011). Valuing pollination services to  
846 agriculture. *Ecological Economics*, 71:80–88.
- 847 Zook, A. E., Eklof, A., Jacob, U., and Allesina, S. (2010). Food webs: Ordering species  
848 according to body size yields high degree of intervality. *Journal of theoretical biology*,  
849 271(1):106–113.
- 850 Zytynska, S. E., Fay, M. F., Penney, D., and Preziosi, R. F. (2011). Genetic variation  
851 in a tropical tree species influences the associated epiphytic plant and invertebrate  
852 communities in a complex forest ecosystem. *Philosophical transactions of the Royal  
853 Society of London. Series B, Biological sciences*, 366:1329–1336.
- 854 Zytynska, S. E., Khudr, M. S., Harris, E., and Preziosi, R. F. (2012). No Title. *Oecologia*,  
855 170(2).




**Figure 1.** The communities of bark lichens were observed in a common garden of replicated genotypes of narrowleaf cottonwood trees (*P. angustifolia*) at the Ogden Nature Center (Ogden, UT). (A) Lichens were sampled within a fixed area ( $100 \text{ cm}^2$ ) on individual trees at two heights, 50cm and 95cm from the ground. (B-I) Close-up photos show the other lichen species observed, respectively: *Xanthomendoza montana*, *Candelariella subdeflexa*, *Rinodina freyi*, *Athallia holocarpa*, *Physcia adscendens*, *Physciella melanachra*, *Physcia undulata* and *Myriolecis hagenii*. Photo Credits: L.J. Lamit (B-D) and R. Reese Næsborg (E-I).




**Figure 2.** Lichen interaction networks were constructed by conducting field observations in  $1 \text{ cm}^2$  cells within a  $100 \text{ cm}^2$  grid on each tree using a checkerboard pattern (grey cells). Thus, a set of  $N$  total cell observations were recorded for each tree with the presence or absence of each species recorded for each cell. Applying the probability-based network modeling method adapted from (Araújo et al., 2011), we calculated the conditional probabilities,  $P(S_i | S_j)$ , for all species pairs and removed (i.e., set equal to zero) species pairs whose joint probabilities,  $P(S_i S_j)$ , were not significant using a confidence interval based comparison of their observed co-occurrence frequency,  $S_i S_j$ , to that expected due to chance alone,  $E[P(S_i S_j)] = P(S_i) P(S_j)$ , and  $P(S_i | S_j)$  reduces to  $P(S_i)$ , the observed individual probability of species  $S_i$ .




**Figure 3.** Lichen networks varied in structure among tree genotypes. Network diagrams of the mean lichen interaction matrices averaged for all trees and for illustrative genotypes (996, WC5 and 1008) showing a range of interaction network structure. Directionality (arrowheads) and sign (red = negative, black = positive) of interactions are shown as edges between species, which are scaled by their magnitude. Species names are abbreviated by the first letter of the genus and specific epithet: Ah = *Athallia holocarpa*, Cs = *Candelariella subdeflexa*, Mh = *Myriolecis hagenii*, Rf = *Rinodina freyi*, Pa = *Physcia adscendens*, Pm = *Physciella melanachra*, Pu = *Physcia undulata*, Xg = *Xanthomendoza galericulata*, Xm = *Xanthomendoza montana*. The sign of the interaction is indicative of greater (positive) or lesser (negative) paired occurrences than expected relative to the overall frequency of occurrence of each species. Ecologically, the links in the network are likely the product of multiple types of interactions (e.g. mutualism, parasitism, competition, facilitation) that could vary over both space and time.



**Figure 4.** The similarity of lichen networks varied among tree genotypes. A. The plot shows genotype centroids of NMDS ordinated ( $R^2 = 0.999$ , stress = 0.008) lichen network similarity ( $\pm 1$  S.E.). Genotype centroids that are closer together tend to have more similar lichen network structure. Arrows showing the direction (arrowhead) and magnitude (length) of the vectors of correlation between bark roughness (BR) and network centralization (Cen) and the ordinated networks. B. Plot showing the standardized ( $\frac{x-\bar{x}}{\sigma}$ ) means ( $\pm 1$  S.E.) for the two of the genetically based lichen network metrics: overall degree (i.e., total number of links) and centralization, which is a measure of the dominance of one species in the network.



**Figure 5.** Dot-plots showing the mean (dot) and  $\pm 1$  SE of in-degree (A) and out-degree (B) centrality for two species, *A. holocarpa* and *X. montana*. *Athallia holocarpa* centrality was highly variable among genotypes. *Xanthomendoza montana* centrality, both in- and out-degree, was only non-zero for two genotypes, and only out-degree centrality displayed a significant response to genotype.



**Figure 6.** Bivariate plots of the negative relationship between bark roughness and two network metrics: A) degree and B) centralization. Each plot displays the genotype mean ( $\pm 1$  S.E.) for both variables and a least-squares regression line calculated using the genotype means. Generally, as roughness increased the number of interactions (degree) and dominance of those interactions (centralization) decreased.