

# **Genotypic variation in a foundation tree results in heritable ecological network structure**

**Matthew K. Lau<sup>1,2</sup>, Louis J. Lamit<sup>1,3,4</sup>, Rikke Reese Næsborg<sup>1,5</sup>, Stuart R. Borrett<sup>6</sup>, Matthew A. Bowker<sup>7</sup>, and Thomas G. Whitham<sup>1,8</sup>**

**<sup>1</sup>Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011, USA**

**<sup>2</sup>Harvard Forest, Harvard University, 324 N Main St, Petersham, MA 01366, USA**

**<sup>3</sup>Department of Biology, Syracuse University, 107 College Place, Syracuse, NY 13244, USA**

**<sup>4</sup>Department of Environmental and Forest Biology, State University of New York College of Environmental Science and Forestry, 1 Forestry Drive, Syracuse, NY 13210, USA**

**<sup>5</sup>Santa Barbara Botanic Garden, 1212 Mission Canyon Road, Santa Barbara, CA 93105**

**<sup>6</sup>Department of Biology and Marine Biology, University of North Carolina Wilmington, 601 South College Road, Wilmington, NC 28403, USA**

**<sup>7</sup>Duke Network Analysis Center, Duke University, Durham, NC 27708, USA**

**<sup>8</sup>School of Forestry, 200 E. Pine Knoll Dr., Northern Arizona University, Flagstaff, AZ 86011, USA**

**<sup>9</sup>Center for Adaptable Western Landscapes, Northern Arizona University, Flagstaff, AZ 86011, USA**

## **ABSTRACT**

Elucidating the genetic basis to ecological network structure is fundamental to understanding evolution in complex ecosystems. Although previous work has demonstrated that genetic variation can influence food webs and trophic chains, we are unaware of a study that has quantified the heritability of network structure of a foundation species associated community. To examine this, in a 20+ year old common garden we observed epiphytic lichens associated with narrowleaf cottonwood (*Populus angustifolia*), a riparian ecosystem foundation tree species. We constructed and conducted genetic analyses of signed, weighted, directed lichen interaction networks on individual trees. We found three primary results. First, genotype identity significantly predicted lichen network similarity; i.e., replicates of the same genotype supported more similar lichen networks than different genotypes. Second, broad-sense heritability estimates showed that plant genotype explained network similarity ( $H^2 = 0.41$ ), degree ( $H^2 = 0.32$ ) and centralization ( $H^2 = 0.33$ ). Third, of several tree phenotypic traits examined, bark roughness was both heritable ( $H^2 = 0.32$ ) and significantly predicted lichen network similarity ( $R^2 = 0.26$ ). These results support a mechanistic, genetic pathway from variation in a heritable tree trait to ecological network structure and demonstrate that evolution has the potential to act at the community level to influence not only abundances of organisms but also interactions at the scale of entire networks.

Keywords: networks | heritability | community | genetics | lichen | cottonwood | *Populus* | common garden

## **INTRODUCTION**

Evolution occurs in the context of complex ecological networks. Community genetics studies have shown that genetic variation in foundation species, which have large effects on ecosystems by modulating and stabilizing local conditions (Ellison et al., 2005), plays a significant role in defining distinct communities of interacting organisms: such as endophytes, pathogens, lichens, arthropods, and soil microbes (Busby et al., 2015; Barbour et al., 2009; Lamit et al., 2015a). Multiple studies have now demonstrated that genetic variation influences numerous functional traits (e.g., phytochemical, phenological, morphological)

29 that in combination result in a multivariate functional trait phenotype (Holeski et al., 2012) in which  
30 individual plant genotypes support different communities and ecosystem processes (Bailey et al., 2009;  
31 Whitham et al., 2012). Recently, the importance of genetic variation in structuring ecological systems  
32 was reviewed, and not only were many instances of strong genetic effects found in many ecosystems but  
33 the effect of intraspecific variation was at times greater than inter-specific variation (Des Roches et al.,  
34 2018). There is now evidence to support that selection occurs among groups of species (Wade, 2007)  
35 and that genetic variation and phylogenetic relatedness contribute to variation in community assembly  
36 (Crutsinger, 2016) and species interactions (Whitham et al., 2006; Bailey et al., 2009; Moya-Laraño,  
37 2011). These evolutionary dynamics have the potential to shape the structure of ecological interaction  
38 networks (Rezende et al., 2007; Guimarães et al., 2007; Gómez et al., 2009).

39 Empirical and theoretical work in network ecology and evolutionary biology point to the need for  
40 examinations of the genetic basis of ecological network structure. Analyses of ecological networks have  
41 demonstrated that indirect effects can lead to self-organization, producing sign-changing, amplifying  
42 and/or dampening effects (Fath and Patten, 1998; Newman, 2006; Sole and Bascompte, 2006) and other  
43 studies have demonstrated that indirect effects of interactions among species can lead to network structures  
44 that amplify or dampen the effects of selection, such as the formation of star-like structures in which there  
45 is a “central” species or core group of species (Lieberman et al., 2005). Also, work by Toju et al. (2014,  
46 2016, 2017) observed consistent patterns of centralized interactions of species modules (i.e., groups of  
47 species that interact more strongly within their group than with other species) focused around hubs of  
48 plant-fungal interactions. In other words, a small number of plant and fungal symbionts tended to have  
49 disproportionate numbers of interactions with other species and likely are the drivers in determining  
50 community assembly, structure and dynamics. Interspecific indirect genetic effects (IIGE) theory (*sensu*  
51 Shuster et al. (2006)) in evolutionary biology also points to the importance of studying the genetics of  
52 interaction network structure. Genetically based differences in network structure among individuals  
53 can be acted upon by natural selection when there are fitness consequences of different networks of  
54 IIGEs, leading to community evolution per Whitham et al. (2020) and, by extension, interaction network  
55 evolution. For example, although the analysis was of abundances rather than interaction networks, Gehring  
56 et al. (2014, 2017) found that the mycorrhizal communities on the roots of drought tolerant and intolerant  
57 trees are dominated by different orders of ectomycorrhizal fungal mutualists that also differ in the benefits  
58 they provide that enhance tree performance. Because drought tolerant genotypes are three times more  
59 likely to survive record droughts, selection acts both on the tree and its fungal community and with  
60 increased drought the community phenotype has changed over time. Also, in an antagonistic interaction  
61 context, Busby et al. (2015) found that with the addition of a damaging leaf pathogen to cottonwoods in a  
62 common garden, the impacts of these strong interactors results in a different and diminished community  
63 of arthropods relative to control trees. These examples collectively support the possibility that selection  
64 acting on the tree may alter the network structure of associated communities in which different networks  
65 are more likely to survive drought and pathogen outbreaks, respectively. Regardless of whether the IIGE  
66 is unilateral (i.e., tree affects the community) or reciprocal (i.e., the community also affects the relative  
67 fitness of the tree), selection at the level of the tree population or its community, or both, can change  
68 network structure and alter community dynamics (Whitham et al., 2020).

69 In this context, the “genetic similarity rule” of community genetics provides a useful framework  
70 we can apply to interaction networks at the nexus of ecological and evolutionary dynamics. In a study  
71 combining experimental common gardens and landscape-scale observations of interactions between  
72 *Populus* spp. (cottonwoods) and arthropods, Bangert et al. (2006) observed that individual genotypes  
73 that are more genetically similar will tend to have similar phytochemical traits and thus tend to have  
74 similar interactions with other species. Although this is likely to have consequences for interactions and  
75 network structure, studies in the network ecology literature generally do not include a genetic component  
76 (Lau et al., 2017) and community genetics studies have primarily focused on community composition  
77 in terms of the abundance of species (Des Roches et al., 2018). Some studies have examined the effects  
78 of genetic variation on trophic chains in plant-associated communities (including *Populus*, *Solidago*,  
79 *Oenothera*, *Salix*) (Bailey et al., 2005; Johnson, 2008; Smith et al., 2011, 2015; Barbour et al., 2016) and  
80 generally found that increasing genotypic diversity leads to increased trophic complexity. We are aware  
81 of only two studies that explicitly examined the effect of genotypic variation on interaction networks  
82 between tree individuals and associated herbivores using ecological network metrics (Lau et al., 2016a;  
83 Keith et al., 2017). Both found that genotypic diversity generates increased network modularity (i.e.,

84 compartmentalization); however, both were examining networks at the scale of forest stands, rather than  
85 networks associated with individual trees; therefore, neither was able to observe replicated networks in  
86 order to statistically test for genetic effects on network structure and quantify the genetic component (i.e.,  
87 heritable variation) in network structure.

88 Here, we investigate how genetic variation in a foundation tree species determines the structure of a  
89 network of interactions among a community of tree associated lichens. We used a long-term (20+ years),  
90 common garden experiment with clonally replicated *Populus angustifolia* individuals of known genetic  
91 identity (Martinsen et al., 2001). We focused on a community of epiphytic lichen species, as previous  
92 research has demonstrated significant compositional effects of genotypic variation on lichen in this system  
93 (Lamit et al., 2011, 2015b,a) and epiphytic organisms in other systems (Winfrey et al., 2011; Zytynska  
94 et al., 2011). Applying a probability-theory based network modeling approach (Araújo et al., 2011), we  
95 constructed a set of interaction network models for the lichens associated with individual trees. Using  
96 these models, we then examined the genetic basis of the structure of these ecological networks via several  
97 network metrics that measure different aspects of network structure at the scale of individual species (i.e.,  
98 nodes) or the entire network observed on each tree genotype. Given the potential importance of focal or  
99 “central” nodes (e.g., species) for determining network dynamics (Lieberman et al., 2005), we focused on  
100 network metrics that measure centrality for individual species and centralization for whole networks. Both  
101 of these metrics measure how much a species is connected in the network relative to other species. As  
102 there is a preponderance of evidence that in natural systems evolution occurs in communities comprised of  
103 networks of interacting species (Lau et al., 2016a; Keith et al., 2017; Thompson, 2013; Bascompte et al.,  
104 2006), we set out to test two hypotheses. First, per the community similarity rule (Bangert et al., 2006)  
105 and IIGE theory (Whitham et al., 2020), we hypothesize that trees of the same genotype (i.e., clones) will  
106 support more similar lichen interaction networks relative to individuals of other genotypes. In other words,  
107 epiphytic lichen network structure is heritable, which can be calculated via comparisons of within and  
108 among group variation in network structure. Second, heritability of lichen network structure is the result  
109 of underlying phenotypic covariation in tree traits important to interactions between trees and lichens and  
110 among lichens. Evidence that such trait covariance generates variation in interactions among community  
111 members provides an intermediate genetics-based mechanism for the underlying factors determining  
112 lichen distribution and abundance. In combination, evaluating these two hypotheses is fundamental to  
113 understanding variation and dynamics of network structure and evolution.

## 114 MATERIALS AND METHODS

### 115 Study System

116 The study was conducted along the Weber River, UT (USA), which is a cottonwood (*Populus* spp.)  
117 dominated riparian ecosystem. Although two native species, *Populus angustifolia* (James) and *Populus*  
118 *fremontii* (S. Watson), occur here and are known to hybridize, in order to focus on intra-specific genetic  
119 variation we only sampled pure or advanced generation back-crosses of *P. angustifolia*. Bark lichens  
120 have been intensively sampled in this system and provide an ideal community in which to observe and  
121 model interaction networks, as their sessile nature permits accurate identification of individuals and their  
122 highly localized, direct contact interactions and slow population turnover rates facilitate the assessment of  
123 interactions among lichen species on individual trees (Lamit et al., 2015b).

124 A long-term, common garden experiment was used to isolate the effect of tree genotype from the  
125 effect of the localized microenvironment associated with each individual and spatial autocorrelation.  
126 Established in 1992, asexually propagated clones of genotyped *P. angustifolia* individuals were obtained  
127 from wild collections and planted in a fully randomized design at the Ogden Nature Center, Ogden, UT.  
128 From the population of established individuals in the common garden, we sampled a total of ten genotypes,  
129 replicated between 3 and 8 times each. These individuals comprised a set of tree genotypes with lichen  
130 communities that have been well studied by previous investigations (Lamit et al., 2011, 2015b,a).

### 131 Bark Lichens and Trait Observations

132 We conducted a modified sampling procedure originally developed by Lamit et al. (2015b). On each  
133 tree, presence or absence of each lichen species was assessed in a total of 50 1 cm<sup>2</sup> cells arrayed in  
134 a 10 cm<sup>2</sup> checkerboard pattern. Given the small size and sessile nature of lichens, we were able to  
135 rapidly assess lichen interactions by quantifying thalli of different species occurring in close proximity.  
136 Sampling was restricted to the northern aspect of the trunk to maximize the abundance of lichens and

control for the effect of trunk aspect. Two adjacent  $100\text{ cm}^2$  quadrats centered at 50 cm and 95 cm from ground level were sampled (Fig 1). The observed lichen community included: *Athallia holocarpa*, *Candelariella subdeflexa*, *Myriolecis hagenii*, *Rinodina freyi*, *Physcia adscendens*, *Physciella melanachra*, *Physcia undulata*, *Xanthomendoza galericulata*, *Xanthomendoza montana*. Several other species were not observed in the present study but are known to occur in this region: *Melanohalea elegantula*, *Melanohalea subolivacea*, *Phaeophyscia ciliata* and *Phaeophyscia orbicularis*.

The cell size and checkerboard sampling pattern was chosen to isolate the individuals in each cell. In a survey of *Xanthomendoza galericulata* in the common garden, we had observed a median thallus size of  $0.12 \pm 0.001\text{ cm}^2$  (1 S.E.) (Supporting Information, Fig. 1). This expected thallus size formed the basis for our sampling design, such that lichen observations were spatially independent of thalli present in other cells but exposed to similar micro-environmental conditions created by the bark and the location of the sampling area on an individual tree. Therefore, we were confident in treating the cell-wise observations in quadrats as independent with respect to lichen-lichen interactions.

We quantified tree traits inside or in close proximity to the lichen quadrats. Selected traits have been demonstrated previously to be under strong genetic control in cottonwoods (Bdeir et al., 2017) and other foundation tree species, such as *Eucalyptus* (Nantongo et al., 2020), and previous work has provided evidence for effects on lichen communities of some of these traits (Lamit et al., 2011). We assessed bark texture/structure, herafter referred to as roughness, in the quadrat as the percent of  $1\text{ cm}^2$  cells with “rough” bark, i.e., bark containing a fractured surface. In addition, we also examined several bark chemistry traits by taking bark samples immediately adjacent to each quadrat. We used previously collected phytochemical data from Lamit et al. (2011), including the concentration of condensed tannins, carbon and nitrogen. Additionally, we quantified bark pH for each tree. Bark samples were collected by excavating adjacent to the quadrat down to a depth of 2 mm. Bark pieces were air dried for storage and later processing. Samples were prepped for pH measurements by crushing with a mortar and pestle until all pieces were approximately 0.5 cm in diameter, creating equivalent surface areas among samples. 0.5 g of crushed bark was placed in a 15 ml Falcon collection tube with 5 ml of deionized water. Tubes were capped and let sit for 24 hrs prior to pH measurement with a SevenGo™ pH meter (Mettler Toledo).

#### Lichen Network Modeling

For each tree, the repeated observations of lichens were used to construct replicated interaction networks, i.e. one for each individual tree. Unipartite networks were generated using the conditional probabilities of each species pair, i.e., the probability of observing one species given an observation of another species  $P(S_i|S_j)$ , based on the method developed by Araújo et al. (2011). To calculate conditional probabilities, we quantified the individual probabilities of species occurrences  $P(S_i)$  and the joint probability of co-occurrences  $P(S_i, S_j)$  using the frequencies of each species and their co-occurrences. Using the axioms of probability, we can calculate the conditional probabilities of each species pair as  $P(S_i|S_j) = \frac{P(S_i, S_j)}{P(S_j)}$ . This yields a matrix that could possibly be asymmetrical, i.e.,  $P(S_i|S_j)$  does not have to be equal to  $P(S_j|S_i)$ . Also, the diagonal,  $P(S_i|S_i)$ , is equal to one for all species present and zero for species that were not observed in any cell.

We then applied an analytical procedure to remove non-significant links between species. This procedure determines if the joint probability of a species pair (i.e.,  $P(S_i, S_j)$ ) is different from zero (Fig. 2). Here, a confidence interval  $CI_{95\%}$  is calculated as  $CI_{95\%} = E[S_i S_j] * Z_{95\%} * \sqrt{V(S_i S_j)}$ , where the expected frequency of co-occurrences  $E(S_i S_j)$  is the total number of cells surveyed ( $N$ ) times the independent probabilities of each species  $P(S_i) * P(S_j)$ ,  $Z_{95\%}$  is the Z-score for 95% from a Z-distribution and the expected variance of  $E(S_i S_j)$  is the total number of cells times the expected probability of  $S_i S_j$  and its compliment (i.e.,  $V(S_i S_j) = N * E[P(S_i, S_j)] * (1 - E[P(S_i, S_j)])$ ). If the observed number of co-occurrence falls within the confidence interval, the joint probability  $P(S_i, S_j)$  is concluded to be equal to the product of the individual probabilities (i.e.,  $P(S_i)P(S_j)$ ), and the conditional probability reduces to the individual probability of that species  $P(S_i)$ . Therefore, unless the co-occurrence of a species pair falls outside the confidence interval, the probability that the observation of one species given the other is no different than simply observing that species alone. This enables us to remove links from a given network by re-scaling the resulting conditional probabilities through subtraction of the individual probabilities from the conditional probabilities (i.e., how different the conditional probability is from the independent probability), which makes any species with a non-significant conditional probability zero.

The resulting matrix ( $\mathbf{D} = D_{ij}$ ) can be interpreted as one species’ impact on another with zero being

no effect and values less than or greater than zero being negative and positive effects, respectively. We will refer to  $\mathbf{D}$  as a signed, weighted interaction matrix. As such,  $\mathbf{D}$  has the properties that it can be asymmetric (i.e.,  $D_{ij}$  does not necessarily equal  $D_{ji}$ ) and that it scales between -1 and 1, and, therefore, does not have the mathematical properties of a probabilistic network (Poisot et al., 2016). Also, as the method does not track individuals within species; therefore, the “intra-specific” observations are the same species being counted across the cells of the grid and a resulting  $D_{ii} = 0$ . In the context of these networks, asymmetry and positive/negative valued connections are distinct quantities. In-coming and out-going connections can be interpreted as “influenced by” and “influenced”, respectively; while positive and negative are within this study interpreted as one species increasing or decreasing, respectively, the probability of another species’ occurrence.

## Analyses, Software and Data

To quantify the structural variation of lichen networks we calculated several metrics at both the level of node and whole networks. Although there are many other network metrics, for the sake of simplicity we focus on a subset that represent the primary interesting features of network structure, see Lau et al. (2017). We calculated the number of interactions or “links” in each network (degree), which provides a measure of the size of the network (Lau et al., 2016a; Borrett and Lau, 2014). We also calculated the centralization of each network using Freeman’s centrality, which measures the evenness of the distribution of interactions among the species in the network, using the `sna` package (Butts, 2019). In a network with low centralization species have similar strengths and numbers of interactions. A network with high centralization tends to have one or a small number of species that interact with other species. We used a related function to calculate the centrality of each species (i.e., node level centrality) in each network as well. To calculate separate metrics for positive and negative links, as the networks contained not only positive and negative connections but also directional connections (both in-coming and out-going), we calculated the same network metrics for all combinations of these types of connections using recently developed methods for signed, weighted and directed networks (Everett and Borgatti, 2014) using the `signnet` package (Schoch, 2020).

We used a combination of parametric and non-parametric, permutation based frequentist statistical analyses to test for the effects of genetic variation on lichen communities and their interaction networks. To assess the effect of genotype on traits as univariate response variables (including the metrics of network structure), we used additive, random effects models with Restricted Maximum Likelihood (REML) conducted in R via the `lme4` and `RLRsim` packages (Bates et al., 2015; Scheipl et al., 2008). To conform to test assumptions, traits were root transformed with the exception of condensed tannin concentration and carbon-nitrogen ratio, which were rank and  $\log_{10}$  transformed, respectively. Differences in node level centrality among species was tested using ANOVA and Tukey-HSD multiple comparison tests. Correlations among trait variables and network metrics were quantified and tested using linear correlations of Pearson’s  $r$ . For multivariate response variables, such as lichen community composition and network structure, we used distance based multivariate statistical approaches. To quantify the similarity of lichen networks among individual trees, we calculated the pairwise Euclidean distance of the  $\mathbf{D}$  interaction matrices among all trees (Newman, 2010). For community composition we applied Bray-Curtis similarity to a matrix of species abundances obtained by aggregating the gridded observations by summing over the binary cell-wise species presence-absences. To test for the effects of genotype and other predictor variables on community and network similarity we conducted Permutational Analysis of Variance (PERMANOVA) with `vegan` (Oksanen et al., 2019) using 100000 permutations. For visualization of multivariate patterns, we used Non-metric Multi-Dimensional Scaling (NMDS) (Goslee and Urban, 2007) to produce dimensionally reduced ordinations of these multivariate responses and fitted vectors for continuous predictor variables to the ordinated values (Oksanen et al., 2019), using 100 random initial configurations with a maximum of 1000 iterations and a change in stress threshold of less than  $10^{-12}$ . This was repeated for one to four dimension configurations, and the configuration with the lowest dimensionality and an unexplained variation less than 10% was selected. For all tests where genotype was used as a predictor, we quantified the heritability of the response variable. Because the trees in the garden were clonal replicates of each genotype, we calculated broad-sense heritability, which is the genotypic variance divided by the total phenotypic variance (Conner and Hartl, 2004), which can be interpreted as a measure of the phenotypic variance due to genotypic variation. All analyses were conducted using R version 4.0.2 (R Core Team, 2020). Code and data for the project

245 are openly available as a reproducible workflow using `drake` (Landau, 2018) archived via Zenodo  
246 <https://doi.org/10.5281/zenodo.4581639>.

## 247 RESULTS

248 In support of our first hypotheses, we found that tree genotype influenced lichen network structure and  
249 that multiple lichen network metrics were heritable. Tree genotype significantly predicted the structural  
250 similarity of lichen networks and, overall, network-level metrics responded significantly to tree genotype,  
251 including network degree and centralization including both in-coming and out-going links or when  
252 separated into in-coming only or out-going only (Table 1, Fig. 4). Metrics including only positive  
253 links also showed a significant effect of tree genotype, including positive degree and positive in-going  
254 centralization. Metrics calculated with negative links were not significant, including degree (negative)  
255 and both in-coming (negative) and out-going centralization (negative). Interestingly, although network  
256 similarity and multiple network metrics were significantly predicted by tree genotype, we did not observe  
257 a significant genotypic effect for community composition ( $\text{Pseudo-}F_{9,27} = 0.751$ ,  $R^2 = 0.20$ ,  $p\text{-value} = 0.888$ ).  
258

	Response	df	RLRT	H2	p-value
Lichen Network Similarity	9	3.5821	0.41	0.0537	
Degree	9	3.5175	0.32	0.0255	
Degree (positive)	9	3.6925	0.32	0.0229	
Degree (negative)	9	0.0327	0.03	0.3859	
Centralization	9	4.0444	0.33	0.0184	
Centralization In-Degree	9	4.4812	0.35	0.0142	
Centralization In-Degree (positive)	9	3.9852	0.33	0.0190	
Centralization In-Degree (negative)	9	0.3304	0.11	0.2508	
Centralization Out-Degree	9	3.8615	0.32	0.0205	
Centralization Out-Degree (positive)	9	3.5585	0.31	0.0248	
Centralization Out-Degree (negative)	9	0.0862	0.05	0.3446	

**Table 1.** Genotypic effects on the associated lichen network structure. *RLRT* is the statistic from the restricted likelihood ratio tests.

259 The genetic response of network centralization was driven by variation in *Athallia holocarpa*. Cen-  
260 tralinity varied significantly among species ( $F_{8,324} = 7.99$ ,  $R^2 = 0.16$ ,  $p\text{-value} < 0.0001$ ). The node-level  
261 metrics for *A. holocarpa* displayed the strongest response to tree genotype with high levels of heritability  
262 of positive centrality for both the in-coming ( $RLRT = 3.61$ ,  $H^2 = 0.32$ ,  $p\text{-value} = 0.0240$ ) and out-going  
263 ( $RLRT = 3.13$ ,  $H^2 = 0.30$ ,  $p\text{-value} = 0.0327$ ) perspectives but not for either negative centrality metrics  
264 in-coming ( $RLRT = 0$ ,  $H^2 = 0$ ,  $p\text{-value} = 1$ ) or out-going ( $RLRT = 0$ ,  $H^2 = 0$ ,  $p\text{-value} = 0.4543$ ). None  
265 of the other species' centralities showed a genotypic response (Supporting Information, Fig. 2) with  
266 the exception of *X. montana* ( $RLRT = 2.92$ ,  $H^2 = 0.32$ ,  $p\text{-value} = 0.0375$ ); however, the centrality of *X.*  
267 *montana* was much lower overall relative to *A. holocarpa* and the variation in *X. montana* centrality was  
268 restricted to two genotypes (Fig. 5).

269 In support of our second hypothesis, analysis of trait covariation revealed that genotype indirectly  
270 influenced lichen network centralization via genetically based variation in bark roughness. The percent  
271 cover of rough bark ( $RLRT = 4.8526$ ,  $H^2 = 0.3221$ ,  $p\text{-value} = 0.0113$ ) and condensed tannins ( $RLRT =$   
272  $3.0522$ ,  $H^2 = 0.3205$ ,  $p\text{-value} = 0.0343$ ) both displayed significant responses to tree genotype. None of  
273 the other bark traits, pH ( $RLRT = 0.00$ ,  $H^2 = 0.00$ ,  $p\text{-value} = 1.0000$ ) or carbon-nitrogen ratio ( $RLRT$   
274 =  $0.0000$ ,  $H^2 = 0.0000$ ,  $p\text{-value} = 1.0000$ ), showed a significant response to tree genotype and none  
275 other than bark roughness was correlated with network similarity (Table 2); therefore, we focused our  
276 subsequent analyses on the indirect effect of genotype on lichen network structure via bark roughness. We  
277 found that bark roughness was significantly correlated with network similarity and other lichen network  
278 metrics, including negative correlations with overall network degree ( $df = 35$ ,  $t = -2.13$ ,  $r = -0.34$ ,  $p\text{-value}$   
279 =  $0.04$ ) and centralization ( $df = 35$ ,  $t = -2.52$ ,  $r = -0.39$ ,  $p\text{-value} = 0.02$ ). In other words, trees with more  
280 similar levels of bark roughness tended to have lichen interaction networks with similar structure (Fig. 6).  
281 To quantify the genetic bases of this effect of bark roughness on network structure, we used the residual

values from regressions of the network metrics and bark roughness in subsequent tests of the effect of tree genotype and found no significant effect of tree genotype for either degree ( $RLRT = 0.00, H^2 = 0.00, p\text{-value} = 1.0000$ ) or centralization ( $RLRT = 0.00, H^2 = 0.00, p\text{-value} = 1.0000$ ), and, thus, the bulk of the genetically based variation in the network metrics can be explained by bark roughness.

	df	SS	R2	Pseudo-F	p-value
Bark Roughness	1	20850.09	0.26	12.9234	0.0101
Condensed Tannins	1	5993.66	0.07	3.7150	0.0813
pH	1	1273.19	0.02	0.7892	0.3712
Carbon:Nitrogen Ratio	1	3896.18	0.05	2.4150	0.1890
Residual	32	51627.33	0.64		
Total	36	80993.59	1.00		

**Table 2.** PERMANOVA Pseudo-F Table of lichen network similarity response to bark traits.

## DISCUSSION

### Evolutionary Importance of Ecological Network Heritability

Although previous studies have examined aspects of ecological networks, such as trophic complexity (Barbour et al., 2016) and forest stand level interaction network structure (Lau et al., 2016b; Keith et al., 2017), this is the first study that we are aware of to examine the heritability of network structure with replicated networks at the genotype scale. Previous work in the evolution of ecological networks have primarily focused on macro-evolutionary dynamics (Rezende et al., 2007; Weber et al., 2017; Valverde et al., 2018; Harmon et al., 2019) or have been simulation based individual-level models that integrate intraspecific variation to the species level (Maliet et al., 2020), even though recent syntheses have pointed to the importance of processes operating across scales of organization (Guimarães, 2020).

There are two important functional ramifications of genetically based variation in network structure. First, heritability of network structure suggests that some amount of interaction network complexity is determined and therefore could be predicted by genetic identity. Variation in space and time create variation in ecological networks that influences evolutionary dynamics via shifts in ecological dynamics, such as population demographics (Guimarães, 2020). Given that ecosystems are comprised of such a large number of species, each having a multitude of interactions, the potential to find traction for making predictions in the context of ecological let alone evolutionary dynamics is daunting. The promise of predictability lies in the presence of asymmetries in ecosystems, such as hierarchy created by foundation species via differences in body size and/or life-history strategies (Ellison et al., 2005). Second, heritability (i.e., genetic determination) means that there is structure in the spatial or temporal variation that is created by individuals of foundation species whose traits are in part determined by underlying trait differences. Although this variation is inherently a function of both genetic and environmental effects (Conner and Hartl, 2004), the community and network-level effects are also a function of the scale of the interaction (Shuster et al., 2006; Lau et al., 2017).

Game theoretic applications to ecological networks provide another theoretical basis that relates more directly to the heritability of network centralization found in the present study. Although developed at the population scale, the network theory could be applied at the community scale as well (Lieberman et al., 2005). One seemingly useful direction is the classification of networks into two general categories, rooted and cyclic, in which rooted networks have interactions in which evolutionary effects emanate from one or multiple origins but these effects do not have feedbacks to the origin, whereas cyclic networks contain feedbacks to one or more origins. This is equivalent to “unidirectional” and “reciprocal” genetic effects in the context of IIIGE theory (Whitham et al., 2020). As we do not have an estimate of the effect of the lichen at the larger scale of the fitness of the tree they occur on, we can not determine whether the lichen networks in this system are cyclic or not. In terrestrial ecosystems, lichens play important ecological roles, such as substrate stabilization (Root et al., 2011) and nitrogen fixation (Nelson et al., 2018). Some epiphytic lichens can have demonstrable effects on the availability of nutrients for the trees that are associated with (Norby and Sigal, 1989). Although none of the lichens the present study’s system is known to fix nitrogen, it is possible that they might add micro-nutrients or provide some other unobserved benefit to their host trees. Elucidating the presence of and quantifying such feedbacks would permit the

325 determination of the cyclic nature evolutionary dynamics. If there are positive effects of lichens on host  
326 trees that might increase their ability to respond to environmental stress, then selection could enhance tree  
327 performance and trees with superior communities are more likely to survive. Gehring et al. (2014, 2017)  
328 showed this with ectomycorrhizal communities in which trees with superior mutualist communities were  
329 more likely to survive drought and community evolution occurred (Whitham et al., 2020). However, such  
330 feedbacks to the higher level of the foundation species are not a requirement for evolution, and, regardless  
331 of whether networks are rooted (no feedbacks) or cyclic (feedbacks present), selection at the community  
332 level leading to evolution can still occur. Specifically within the context of the current study, even if  
333 lichens and their interactions do not feed back to affect the performance and ultimate fitness of the tree on  
334 which they reside, non-random death of trees, such as those observed for drought in arid systems Sthultz  
335 et al. (2009); Gehring et al. (2017), can still result in selection at the community level and evolution. For  
336 example, when a tree dies from some event (e.g., a drought, fire, storm, etc.), its lichen network is selected  
337 against while intact networks persist on other trees that survive this selection event.

338 The demonstration of evolution by natural selection requires three key elements (Conner and Hartl,  
339 2004), which multilevel selection theory posits can occur simultaneously at multiple levels of ecological  
340 organization (Whitham et al., 2003, 2020). First, there must be variation, which at the community level  
341 means variation in species abundance, richness, interactions and/or interaction network structure. Second,  
342 these differences must be genetically based and heritable in which community structure is passed from  
343 one generation to the next. For example, numerous studies show that related individuals tend to support  
344 the same communities of insects and microbes, and ecosystem processes of biodiversity, nutrient cycling  
345 and stability, whereas unrelated individuals support more different communities and ecosystem processes  
346 (Bangert et al., 2006, 2008; Barbour et al., 2009; Whitham et al., 2020). Importantly, the current study  
347 shows that lichen networks of individual trees are heritable traits, which provides the important mechanism  
348 for selection to act. Third, selection must favor some communities over others leading to change over  
349 time (i.e., community evolution). The differential survival and performance of individual tree genotypes  
350 will simultaneously result in selection occurring on the lichen community and network structure that  
351 it supports. This is consistent with holobiont theory and empirical studies (Zilber-Rosenberg I, 2008;  
352 Gilbert SF, 2012) in which the holobiome (usually a multicellular host and its symbionts) is the primary  
353 unit of selection (Bordenstein SR, 2015; Johnson and S., 2021). Since our findings show that networks  
354 are heritable, another metric of community evolution would be to quantify how networks change over  
355 time in response to an invasive species, climate change, or some other agent of selection, which represents  
356 a frontier for future ecological network studies.

## 357 **0.1 Community Composition and Quantifying Network Structure**

358 Notably, our findings point strongly to the importance of quantifying interactions and not simply using  
359 community composition as a proxy. We didn't observe compositional differences using the same data from  
360 which the lichen networks were derived. Thus, if we only had analyzed the composition data, we would  
361 have concluded no response of the lichen community to tree genotype, even though the interactions among  
362 lichen species vary among genotypes. As such differences in network structure could occur without  
363 observable differences in species richness or community composition, which have been the primary  
364 focus of the majority of community genetics studies (Whitham et al., 2020). Community composition  
365 of lichens has previously been observed to be different among tree genotypes in the same experimental  
366 garden (Lamit et al., 2011, 2015b). The different results observed in the present study is likely a result of  
367 differences in lichen quantification and the tree genotypes observed leading to overall higher abundances  
368 of observed lichens to assure the possibility of observing lichen interactions. Thus, the findings of the  
369 current study do not negate the previous study, as the quantification methods were different for community  
370 composition, but they do support the conclusion that composition is not a proxy for interactions in complex  
371 communities.

372 It is possible that these underlying differences in interactions among lichens could lead to differences  
373 in community composition at a future point in time via their effects on species abundances (Shuster  
374 et al., 2006); however, this is not needed for evolutionary dynamics to occur via selection that produces  
375 shifts in trait distributions without shifting species abundance distributions, which is possible under  
376 stabilizing, disruptive and directional selection (Conner and Hartl, 2004). Thus, it is imperative that  
377 further community genetics research assess or at least be aware of the potential effects of variation in  
378 interactions and not just observe species abundances, otherwise community level genetic effects may be

379 underestimated, especially when cumulative interaction effects are taken into account (Borrett et al., 2007,  
380 2010).

381 Furthermore, the demonstration of the heritability of interaction networks, without significant dif-  
382 ferences in community composition, provides clear empirical evidence that IIGEs need to encompass  
383 the structure of interaction networks. Although IIGE theory provides a quantitative framework within  
384 which to approach evolutionary theory at higher levels of biological organization (from populations to  
385 communities and ecosystems), this theory has focused on modeling the strong effects of foundation  
386 species (Shuster et al., 2006; Whitham et al., 2012, 2020) and has not yet integrated developments from  
387 the ecological or evolutionary network theory literature. Thus, it has not developed a way to examine  
388 complex interactions among species; however, previous studies have demonstrated this network context  
389 is likely to be important, as altering the structure of interaction networks provides a means for genetic  
390 effects to be damped or magnified within the system of interacting species (Smith et al., 2011; Keith  
391 et al., 2017).

392 Such a synthesis necessitates a much greater effort than can be afforded in this paper; however, in terms  
393 of quantifying interaction networks, we recommend greater focus on foundation species. They are by  
394 definition “central” within the system of interactions and their direct and indirect effects should generally  
395 be greater than other species. Thus, when the more centralized (foundation) species have genetically  
396 based interactions, genetic effects will tend to be propagated and possibly magnified in the community.  
397 Here, we found that even though species with relatively higher abundances (e.g., *X. galericulata*) were  
398 present in the community, the response of the network was not predominately due to their interactions. In  
399 addition, although there were species that tended to be central within the lichen networks, the similarity  
400 of networks showed the highest heritability compared to any of the centrality metrics, which suggests that  
401 there are aspects to network structure that not solely due to the central species.

## 402 **0.2 Network Structure and Scaling**

403 The current study shows the utility of considering the spatial and temporal scales of ecological and  
404 evolutionary dynamics. In line with previous empirical studies (Bangert et al., 2006; Zook et al., 2010;  
405 Zytnyska et al., 2012), we found that the assembly of ecological networks can have a measurable genetic  
406 basis depending on the spatial scale of interactions, due in part, to asymmetries in size and longevity of  
407 organisms. Intra-specific, genotypic diversity among cottonwood trees appears to be a major factor in the  
408 creation of meta-communities of lichens on individual trees that form interaction modules with different  
409 dynamics. When communities are comprised of individuals whose habitat is primarily determined by  
410 another organism, these communities inherently form modules within the larger ecosystem, as they tend  
411 to interact more with each other than with other individuals (Lau et al., 2017).

412 Our study demonstrates that the localized environmental differences determined by the genetic varia-  
413 tion within a single tree species can not only impact community composition, as repeatedly demonstrated  
414 in other community genetics studies (Whitham et al., 2006; Des Roches et al., 2018), but also shape the  
415 structure of interactions among individuals. Some network structures are likely to be more stable, either  
416 in response to disturbance or via self-organized dynamics. For example, centralized networks, although  
417 more efficient, are theorized to be more susceptible to targeted “attacks” in the terminology of defense  
418 networks. As mentioned previously, one class of networks that are theorized to have amplifying effects on  
419 networks have centralized “star” shapes with one or a few species at the center and radiating interactions  
420 out from the central core (Lieberman et al., 2005). This is structurally what we have observed with the  
421 networks that tend to occur on some of the genotypes in our study, i.e., the more centralized networks.  
422 It is likely that these networks could function as hot-spots of evolutionary dynamics resulting from the  
423 amplifying effect of the centralized network structure found on that tree genotype, as multiple studies  
424 have found significant impacts of the removal of foundation species in different systems (Keith et al.,  
425 2017; Des Roches et al., 2018).

426 The importance of the scale of network organization to create hierarchical structure (Guimarães, 2020)  
427 and the potential for foundation species to create this structure in the vast majority of ecosystems (Ellison  
428 et al., 2005; Whitham et al., 2006) suggests that future work would be aided by determining species  
429 with large differences in body-size and longevity. Such differences are likely contributing to interaction  
430 network structure (e.g., modules and centralized species). Many ecological network studies that have an  
431 evolutionary context have focused on asymmetry (e.g., nestedness) and the quantification of its structure  
432 in communities (Bascompte et al., 2006; Díaz-Castelazo et al., 2010; Guimarães et al., 2011; Thompson,

433 2013). Although there is growing evidence that the nestedness of mutualistic networks is not necessarily  
434 the result of selection for systems-level properties but could be either the numerical result of asymptotic  
435 abundance distributions leading to uneven interaction frequencies (Staniczenko et al., 2013) and/or a  
436 by-product of selection and divergence creating network “spandrels” in ecosystems (Valverde et al., 2018),  
437 the findings of these studies have generally supported the hypothesis that nestedness (the degree to which  
438 species tend to interact with similar subsets of the community) tends to promote stability in mutualistic,  
439 primarily bipartite (i.e., two-mode), networks and that modularity contributes to the stabilization of  
440 antagonistic networks (Elias et al., 2013; Grilli et al., 2016). We were not able to examine nestedness  
441 or modularity of the lichen networks in the current study, as metrics for analyzing weighted, directed  
442 and signed networks could not be found. Future network theoretic developments will hopefully develop  
443 appropriate metrics to conduct these analyses.

## 444 Conclusion

445 We found support for both of our hypotheses. First, tree genotype influenced the network structure of  
446 lichen communities associated with narrowleaf cottonwoods in a riparian forest ecosystem. Network  
447 similarity and metrics of network structure tended to be more similar on trees of the same genotype.  
448 Generally, this genetic effect was manifested in positive interactions and largely driven by *A. holocarpa*.  
449 Second, the genetically based trait, bark roughness, was observed to affect network variation, largely  
450 via shifts in positive in-coming and out-going interactions. Although bark roughness has previously  
451 been shown to be an important tree trait influencing bark lichens (Lamit et al., 2015b), this is the first  
452 demonstration of a link from plant genetics to lichen network structure. As such, these results have  
453 important implications for the influence of genetically based variation in ecosystems with networks of  
454 interacting species. As heritable variation is the raw material for natural selection to act upon, a genetic  
455 basis for interaction network structure indicates that conserving genetic variation is important to consider  
456 in efforts to restore or preserve complex species interactions and their associated ecosystem functions  
457 (Whitham et al., 2012; Evans et al., 2013; Whitham et al., 2020). Network modeling and analysis will  
458 provide useful tools for the identification of species within network modules that are most important to  
459 study in systems where little is known about the natural history of organisms in an ecosystem. Collectively,  
460 such investigations will bring us closer to understanding the evolutionary drivers of Darwin’s entangled  
461 bank and the interconnectedness of species in complex communities (Darwin, 1859; Dátilo et al., 2016).

## 462 Acknowledgments

463 This work was supported by National Science Foundation grants DEB-0425908 and DEB-1340852, the  
464 NSF funded Northern Arizona University Genes to Ecosystems Integrative Graduate Research Traineeship  
465 (IGERT) and the ARCS Foundation (Phenoix Chapter). Thanks also to the Ogden Nature Center staff  
466 who helped to maintain the common garden experiments and Drs. Todd Wojtowicz, Luke Evans and  
467 David Solance Smith for assistance with lichen network observations.

## 468 REFERENCES

- 469 Araújo, M. B., Rozenfeld, A., Rahbek, C., and Marquet, P. A. (2011). Using species co-occurrence  
470 networks to assess the impacts of climate change. *Ecography*, 34:897–908.
- 471 Bailey, J. K., Schweitzer, J. A., Ubeda, F., Koricheva, J., LeRoy, C. J., Madritch, M. D., Rehill, B. J.,  
472 Bangert, R. K., Fischer, D. G., Allan, G. J., and Whitham, T. G. (2009). From genes to ecosystems: a  
473 synthesis of the effects of plant genetic factors across levels of organization. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 364(1523):1607–16.
- 474 Bailey, J. K., Wooley, S. C., Lindroth, R. L., and Whitham, T. G. (2005). Importance of species  
475 interactions to community heritability: a genetic basis to trophic-level interactions. *Ecology Letters*,  
476 0(0):051122062725008.
- 477 Bangert, R. K., Lonsdorf, E. V., Wimp, G. M., Shuster, S. M., Fischer, D., Schweitzer, J. A., Allan, G. J.,  
478 Bailey, J. K., and Whitham, T. G. (2008). Genetic structure of a foundation species: Scaling community  
479 phenotypes from the individual to the region.
- 480 Bangert, R. K., Turek, R. J., Rehill, B., Wimp, G. M., Schweitzer, J. A., Allan, G. J., Bailey, J. K.,  
481 Martinsen, G. D., Keim, P., Lindroth, R. L., and Whitham, T. G. (2006). A genetic similarity rule  
482 determines arthropod community structure. *Molecular ecology*, 15:1379–1391.

- 484 Barbour, M. A., Fortuna, M. A., Bascompte, J., Nicholson, J. R., Julkunen-Tiitto, R., Jules, E. S., and  
485 Crutsinger, G. M. (2016). Genetic specificity of a plant-insect food web: Implications for linking  
486 genetic variation to network complexity. *Proceedings of the National Academy of Sciences of the*  
487 *United States of America*, 113(8):2128–2133.
- 488 Barbour, R. C., O'Reilly-Wapstra, J. M., Little, D. W. D., Jordan, G. J., Steane, D. A., Humphreys, J. R.,  
489 Bailey, J. K., Whitham, T. G., Potts, B. M., De Little, D. W., Jordan, G. J., Steane, D. A., Humphreys,  
490 J. R., Bailey, J. K., Whitham, T. G., and Potts, B. M. (2009). A geographic mosaic of genetic variation  
491 within a foundation tree species and its community-level consequences. *Ecology*, 90(7):1762–1772.
- 492 Bascompte, J., Jordano, P., and Olesen, J. M. (2006). Asymmetric Coevolutionary Networks Facilitate  
493 Biodiversity Maintenance\r10.1126/science.1123412. *Science*, 312:431–433.
- 494 Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4.  
495 *Journal of Statistical Software*, 67(1):1–48.
- 496 Bdeir, R., Muchero, W., Yordanov, Y., Tuskan, G. A., Busov, V., and Gailing, O. (2017). Quantitative trait  
497 locus mapping of Populus bark features and stem diameter. *BMC Plant Biology*.
- 498 Bordenstein SR, T. K. (2015). Host biology in light of the microbiome: ten principles of holobionts and  
499 hologenomes. *PLOS Biol.*, 13:e1002226.
- 500 Borrett, S. R., Fath, B. D., and Patten, B. C. (2007). Functional integration of ecological networks through  
501 pathway proliferation. *Journal of Theoretical Biology*, 245(1):98–111.
- 502 Borrett, S. R. and Lau, M. K. (2014). enaR: An R package for Ecosystem Network Analysis. *Methods in*  
503 *Ecology and Evolution*, 5(11):1206–1213.
- 504 Borrett, S. R., Whipple, S. J., and Patten, B. C. (2010). Rapid development of indirect effects in ecological  
505 networks. *Oikos*, 119(7):1136–1148.
- 506 Busby, P. E., Lamit, L. J., Keith, A. R., Newcombe, G., Gehring, C. A., Whitham, T. G., and Dirzo,  
507 R. (2015). Genetics-based interactions among plants, pathogens, and herbivores define arthropod  
508 community structure. *Ecology*, 96(7):1974–1984.
- 509 Butts, C. T. (2019). *sna: Tools for Social Network Analysis*. R package version 2.5.
- 510 Conner, K. and Hartl, D. L. (2004). *A Primer of Ecological Genetics: a textbook*. Sinauer Associates,  
511 Inc., Sunderland, Massachusetts, 1st edition.
- 512 Crutsinger, G. M. (2016). A community genetics perspective: Opportunities for the coming decade. *New*  
513 *Phylogist*.
- 514 Darwin, C. (1859). *On the Origin of Species*. Murray, London.
- 515 Dátillo, W., Lara-Rodríguez, N., Jordano, P., Guimarães, P. R., Thompson, J. N., Marquis, R. J., Medeiros,  
516 L. P., Ortiz-Pulido, R., Marcos-García, M. A., and Rico-Gray, V. (2016). Unravelling Darwin's  
517 entangled bank: Architecture and robustness of mutualistic networks with multiple interaction types.  
518 *Proceedings of the Royal Society B: Biological Sciences*.
- 519 Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., Schweitzer, J. A.,  
520 and Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology and*  
521 *Evolution*, 2(1):57–64.
- 522 Díaz-Castelazo, C., Guimarães, P. R., Jordano, P., Thompson, J. N., Marquis, R. J., and Rico-Gray, V.  
523 (2010). Changes of a mutualistic network over time: Reanalysis over a 10-year period. *Ecology*.
- 524 Elias, M., Fontaine, C., and Frank Van Veen, F. J. (2013). Evolutionary history and ecological processes  
525 shape a local multilevel antagonistic network. *Current Biology*, 23(14):1355–1359.
- 526 Ellison, A. M., Bank, M. S., Clinton, B. D., Colburn, E. A., Elliott, K., Ford, C. R., Foster, D. R., Kloeppele,  
527 B. D., Knoepp, J. D., Lovett, G. M., Mohan, J., Orwig, D. A., Rodenhouse, N. L., Sobczak, W. V.,  
528 Stinson, K. A., Stone, J. K., Swan, C. M., Thompson, J., Von Holle, B., and Webster, J. R. (2005). Loss  
529 of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers*  
530 *in Ecology and the Environment*, 3(9):479–486.
- 531 Evans, D. M., Pocock, M. J. O., and Memmott, J. (2013). The robustness of a network of ecological  
532 networks to habitat loss. *Ecology letters*, 16:844–52.
- 533 Everett, M. G. and Borgatti, S. P. (2014). Networks containing negative ties. *Social Networks*, 38(1):111–  
534 120.
- 535 Fath, B. D. and Patten, B. C. (1998). Network synergism: Emergence of positive relations in ecological  
536 systems. *Ecological Modelling*, 107(2):127 – 143.
- 537 Gehring, C., Flores-Rentería, D., Sthultz, C. M., Leonard, T. M., Flores-Rentería, L., Whipple, A. V.,  
538 and Whitham, T. G. (2014). Plant genetics and interspecific competitive interactions determine


- 539 ectomycorrhizal fungal community responses to climate change. *Molecular Ecology*, 23(6):1379–1391.
- 540 Gehring, C. A., Sthultz, C. M., Flores-Rentería, L., Whipple, A. V., and Whitham, T. G. (2017). Tree  
541 genetics defines fungal partner communities that may confer drought tolerance. *Proceedings of the  
542 National Academy of Sciences*.
- 543 Gilbert SF, Sapp J, T. A. (2012). A symbiotic view of life: We have never been individuals. *Q. Rev. Biol.*,  
544 87:325–41.
- 545 Gómez, J. M., Abdelaziz, M., Camacho, J. P. M., Muñoz-Pajares, A. J., and Perfectti, F. (2009). Local  
546 adaptation and maladaptation to pollinators in a generalist geographic mosaic. *Ecology Letters*,  
547 12(7):672–682.
- 548 Goslee, S. C. and Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of ecological  
549 data. *Journal of Statistical Software*, 22:1–19.
- 550 Grilli, J., Rogers, T., and Allesina, S. (2016). Modularity and stability in ecological communities. *Nature  
551 Communications*, 7.
- 552 Guimarães, P. R. (2020). The Structure of Ecological Networks Across Levels of Organization. *Annual  
553 Review of Ecology, Evolution, and Systematics*, 51(1).
- 554 Guimarães, P. R., Jordano, P., and Thompson, J. N. (2011). Evolution and coevolution in mutualistic  
555 networks. *Ecology letters*, 14(9):877–85.
- 556 Guimarães, P. R., Rico-Gray, V., Oliveira, P. S., Izzo, T. J., dos Reis, S. F., and Thompson, J. N. (2007).  
557 Interaction Intimacy Affects Structure and Coevolutionary Dynamics in Mutualistic Networks. *Current  
558 Biology*, 17(20):1797–1803.
- 559 Harmon, L. J., Andreazza, C. S., Débarre, F., Drury, J., Goldberg, E. E., Martins, A. B., Melián, C. J.,  
560 Narwani, A., Nuismer, S. L., Pennell, M. W., Rudman, S. M., Seehausen, O., Silvestro, D., Weber, M.,  
561 and Matthews, B. (2019). Detecting the macroevolutionary signal of species interactions. *Journal of  
562 Evolutionary Biology*, 32(8):769–782.
- 563 Holeski, L. M., Hillstrom, M. L., Whitham, T. G., and Lindroth, R. L. (2012). Relative importance of  
564 genetic, ontogenetic, induction, and seasonal variation in producing a multivariate defense phenotype  
565 in a foundation tree species. *Oecologia*, 170:695–707.
- 566 Johnson, M. T. J. (2008). Bottom-up effects of plant genotype on aphids, ants, and predators. *Ecology*.
- 567 Johnson, N. C. and S., G. K. (2021). Understanding multilevel selection may facilitate management of  
568 arbuscular mycorrhizae in sustainable agroecosystems. *Front. Plant. Sci.*, 11:627345.
- 569 Keith, A. R., Bailey, J. K., Lau, M. K., and Whitham, T. G. (2017). Genetics-based interactions of  
570 foundation species affect community diversity, stability and network structure. *Proceedings of the  
571 Royal Society B: Biological Sciences*, 284(1854):20162703.
- 572 Lamit, L., Bowker, M., Holeski, L., Næsborg, R. R., Wooley, S., Zinkgraf, M., Lindroth, R., Whitham, T.,  
573 and Gehring, C. (2011). Genetically-based trait variation within a foundation tree species influences a  
574 dominant bark lichen. *Fungal Ecology*, 4(1):103–109.
- 575 Lamit, L. J., Busby, P. E., Lau, M. K., Compson, Z. G., Wojtowicz, T., Keith, A. R., Zinkgraf, M. S.,  
576 Schweitzer, J. A., Shuster, S. M., Gehring, C. A., and Whitham, T. G. (2015a). Tree genotype  
577 mediates covariance among communities from microbes to lichens and arthropods. *Journal of Ecology*,  
578 103(4):840–850.
- 579 Lamit, L. J., Lau, M. K., Næsborg, R. R., Wojtowicz, T., Whitham, T. G., and Gehring, C. A. (2015b).  
580 Genotype variation in bark texture drives lichen community assembly across multiple environments.  
581 *Ecology*, 96(4):960–971.
- 582 Landau, W. M. (2018). The drake r package: a pipeline toolkit for reproducibility and high-performance  
583 computing. *Journal of Open Source Software*, 3(21).
- 584 Lau, M. K., Borrett, S. R., Baiser, B., Gotelli, N. J., and Ellison, A. M. (2017). Ecological network  
585 metrics: opportunities for synthesis. *Ecosphere*, 8(8):e01900.
- 586 Lau, M. K., Keith, A. R., Borrett, S. R., Shuster, S. M., and Whitham, T. G. (2016a). Genotypic variation  
587 in foundation species generates network structure that may drive community dynamics and evolution.  
588 *Ecology*, 97(3):15–0600.
- 589 Lau, M. K., Keith, A. R., Borrett, S. R., Shuster, S. M., and Whitham, T. G. (2016b). Genotypic variation  
590 in foundation species generates network structure that may drive community dynamics and evolution.  
591 *Ecology*.
- 592 Lieberman, E., Hauert, C., and Nowak, M. A. (2005). Evolutionary dynamics on graphs. *Nature*,  
593 433(7023):312–316.

- 594 Maliet, O., Loeuille, N., and Morlon, H. (2020). An individual-based model for the eco-evolutionary  
 595 emergence of bipartite interaction networks. *Ecology Letters*, 23(11):1623–1634.
- 596 Martinsen, G. D., Whitham, T. G., Turek, R. J., and Keim, P. (2001). Hybrid populations selectively filter  
 597 gene introgression between species. *Evolution*, 55(7):1325–1335.
- 598 Moya-Laraño, J. (2011). Genetic variation, predator-prey interactions and food web structure. *Philosophical  
 599 transactions of the Royal Society of London. Series B, Biological sciences*, 366(1569):1425–37.
- 600 Nantongo, J. S., Potts, B. M., Fitzgerald, H., Newman, J., Elms, S., Aurik, D., Dungey, H., and O'Reilly-  
 601 Wapstra, J. M. (2020). Quantitative genetic variation in bark stripping of *pinus radiata*. *Forests*,  
 602 11(12):1356.
- 603 Nelson, P. R., McCune, B., Wheeler, T., Geiser, L. H., and Crisafulli, C. M. (2018). Lichen community  
 604 development along a volcanic disturbance gradient at Mount St. Helens. In *Ecological Responses at  
 605 Mount St. Helens: Revisited 35 years after the 1980 Eruption*, pages 185–198. Springer New York.
- 606 Newman, M. (2010). *Networks: An Introduction*. Oxford University Press.
- 607 Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of the National  
 608 Academy of Sciences of the United States of America*, 103(23):8577–82.
- 609 Norby, R. J. and Sigal, L. L. (1989). Nitrogen fixation in the lichen *Lobaria pulmonaria* in elevated  
 610 atmospheric carbon dioxide. *Oecologia*, 79(4):566–568.
- 611 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara,  
 612 R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H. (2019). *vegan:  
 613 Community Ecology Package*. R package version 2.5-6.
- 614 Poisot, T., Cirtwill, A. R., Cazelles, K., Gravel, D., Fortin, M., and Stouffer, D. B. (2016). The structure  
 615 of probabilistic networks. *Methods in Ecology and Evolution*, 7(3):303–312.
- 616 R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for  
 617 Statistical Computing, Vienna, Austria.
- 618 Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P., and Bascompte, J. (2007). Non-random  
 619 coextinctions in phylogenetically structured mutualistic networks. *Nature*, 448(7156):925–8.
- 620 Root, H. T., Miller, J. E. D., and McCune, B. (2011). Biotic soil crust lichen diversity and conservation in  
 621 shrub-steppe habitats of Oregon and Washington. *The Bryologist*, 114(4):796.
- 622 Scheipl, F., Greven, S., and Kuechenhoff, H. (2008). Size and power of tests for a zero random effect  
 623 variance or polynomial regression in additive and linear mixed models. *Computational Statistics &  
 624 Data Analysis*, 52(7):3283–3299.
- 625 Schoch, D. (2020). *signnet: An R package to analyze signed networks*.
- 626 Shuster, S. M., Lonsdorf, E. V., Wimp, G. M., Bailey, J. K., and Whitham, T. G. (2006). COMMUNITY  
 627 HERITABILITY MEASURES THE EVOLUTIONARY CONSEQUENCES OF INDIRECT GENETIC  
 628 EFFECTS ON COMMUNITY STRUCTURE. *Evolution*, 60(5):991.
- 629 Smith, D. S., Bailey, J. K., Shuster, S. M., and Whitham, T. G. (2011). A geographic mosaic of trophic  
 630 interactions and selection: trees, aphids and birds. *Journal of evolutionary biology*, 24(2):422–9.
- 631 Smith, D. S., Lamit, L. J., Lau, M. K., Gehring, C. A., Shuster, S. M., and Whitham, T. G. (2015).  
 632 Introduced elk alter traits of a native plant and its plant-associated arthropod community. *Acta  
 633 Oecologica*, 67:8–16.
- 634 Sole, R. and Bascompte, J. (2006). *Self-Organization in Complex Ecosystems*. Princeton University Press,  
 635 Princeton, N.J.
- 636 Staniczenko, P. P., Kopp, J. C., and Allesina, S. (2013). The ghost of nestedness in ecological networks.  
 637 *Nature Communications*, 4(1):1–6.
- 638 Sthultz, C., Gehring, C., and Whitham, T. (2009). Deadly combination of genes and drought: Increased  
 639 mortality of herbivore-resistant trees in a foundation species. *Global Change Biology*, 15(8):1949–1961.  
 640 Copyright: Copyright 2009 Elsevier B.V., All rights reserved.
- 641 Thompson, J. N. (2013). *Relentless Evolution*. University of Chicago Press.
- 642 Toju, H., Guimarães, P. R., Olesen, J. M., and Thompson, J. N. (2014). Assembly of complex plant-fungus  
 643 networks. *Nature Communications*.
- 644 Toju, H., Yamamichi, M., Guimarães, P. R., Olesen, J. M., Mougi, A., Yoshida, T., and Thompson, J. N.  
 645 (2017). Species-rich networks and eco-evolutionary synthesis at the metacommunity level.
- 646 Toju, H., Yamamoto, S., Tanabe, A. S., Hayakawa, T., and Ishii, H. S. (2016). Network modules and hubs  
 647 in plant-root fungal biomes. *Journal of the Royal Society Interface*.
- 648 Valverde, S., Piñero, J., Corominas-Murtra, B., Montoya, J., Joppa, L., and Solé, R. (2018). The


- 649 architecture of mutualistic networks as an evolutionary spandrel. *Nature Ecology and Evolution*,  
650 2(1):94–99.
- 651 Wade, M. J. (2007). The co-evolutionary genetics of ecological communities.
- 652 Weber, M. G., Wagner, C. E., Best, R. J., Harmon, L. J., and Matthews, B. (2017). Evolution in a  
653 Community Context: On Integrating Ecological Interactions and Macroevolution.
- 654 Whitham, T., Young, W., Martinsen, G., Gehring, C., Schweitzer, J., Shuster, S., Wimp, G., Fischer, D.,  
655 Bailey, J., Lindroth, R., Woolbright, S., and Kuske, C. (2003). Community and ecosystem genetics: A  
656 consequence of the extended phenotype. *Ecology*, 84(3):559–573. Copyright: Copyright 2017 Elsevier  
657 B.V., All rights reserved.
- 658 Whitham, T. G., Allan, G. J., Cooper, H. F., and Shuster, S. M. (2020). Intraspecific Genetic Variation  
659 and Species Interactions Contribute to Community Evolution. *Annual Review of Ecology, Evolution,  
660 and Systematics*, 51(1).
- 661 Whitham, T. G., Bailey, J. K., Schweitzer, J. A., Shuster, S. M., Bangert, R. K., LeRoy, C. J., Lonsdorf,  
662 E. V., Allan, G. J., DiFazio, S. P., Potts, B. M., Fischer, D. G., Gehring, C. A., Lindroth, R. L., Marks,  
663 J. C., Hart, S. C., Wimp, G. M., and Wooley, S. C. (2006). A framework for community and ecosystem  
664 genetics: from genes to ecosystems. *Nature reviews. Genetics*, 7:510–523.
- 665 Whitham, T. G., Gehring, C. A., Lamit, L. J., Wojtowicz, T., Evans, L. M., Keith, A. R., and Smith, D. S.  
666 (2012). Community specificity: Life and afterlife effects of genes.
- 667 Winfree, R., Gross, B. J., and Kremen, C. (2011). Valuing pollination services to agriculture. *Ecological  
668 Economics*, 71:80–88.
- 669 Zilber-Rosenberg I, R. E. (2008). Role of microorganisms in the evolution of animals and plants: the  
670 hologenome theory of evolution. *FEMS Microbiol. Rev.*, 32:723–735.
- 671 Zook, A. E., Eklof, A., Jacob, U., and Allesina, S. (2010). Food webs: Ordering species according to  
672 body size yields high degree of intervality. *Journal of theoretical biology*, 271(1):106–113.
- 673 Zytynska, S. E., Fay, M. F., Penney, D., and Preziosi, R. F. (2011). Genetic variation in a tropical tree  
674 species influences the associated epiphytic plant and invertebrate communities in a complex forest  
675 ecosystem. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*,  
366:1329–1336.
- 676 Zytynska, S. E., Khudr, M. S., Harris, E., and Preziosi, R. F. (2012). No Title. *Oecologia*, 170(2).




**Figure 1.** The communities of bark lichens were observed in a common garden of replicated genotypes of narrowleaf cottonwood trees (*P. angustifolia*) at the Ogden Nature Center (Ogden, UT). (A) Lichens were sampled within a fixed area ( $100 \text{ cm}^2$ ) on individual trees at two heights, 50cm and 95cm from the ground. (B-I) Photos showing lichen species observed, respectively: *Xanthomendoza montana*, *Candelariella subdeflexa*, *Rinodina freyi*, *Athallia holocarpa*, *Physcia adscendens*, *Physciella melanachra*, *Physcia undulata* and *Myriolecis hagenii*. Photo Credits: L.J. Lamit (B-D) and R. Reese Næsborg (E-I).




**Figure 2.** Lichen interaction networks were constructed by conducting field observations in 1 cm<sup>2</sup> cells within a 100 cm<sup>2</sup> grid on each tree using a checkerboard pattern (grey cells). Thus, a set of  $N$  total cell observations were recorded for each tree with the presence or absence of each species recorded for each cell. Applying the probability-based network modeling method adapted from (Araújo et al., 2011), we calculated the conditional probabilities,  $P(S_i | S_j)$ , for all species pairs and removed (i.e., set equal to zero) species pairs whose joint probabilities,  $P(S_i S_j)$ , were not significant using a confidence interval based comparison of their observed co-occurrence frequency,  $S_i S_j$ , to that expected due to chance alone,  $E[P(S_i S_j)] = P(S_i)P(S_j)$ , and  $P(S_i | S_j)$  reduces to  $P(S_i)$ , the observed individual probability of species  $S_i$ .




**Figure 3.** Lichen networks varied in structure among tree genotypes. Network diagrams of the mean lichen interaction matrices averaged for all trees and for illustrative genotypes (996, WC5 and 1008) showing a range of interaction network structure. Directionality (arrowheads) and sign (red = negative, black = positive) of interactions are shown as edges between species, which are scaled by their magnitude. Species names are abbreviated by the first letter of the genus and specific epithet: Ah = *Athallia holocarpa*, Cs = *Candelariella subdeflexa*, Mh = *Myriolecis hagenii*, Rf = *Rinodina freyi*, Pa = *Physcia adscendens*, Pm = *Physciella melanchra*, Pu = *Physcia undulata*, Xg = *Xanthomendoza galericulata*, Xm = *Xanthomendoza montana*. The sign of the interaction is indicative of greater (positive) or lesser (negative) paired occurrences than expected relative to the overall frequency of occurrence of each species. Ecologically, the links in the network are likely the product of multiple types of interactions (e.g. mutualism, parasitism, competition, facilitation) that could vary over both space and time.



**Figure 4.** The similarity of lichen networks varied among tree genotypes. A. The plot shows genotype centroids of NMDS ordinated ( $R^2 = 0.999$ , stress = 0.008) lichen network similarity ( $\pm 1$  S.E.). Genotype centroids that are closer together tend to have more similar lichen network structure. Arrows showing the direction (arrowhead) and magnitude (length) of the vectors of correlation between bark roughness (BR) and network centralization (Cen) and the ordinated networks. B. Plot showing the standardized ( $\frac{x-\bar{x}}{\sigma}$ ) means ( $\pm 1$  S.E.) for the two of the genetically based lichen network metrics: overall degree (i.e., total number of links) and centralization, which is a measure of the dominance of one species in the network.



**Figure 5.** Dot-plots showing the mean (dot) and  $\pm 1$  SE of in-degree (A) and out-degree (B) centrality for two species, *A. holocarpa* and *X. montana*. *Athallia holocarpa* centrality was highly variable among genotypes. *Xanthomendoza montana* centrality, both in- and out-degree, was only non-zero for two genotypes, and only out-degree centrality displayed a significant response to genotype.



**Figure 6.** Bivariate plots of the negative relationship between bark roughness and two network metrics: A) degree and B) centralization. Each plot displays the genotype mean ( $\pm 1$  S.E) for both variables and a least-squares regression line calculated using the genotype means. Generally, as roughness increased the number of interactions (degree) and dominance of those interactions (centralization) decreased.