

Genotypic variation in foundation trees generates ecological network structure

Matthew K. Lau^{1,2,*}, Louis J. Lamit², Rikke R. Naesbourg³, Stuart R. Borrett⁴, Matthew A. Bowker⁵, and Thomas G. Whitham¹

¹Department of Biological Sciences and Merriam-Powell Center for Environmental Research, Northern Arizona University, Flagstaff, AZ 86011, USA

²Harvard Forest, Harvard University, 324 N Main St, Petersham, MA 01366, USA

³University of California Berkeley, Berkeley, CA, USA

⁴Department of Biology and Marine Biology, University of North Carolina Wilmington, 601 South College Road, Wilmington, NC, 28403, USA

⁵School of Forestry, Northern Arizona University, Flagstaff, AZ 86011, USA

*matthewklau@fas.harvard.edu

ABSTRACT

Biological evolution is argued to occur in the context of complex networks of interacting species in which natural selection defines the structure of ecological networks^{1,2,3}. However, fundamental to this evolutionary process is the discovery of a genetic basis to ecological network structure, which remains largely unknown. Here, we use both a long-term experimental common garden⁴ with genotyped individuals and a natural riparian forest of the foundation tree species⁵ *Populus angustifolia*, to test how genetic variation contributes to the interaction network structure of a model community comprised of epiphytic lichens. We found three main results: 1) lichen communities showed significant unipartite (i.e., one mode) network structure that was similar between the common garden and a natural stand, 2) individual tree genotype significantly influenced lichen species interactions, which was strongly correlated with bark roughness, a genetically based trait in cottonwoods⁶ known to influence epiphytic lichen⁷, and 3) bipartite (two mode) genotype-species networks, comprised of the foundation species and its associated lichen community, showed significant modular structure in both the common garden and natural stand. These results demonstrate strong support for a genetic basis to ecological network structure and the potential for selection to act in complex ecosystems. This work sets the stage for studies that address greater complexity in the evolution of biological systems and provides a framework for the discovery of evolutionarily dynamic compartments in ecosystems.

1 Introduction

- 2 Evolution occurs in the context of complex networks of interacting species. In ecological communities, community dynamics
- 3 depend on key interactions⁶ that occur in species interaction networks, such as: trophic⁷ and mutualistic⁸ interaction networks.

4 Phylogenetic patterns in ecological networks support the importance of evolutionary processes in shaping species interactions^{1,9}.
5 Community genetics studies¹⁰ have shown that genetic variation in foundation species⁴ (species, such as trees, that largely
6 define the composition of communities by creating locally stable conditions and modulating resources; contributes to variation
7 in interactions with dependent communities in both terrestrial and aquatic ecosystems¹¹. more specifically, genetic variation
8 affects diverse chemical, phenological, morphological and other traits producing a multivariate phenotype¹² that makes each
9 individual unique so that different communities assemble on different genotypes resulting in different species interaction^{13,14}.
10 although, some empirical studies have shown that genetic variation in a single species can impact a tri-trophic interaction¹⁵;
11 currently no studies have yet investigated the impact of intraspecific genotypic variation on network structure, which is integral
12 to the process of evolution by natural selection.

13 **add newer lit studies here.**

- 14 • McKnight's paper: ecological networks are connected across aquatic and terrestrial systems and impact evolutionary
15 processes
- 16 • Arts 2017 paper: Ecological networks are impacted by the interaction of foundation species
- 17 • Daves 2016 paper
- 18 • Jamie's 2016 paper
- 19 • Nat Eco Evo 2017:
- 20 • Liza and Amy 2017:
- 21 • Ghering 2017: Tree genetics defines fungal partner communities that may confer drought tolerance
- 22 • Bothwell 2017: Precipitation gradients and river networks drive genetic connectivity and diversity in a foundation riparian
23 tree
- 24 • Poisot
- 25 • Josh Tylinakis

26 Andreazzi, C.S, J. N. Thompson, and P. R. Guimarães, Jr. 2017. Network structure and selection asymmetry drive
27 coevolution in species-rich antagonistic interactions. *American Naturalist* 190:99-115

28 Toju, H. M. Yamamichi, P. R. Guimarães, Jr., J. M. Olesen, A. Mougi, T. Yoshida, and J. N. Thompson. 2017. Species-
29 rich networks and eco-evolutionary synthesis at the metacommunity level. *Nature Ecology and Evolution* 1:0024. DOI:
30 10.1038/s41559-016-0024

31 Dáttilo, W., N. Lara-Rodriguez, P. Jordano, P. R. Guimarães, Jr., J. N. Thompson, R. J. Marquis, L. P. Medeiros, R.
32 Ortiz-Pulido, M. A. Marcos-García, and V. Rico-Gray. 2016. Unraveling Darwin's entangled bank: architecture and robustness
33 of mutualistic networks with multiple interaction types. *Proceedings of the Royal Society B* 283: 20161564.

Toju, H., P. R. Guimarães, Jr., J. M. Olesen, and J. N. Thompson. 2015. Plant communities and below-ground plant-fungal networks. *Sciences Advances* 1:e1500291

Toju, H., P. R. Guimarães, Jr., J. M. Olesen, and J. N. Thompson. 2014 Assembly of complex plant-fungal networks. *Nature Communications* 5:5273 DOI:10.1038/ncomms6273

Guimarães, P. R., Jr., P. Jordano, and J. N. Thompson. 2011. Evolution and coevolution in mutualistic networks. *Ecology Letters* 14:877-885.

Cuautle, M., and J. N. Thompson. 2010. Evaluating the co-pollinator network structure of two sympatric *Lithophragma* species with different morphology. *Oecologia* 162:71-80.

Díaz-Castelazo, C., P. R. Guimarães, Jr., P. Jordano, J. N. Thompson, R. J. Marquis, and V. Rico-Gray. 2010. Changes of a mutualistic network over time: reanalysis over a 10-year period. *Ecology* 91:793-801.

Guimarães, P. R., Jr., V. Rico-Gray, P.S. Oliveira, T. J. Izzo, S. F. dos Reis, and J. N. Thompson. 2007. Interaction intimacy affects structure and coevolutionary dynamics in mutualistic networks. *Current Biology* 17:1797-1803

Guimarães, P.R., V. Rico-Gray, S.F. dos Reis and J.N. Thompson (2006). Asymmetries in specialization in ant-plant mutualistic networks. *Proc. R. Soc. B* 273: 2041–2047.

Tylianakis 2010 Conservation of species interaction networks.

- Functions-Metrics (Figure 2:

- Richness/Connectance = Increased function and function stability

- Nestedness = Buffer extinctions in mutualistic networks

- Compartmentalization = greater stability, slow spread of disturbance (i.e. trophic cascades)

- Proportion of Weak Links = stability, fewer cascades

- Connectivity Distribution = Indicate Assembly, Robustness to 2nd extinctions

- Focus on metrics that saturate quickly with sampling

- Connectivity, Compartmentalization, Nestedness

- Need more research on the impacts of perturbations on these networks

Here, we investigate how genetic variation in a foundation tree species determines the structure of a network of interactions among species. Using a long-term (20 years+), common garden experiment with replicated individuals of known genetic identity and a naturally established stand of *P. angustifolia*. We focused on a model community of 9 epiphytic lichens species, as previous research has demonstrated significant compositional responses of epiphytes to genotypic variation^{5,16}. In addition, the life-history characteristics of lichen, having highly localized, direct contact interactions and slow population turnover rates,

63 allowed us to assess interactions among lichen species rapidly on individual trees. Applying a dual-scale (lichen-lichen and
64 genotype-lichen interactions) network analysis approach, we then examined the genetically based impacts of *P. angustifolia* on
65 network structure. We focused our network analysis on modularity (i.e. the formation of compartments) and nestedness (i.e. the
66 overlap in sets of interactions among species) both of which are of interest in the context of the interplay between ecological
67 and evolutionary dynamics as have established theoretical and empirical support as relevant structural metrics of ecological
68 dynamics and can serve as indicators of the units of evolution in a community context.

69 Results

70 Experimental Garden and Field Observations of Networks

71 In both the experimental garden and the natural stand, we discovered that genotypic variation in a *P. angustifolia* predictably
72 influenced the structure of the lichen species interaction network and contributed to the formation of evolutionary modules
73 comprised of tree genotypes and the lichen community. We observed significant unipartite (one-mode) network structure¹⁷ in
74 the lichen species interaction networks that was similar between the experimental garden and the natural stand (Fig. 1a and
75 1b; Garden: $z = -6.31$, $p = 0.0002$; Natural: $z = -3.15$, $p = 0.002$). The two networks displayed high multivariate structural
76 similarity (Mantel $R = 0.51$, $p = 0.029$). Node level eigen-centrality²¹, a measure of species importance that integrates indirect
77 connections, showed strong correlation between the two stands (Fig. 1c; $r = 0.7$, $t = 2.6135$, $df = 7$, $p = 0.035$). Centrality was
78 also highly correlated with total abundance in both networks (Fig. 1d; Garden: $r = 0.77$, $t = 3.2427$, $df = 7$, $p = 0.014$; Natural: r
79 $= 0.86$, $t = 4.43$, $df = 7$, $p = 0.003$). In combination, the similarity of both the whole and node level network structure between
80 the common garden and the wild indicates that the common garden environment captures much of the natural variation that
81 exists in nature and accurately reflects natural processes.

82 In the common garden, where the effect of environmental variation was controlled, genotype was an important factor
83 contributing to network structure. Genotype was a significant predictor of interactions on individual trees (Fig. 2a; $F = 3.4213$,
84 $num\ df = 12.000$, $denom\ df = 14.668$, $p\text{-value} = 0.01426$). Similar to the effect of a genetically controlled trait (bark roughness)
85 on a dominant lichen⁴, we found that individual tree genotypes with similar levels of bark roughness had similar levels of lichen
86 interactions (Fig. 2a; Mantel $R = 0.08$, $p = 0.013$), which was similar to the correlation observed between bark roughness and
87 lichen interactions in the natural stands (Fig 2b: $r = -0.53$, $p = 0.050$).

88 We also examined how *P. angustifolia* genotypic differentiation contributes to the formation of groups of tree genotypes
89 and lichen species and found significant modular structure. Using a bipartite (two-mode) network approach in which genotype-
90 species networks were modeled using the species maximum relativized values of each lichen species across all *P. angustifolia*
91 genotypes, we found significant modularity in the common garden stand (Fig. 3a; $z = 9.64$, $p < 0.001$). When using the same
92 analyses on individual trees in the natural stand, we also found significant modularity (Fig. 3b; $z = 7.42$, $p < 0.001$). Furthermore,
93 nestedness of both of these networks was significantly lower than expected under a null model (Garden: $z = -2.30$, $p < 0.001$;
94 Natural Stand: $z = -2.84$, $p < 0.001$), most likely as a result of module formation.

Discussion

These findings support the hypothesis that genotypic variation in a foundation species can contribute to the structure of a network of interacting species. Several lines of evidence support this conclusion. First, the wild stand showed significant interaction network structure (Fig. 1a and b); and both tree genotype and the genetically based tree trait, bark roughness, was a strong predictor of co-occurrence patterns (Fig. 2a). Second, the common garden network (Fig. 1b) structure showed a high degree of similarity to the wild stand network structure (Fig. 1c and d). Third, tree genotype was a significant predictor of SES values (Fig. 2a), displaying significant correlation with a genetically linked trait, bark roughness, both in the common garden (Fig. 2a) and in a naturally established stand of trees (Fig. 2b). Last, both of the bipartite genotype-species networks in the common garden and natural stand displayed significant modularity, suggesting that genotypic variation is leading to the formation of evolutionarily dynamic compartments within the community. Thus, just as numerous studies have shown that plant genotype can affect species richness, abundance, diversity, and composition and previous work has demonstrated that evolutionary processes shape ecological networks^{2,3}, our study includes genetics in an empirical investigation that combines both experimental common garden findings along with studies in the wild that are in close agreement.

Our results point to the importance of understanding the community level effects of genetic variation and corroborate previous findings of the importance of plant genetics in shaping ecosystems⁹. This study highlights the potential for indirect effects of genetic variation to propagate through networks of interacting species and trophic levels. Altering the structure of interaction networks presents a means for genetic effects to be magnified within the system of interacting species. At the scale of ecosystems, trophic networks or food webs direct and control the rates of energy and nutrient flux²². One important example², showed that the interactions among species across three trophic levels depended on cottonwood (*Populus* spp.) genotype.

- Networks working at multiple scales re-inforce connectivity
- Gene networks (Zink)
- Community networks (Keith2017, Lau 2016)
- Stand scale (This work)
- Landscape scale (Bothwell2017)

Although our study was conducted with a community of lichens, these results should be generalized to other groups of diverse organisms around the world that also exhibit significant genetic signals at the community level^{2,23}, although spatial scale of interactions should be considered²⁴. As heritable variation is the raw material for natural selection to act upon, a genetic basis for interaction network structure indicates evolutionary dynamics should be considered at the community level and that conserving genetic variation is important to consider in efforts to restore or preserve complex species interactions and the associated ecosystem functions²⁵ that they provide.

Methods

Field observations in common garden and natural riparian forest stands

The study was conducted along the Weber River, UT (USA), which is a cottonwood (*Populus* spp.) dominated riparian ecosystem. Although two native species, *Populus angustifolia* (James) and *Populus fremontii* (S. Watson), occur here and are known to hybridize, only pure or highly advanced backcrosses of *P. angustifolia* were sampled in order to avoid the effect of the hybridization between these two species.

A common garden setting was used to isolate the effect of tree genotype from the effect of the localized microenvironment associated with each individual and spatial autocorrelation. Asexually propagated clones of genotyped *P. angustifolia* individuals⁴ were obtained from wild collections and planted randomly in a single field (0.025 km²) at the Ogden Nature Center, Ogden, UT in 1992. A total of thirteen genotypes replicated between 3 and 8 times each, were chosen for sampling. Genotype names were previously published⁴. Observations were made in the common garden in October 2010 and May 2011.

The natural stand of *P. angustifolia* near the city of Uintah, UT (GPS: N41.13903, W110.94400) was used for the wild stand survey. We conducted sampling of the stand in May 2012. A total of 14 trees were chosen randomly over a 0.10 km² area with a minimal distance of 5.56 m between trees across a range of tree core based ages from 15 to 60 years.

Bark and Lichen Community Observations

On each tree, presence or absence of each lichen species was assessed in 50 total 1 cm² cells arrayed in a checkerboard pattern. Two adjacent 10 cm² quadrats centered at 50 cm and 85 cm from ground level were sampled. The checkerboard sampling pattern was chosen to isolate each cell based on an average thallus size of 1 cm². Samples were restricted to the northern aspect of the trunk to maximize the abundance of lichen and control for the effect of aspect. The thalli in each cell are expected to be spatially independent of the other cells in the quadrat, but exposed to similar micro-environmental conditions. Bark roughness was measured on each tree following⁵.

The bark lichen community in this system is comprised of fourteen species; however, only 9 species were observed within our study quadrats. The lichen community included (abbreviations are given for species present in study): Xg = *Xanthomendoza galericulata*, Xm = *X. montana*, Ch = *Caloplaca holocarpa*, Cs = *Candelariella subdeflexa*, Rg = *Rinodina glauca*, Lh = *Lecanora hagenii*, Ls = *Lecanora* (unknown species), Pm = *Phyciella melanchra*, Pa = *Phycia adscendens*, Pu = *Phycia undulata*, *Phaeophyscia orbicularis*, *Phaeophyscia ciliata*, *Melanelia subolivacea*, *Meanelia elegantula*, including both crustose and foliose lichen species that exhibit low inter-annual variation¹⁰. We were able to rapidly assess lichen interactions by quantifying thalli in closed contact as assessed using 1 cm² cells. Species accumulation curves showed that communities in the wild and the common garden were thoroughly sampled and with very similar species richness (Supplementary Materials).

Network modeling and analyses

We used the observations of lichen in the 1cm² cells on individual trees of *P. angustifolia* both in the common garden and the natural stand. Uni-partite networks were generated using an analytical procedure that removes non-significant interactions

between species¹⁷. We used a null model based approach for all other analyses of network structure. A conservative null model that constrained both the row and column marginal totals was used in order to account for the effects of variation in species' total abundances²⁴. From a total of 5000 null matrices, a standardized score was calculated for each statistic ($z = \frac{x_{obs} - \bar{x}_{sim}}{sd_{sim}}$), including the C-score¹⁸, nestedness¹⁹ and modularity²⁰. Here, we follow the convention of the co-occurrence literature and refer to the standardized C-Score as the Standardized Effect Size (SES)².

A correlation test with Pearson's r was used to test for the correlation between the wild and common garden networks. A Welch Analysis of Variance (ANOVA), which relaxes the assumption of homogeneity of variance, was used to test for the effects of genotype on tree scale SES values. A permutation based Mantel Test was used to test for the effect of bark roughness on SES values in the common garden. All analyses were conducted using the programming language R².

References

1. Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P. & Bascompte, J. Non-random coextinctions in phylogenetically structured mutualistic networks. *Nat.* **448**, 925–8 (2007). URL <http://dx.doi.org/10.1038/nature05956>. DOI 10.1038/nature05956.
2. Guimarães, P. R., Jordano, P. & Thompson, J. N. Evolution and coevolution in mutualistic networks. *Ecol. Lett.* **14**, 877–85 (2011). URL <http://www.ncbi.nlm.nih.gov/pubmed/21749596>. DOI 10.1111/j.1461-0248.2011.01649.x.
3. Moya-Laraño, J. Genetic variation, predator-prey interactions and food web structure. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **366**, 1425–37 (2011). DOI 10.1098/rstb.2010.0241.
4. Ellison, A. M. *et al.* Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Front. Ecol. Environ.* **3**, 479–486 (2005). URL [http://www.esajournals.org/doi/abs/10.1890/1540-9295\(2005\)003\[0479:LOFSCF\]2.0.CO;2?journalCode=fron](http://www.esajournals.org/doi/abs/10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2?journalCode=fron). DOI 10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2.
5. Fontaine, C. *et al.* The ecological and evolutionary implications of merging different types of networks. *Ecol. Lett.* **14**, 1170–81 (2011). URL <http://www.ncbi.nlm.nih.gov/pubmed/21951949>. DOI 10.1111/j.1461-0248.2011.01688.x.
6. Bascompte, J., Jordano, P. & Olesen, J. M. Asymmetric Coevolutionary Networks Facilitate Biodiversity Maintenance. *Sci.* (80-.). **312**, 431–433 (2006). URL <http://www.sciencemag.org/cgi/content/abstract/312/5772/431>. DOI 10.1126/science.1123412.
7. Rafferty, N. E. & Ives, A. R. Phylogenetic trait-based analyses of ecological networks. *Ecol.* **94**, 2321–33 (2013).
8. Whitham, T. G. *et al.* A framework for community and ecosystem genetics: from genes to ecosystems. *Nat. Rev. Genet.* **7**, 510–523 (2006). DOI 10.1038/nrg1877.

- 188 **9.** Lamit, L. *et al.* Genetically-based trait variation within a foundation tree species influences a dominant bark
189 lichen. *Fungal Ecol.* **4**, 103–109 (2011). URL [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S1754504810000668)
190 [S1754504810000668](http://www.sciencedirect.com/science/article/pii/S1754504810000668). DOI 10.1016/j.funeco.2010.09.007.
- 191 **10.** Bailey, J. K. *et al.* From genes to ecosystems: a synthesis of the effects of plant genetic factors across levels of organi-
192 zation. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **364**, 1607–16 (2009). URL [http://www.pubmedcentral.nih.](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2690499)
193 [gov/articlerender.fcgi?artid=2690499](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2690499){&}tool=pmcentrez{&}rendertype=abstract. DOI
194 10.1098/rstb.2008.0336.
- 195 **11.** Holeski, L. M., Hillstrom, M. L., Whitham, T. G. & Lindroth, R. L. Relative importance of genetic, ontogenetic, induction,
196 and seasonal variation in producing a multivariate defense phenotype in a foundation tree species. *Oecologia* **170**, 695–707
197 (2012). DOI 10.1007/s00442-012-2344-6.
- 198 **12.** Whitham, T. G. *et al.* Community specificity: Life and afterlife effects of genes (2012). DOI 10.1016/j.tplants.2012.01.005.
- 199 **13.** Burkle, L. A., Souza, L., Genung, M. A. & Crutsinger, G. M. Plant genotype, nutrients, and G x E interactions structure
200 floral visitor communities. *Ecosphere* **4**, art113 (2013). URL <http://dx.doi.org/10.1890/ES13-00039.1>.
201 DOI 10.1890/ES13-00039.1.
- 202 **14.** Smith, D. S., Bailey, J. K., Shuster, S. M. & Whitham, T. G. A geographic mosaic of trophic interactions and selection: trees,
203 aphids and birds. *J. Evol. Biol.* **24**, 422–9 (2011). URL <http://www.ncbi.nlm.nih.gov/pubmed/21091573>.
204 DOI 10.1111/j.1420-9101.2010.02178.x.
- 205 **15.** Winfree, R., Gross, B. J. & Kremen, C. Valuing pollination services to agriculture. *Ecol. Econ.* **71**, 80–88
206 (2011). URL <http://www.sciencedirect.com/science/article/pii/S092180091100334X>. DOI
207 10.1016/j.ecolecon.2011.08.001.
- 208 **16.** Zytynska, S. E., Fay, M. F., Penney, D. & Preziosi, R. F. Genetic variation in a tropical tree species influences the associated
209 epiphytic plant and invertebrate communities in a complex forest ecosystem. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **366**,
210 1329–1336 (2011). DOI 10.1098/rstb.2010.0183.
- 211 **17.** Araújo, M. B., Rozenfeld, A., Rahbek, C. & Marquet, P. A. Using species co-occurrence networks to assess the impacts of
212 climate change. *Ecography (Cop.)*. **34**, 897–908 (2011). URL [http://dx.doi.org/10.1111/j.1600-0587.](http://dx.doi.org/10.1111/j.1600-0587.2011.06919.x)
213 [2011.06919.x](http://dx.doi.org/10.1111/j.1600-0587.2011.06919.x){&}5Cnhttp://doi.wiley.com/10.1111/j.1600-0587.2011.06919.x. DOI
214 10.1111/j.1600-0587.2011.06919.x.
- 215 **18.** DeAngelis, D. L., Bartell, S. M. & Brenkert, A. L. Effects of Nutrient Recycling and Food-Chain Length on Resilience
216 (1989). DOI 10.1086/285011.
- 217 **19.** Borgatti, S. P. & Everett, M. G. A Graph-theoretic perspective on centrality. *Soc. Networks* **28**, 466–484 (2006). DOI
218 10.1016/j.socnet.2005.11.005.

- 219 **20.** Rowntree, J. K., Shuker, D. M. & Preziosi, R. F. Forward from the crossroads of ecology and evolution. *Philos. Trans. R.*
220 *Soc. Lond. B. Biol. Sci.* **366**, 1322–8 (2011). URL [http://rstb.royalsocietypublishing.org/content/](http://rstb.royalsocietypublishing.org/content/366/1569/1322.full)
221 [366/1569/1322.full](http://rstb.royalsocietypublishing.org/content/366/1569/1322.full). DOI 10.1098/rstb.2010.0357.
- 222 **21.** Zook, A. E., Eklof, A., Jacob, U. & Allesina, S. Food webs: Ordering species according to body size yields high degree of
223 intervality. *J. Theor. Biol.* **271**, 106–113 (2010). URL [http://www.sciencedirect.com/science/article/](http://www.sciencedirect.com/science/article/pii/S0022519310006405)
224 [pii/S0022519310006405](http://www.sciencedirect.com/science/article/pii/S0022519310006405). DOI 10.1016/j.jtbi.2010.11.045.
- 225 **22.** Evans, D. M., Pocock, M. J. O. & Memmott, J. The robustness of a network of ecological networks to habitat loss. *Ecol.*
226 *Lett.* **16**, 844–52 (2013). URL <http://www.ncbi.nlm.nih.gov/pubmed/23692559>. DOI 10.1111/ele.12117.
- 227 **23.** Stone, L. & Roberts, A. The checkerboard score and species distributions. *Oecologia* **85**, 74–79 (1990). DOI
228 10.1007/BF00317345.
- 229 **24.** Atmar, W. & Patterson, B. D. The measure of order and disorder in the distribution of species in fragmented
230 habitat. *Oecologia* **96**, 373–382 (1993). URL <http://link.springer.com/10.1007/BF00317508>. DOI
231 10.1007/BF00317508.
- 232 **25.** Newman, M. E. J. Modularity and community structure in networks. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 8577–82 (2006).
233 URL <http://www.pnas.org/content/103/23/8577.short>. DOI 10.1073/pnas.0601602103.

234 **Acknowledgments**

235 This work was supported by the National Science Foundation grant (DEB-0425908) and Integrative Graduate Research
236 Traineeship (IGERT) fellowships for M.L. and L.L. The Ogden Nature Center staff helped to maintain the common gardens.
237 Lichen sampling was supported by Todd Wojtowicz, Luke Evans and David Solance Smith.

238 **Author contributions statement**

239 M.L. and L.L. conceived the study, M.L. and L.L. conducted the field work, R.N. assisted in lichen identifications, M.L. wrote
240 the first draft of the manuscript, S.B. and T.W. contributed substantively to the conceptual development, T.W. established the
241 common garden. All authors contributed to revisions of the manuscript.

242 **Additional information**




Figure 1. Significant unipartite network structure was observed for epiphytic lichens on trees of known genotype in a common garden (ONC = Ogden Nature Center, Utah, USA) (A) and individual trees in a natural stand (Uintah, Utah, USA) (B) of the foundation species, *Populus angustifolia*. Both networks are shown here with lichen species as nodes (see Methods for complete species names) scaled by the log of their total abundances and significant co-occurrence patterns between species shown as edges scaled by their log frequencies. The bivariate plot (C) shows the significant correlation in Eigen Centrality between the two networks. (D) The total abundance of lichen species was a significant driver of network structure for both networks.




Figure 2. Tree genotype influenced lichen co-occurrence patterns in the common garden and the natural stand through a genetically controlled tree trait. The lichen co-occurrence patterns were highly correlated with the genetically based phenotypic trait; bark roughness (i.e., the percentage of textured bark), in both the common garden and natural stand. The scatterplot (A) shows the mean (± 1 SE) percent rough bark (broad-sense heritability, $H^2 = 0.36$, $\chi^2 = 9.214$, $p = 0.002$) and SES for each genotype for trees in the common garden with SES values becoming more negative (i.e., species interactions increased), indicating stronger co-occurrence patterns, as bark roughness increases. The lichen communities on individual trees in the Unitah natural stand (B) displayed a similar pattern with the SES values becoming increasingly more negative on trees with more rough bark.




Figure 3. Bipartite networks displayed significant modularity with modules comprised of both genotypes and species. The left most set of nodes shows tree genotypes (see Methods for genotype names) for the common garden (A) or individuals in the natural stand (B) connected to lichen species on the right. Both sets of nodes are scaled by their marginal totals (i.e., total observed individuals for tree nodes and total abundance for lichen species) and arranged by ascending totals from bottom to top. Node color shows the significant module membership for both trees and lichen species with module color having no direct relationship between the two networks, as modules were determined for each network independently.