


Stereo-electroencephalography Brain-computer Interface Study



**From BioMechatronics and BioRobotics Laboratory
of Shanghai Jiaotong University**
<http://bbl.sjtu.edu.cn/>

Tao Xie, Zehan Wu|2016.03.25








- **Day one:** Only use BP to detect signal, the result is good;
- **Day two:** Detect signal with Neuroscan, system is not stable, and the result is not good;
- **Day three:** with Neuroscan again, system is stable.

Self motion








Subject perform brisk wrist motion with 5~10 sec interval.
sEEG shows significant ERD/S and MRCP only for the
electrode within motorsensory area.

● Cue real motion (black: keep 1 sec; red: keep 3 sec)




● Cue imagery motion (black: keep 1 sec; red: keep 3 sec)




● Vibration

Chan_12 (vib with motion)



Chan_12 (vib only)



Anyhow, thank **Shize Jiang**, and we have a happy work together!

- Neuroscan system
- g.tec system

Online and offline study

日本光电脑电采集系统

Offline study

Different electrodes location focus on different study point

- Self-motion, cue-motion (real & imagery)
- Force level study
- Vib-stimulation
- Visual-stimulation



Contents lists available at [SciVerse ScienceDirect](#)

NeuroImage

journal homepage: www.elsevier.com/locate/ynimg

Toward a minimally invasive brain–computer interface using a single subdural channel: A visual speller study

Dan Zhang ^{a,1}, Huaying Song ^{a,1}, Rui Xu ^a, Wenjing Zhou ^c, Zhipei Ling ^{b,*}, Bo Hong ^{a,**}

^a Department of Biomedical Engineering, School of Medicine, Tsinghua University, Beijing 100084, China

^b Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China

^c Department of Neurosurgery, Affiliated Yuquan Hospital, School of Medicine, Tsinghua University, Beijing 100084, China

Thanks