

Situational awareness and forecasting

FHI COVID-19 modelling team

16 April 2020

What this report contains:

This report presents results based on a mathematical model describing the geographical spread of COVID-19 in Norway. The model consists of three layers:

- Population structure in each municipality
- Mobility data for inter-municipality movements (Telenor mobile phone data)
- Infection transmission model

The model produces estimates of the current epidemiological situation at the municipality, county (fylke) and national levels, a forecast of the situation for the next three weeks and a long term prediction.

How we calibrate the model:

The model is fitted to Norwegian COVID-19 hospitalization prevalence data since March 10 until today. We seed new infections into the model using imported COVID-19 cases in Norway from February 26 until March 18.

How you should interpret the results:

The model is stochastic (random). To predict the probability of various outcomes, we run the model many times. We present the results in terms of mean values, 95% confidence intervals, median and interquartile ranges. We underline that the confidence bands presented might be broader: First, there are uncertainties related to the natural history of SARS-CoV-2, including the importance of asymptomatic and presymptomatic infection. Second, there are uncertainties related to the timing of hospitalization relative to symptom onset, the severity of the COVID-19 infections by age, and the duration of hospitalization and ventilator treatment in ICU. These uncertainties are not fully explored in the present results. We will update the model parameters in accordance with new evidence and local data as they becomes available and results can change also significantly. See more details at the end of this report.

The mobility data is updated until April 13. It accounts for the changes in the movement patterns between municipalities that have occurred since March 12.

In the forecasting, we use the reproduction numbers that fit the hospitalization data best. The basic reproductive number, R_0 , is used until March 14. A new effective reproductive number, R_{eff} , acts from March 15 until today, and in the future when we predict. R_0 is calibrated to hospitalization data (number of occupied beds) until March 20; R_{eff} from March 21 until today. We weight the last days (from March 28) more than the first days. We use a distribution of all reproduction numbers to guarantee appropriate uncertainty of their estimates. However, uncertainties related to the model parameters, as well as the transient period in week 11 imply that reported effective reproductive values should be interpreted with caution. We notice that our predictions of intensive care prevalence are

slightly overestimated compared to current data. We will update the parameters related to permanence in hospital and ICU as soon as NPR data will be linked with MSIS.

Note that, in this report, the term ventilator treatment or ICU includes only those patients that require either invasive mechanical ventilation or ECMO (Extracorporeal membrane oxygenation).


1 Estimated Reproductive Numbers

Calibration of our model with hospitalisation data leads to the following estimates:


Table 1: Calibration results

Parameter	Mean	Median	Confidence interval (95 %)
Amplification factor	2.32	2.33	(1.71-2.96)
R_0	3.06	3.06	(2.64-3.64)
R_{eff}	0.68	0.65	(0.5-0.83)

Estimated densities of these three parameters are plotted below:



Our model estimates the number of hospitalised Covid-19 patients, plotted below with blue median and interquartile bands, which are compared with the actual hospitalisation data, in red. The uncertainty captures the uncertainty in the calibrated parameters in addition to the stochastic elements of our model.



True total number of hospitalisations (red) and predicted values (blue)

2 Estimated cumulative number of infected individuals

Table 2: Estimated cumulative number of infections, 2020-04-16

Region	Total	Symptomatic	No. confirmed	Fraction reported	Min. fraction
Norway	45801 (38671; 53308)	28045 (23850; 32465)	6782	15%	13%
Agder	2807 (2014; 3737)	1718 (1224; 2288)	275	10%	7%
Innlandet	2334 (1646; 3297)	1417 (982; 2003)	403	17%	12%
Møre og Romsdal	910 (575; 1292)	563 (354; 804)	118	13%	9%
Nordland	738 (434; 1136)	452 (269; 698)	104	14%	9%
Oslo	10615 (8906; 12523)	6462 (5441; 7609)	2031	19%	16%
Rogaland	6093 (4750; 7707)	3731 (2918; 4708)	383	6%	5%
Troms og Finnmark	1349 (662; 2444)	821 (419; 1495)	203	15%	8%
Trøndelag	2044 (1449; 2803)	1252 (885; 1694)	406	20%	14%
Vestfold og Telemark	3729 (2574; 5664)	2271 (1592; 3427)	251	7%	4%
Vestland	4926 (3654; 6408)	3005 (2230; 3923)	743	15%	12%
Viken	10255 (8393; 12367)	6354 (5216; 7619)	1864	18%	15%

Fraction reported=Number confirmed/number predicted; Minimal fraction reported=number confirmed/upper CI

3 Predicted incidence of infected individuals, next three weeks

Predicted incidence (asymptomatic and symptomatic) for Norway per day, with confidence intervals.




Table 3: Predicted incidence per day.

Region	1 week prediction (23 April)	2 weeks prediction (30 April)	3 weeks prediction (07 May)
Norway	354 (134-684)	268 (85-606)	204 (56-524)
Agder	22 (7-45)	17 (4-41)	13 (2-36)
Innlandet	24 (8-51)	18 (5-41)	14 (2-37)
Møre og Romsdal	8 (1-20)	6 (0-17)	5 (0-15)
Nordland	6 (1-16)	5 (0-14)	4 (0-11)
Oslo	57 (23-112)	44 (13-97)	33 (8-80)
Rogaland	44 (14-91)	34 (10-78)	26 (6-66)
Troms og Finnmark	10 (1-25)	8 (1-21)	6 (0-18)
Trøndelag	18 (5-37)	13 (2-31)	10 (1-27)
Vestfold og Telemark	29 (9-58)	22 (6-50)	17 (3-44)
Vestland	39 (14-77)	30 (8-67)	23 (5-56)
Viken	97 (38-189)	73 (22-165)	55 (14-139)


4 Predicted hospitalisation, next three weeks, including patients in ventilator treatment

Table 4: Number of hospitalisation beds occupied by Covid-19 patients.

Region	1 week prediction (23 April)	2 weeks prediction (30 April)	3 weeks prediction (07 May)
Norge	135 (78-206)	101 (49-172)	77 (32-145)
Agder	8 (1-18)	6 (0-17)	5 (0-13)
Innlandet	9 (1-20)	7 (1-17)	6 (0-15)
Møre og Romsdal	3 (0-9)	2 (0-9)	2 (0-7)
Nordland	3 (0-8)	2 (0-7)	2 (0-6)
Oslo	26 (10-46)	17 (4-33)	11 (2-25)
Rogaland	17 (5-31)	12 (3-26)	9 (1-21)
Troms og Finnmark	4 (0-12)	3 (0-10)	2 (0-7)
Trøndelag	6 (0-15)	5 (0-12)	4 (0-12)
Vestfold og Telemark	13 (3-26)	10 (1-22)	7 (0-17)
Vestland	15 (4-29)	11 (2-24)	8 (1-20)
Viken	32 (14-52)	26 (9-46)	21 (6-44)

Yesterday's real value for Norway: 175

Predicted daily number of COVID-19 patients in hospital in Norway (95% confidence intervals and interquartile range), next three weeks, including patients ventilator treatment.



Similar table and figure for each county (fylke) available on request.


5 Predictive number of patients in ventilator treatment: next three weeks

Table 5: Number of ICU beds occupied by Covid-19 patients.

Region	1 week prediction (23 April)	2 weeks prediction (30 April)	3 weeks prediction (07 May)
Norge	51 (31-76)	38 (20-63)	29 (13-51)
Agder	3 (0-7)	2 (0-7)	2 (0-5)
Innlandet	4 (0-8)	3 (0-7)	2 (0-6)
Møre og Romsdal	1 (0-4)	1 (0-4)	1 (0-3)
Nordland	1 (0-3)	1 (0-3)	1 (0-3)
Oslo	9 (3-16)	6 (1-13)	4 (0-9)
Rogaland	6 (2-12)	5 (1-10)	3 (0-8)
Troms og Finnmark	1 (0-5)	1 (0-4)	1 (0-3)
Trøndelag	2 (0-6)	2 (0-5)	1 (0-5)
Vestfold og Telemark	5 (1-11)	4 (0-9)	3 (0-7)
Vestland	6 (1-12)	4 (0-9)	3 (0-8)
Viken	12 (5-19)	9 (3-17)	8 (2-16)

Yesterday's real value for Norway: 52

Predicted daily number of COVID-19 patients in ventilator treatment in Norway (95% confidence intervals and interquartile range), next three weeks.



Similar table and figure for each county (fylke) available on request.

6 Predicted prevalence of infectious individuals, next three weeks:

Predicted daily prevalence of asymptomatic, presymptomatic and symptomatic individuals, aggregated, whole Norway, (95% confidence interval).





Table 6: Predicted prevalence. Number of infectious individuals (asymptomatic plus pre-symptomatic plus symptomatic) per day. Means and 95 perc. CI for three weeks prediction.


Region	Mean, 23 April	Mean, 30 April	Mean, 07 May	low CI, 07 May	high CI, 07 May
Norway	2586	1953	1483	445	3519
Agder	158	120	93	26	223
Innlandet	173	131	101	29	229
Møre og Romsdal	58	45	35	7	89
Nordland	46	34	26	5	68
Oslo	422	318	241	71	556
Rogaland	325	245	185	54	421
Troms og Finnmark	74	56	43	9	114
Trøndelag	127	96	73	19	171
Vestfold og Telemark	209	158	120	37	278
Vestland	286	216	164	50	365
Viken	708	535	403	122	906

Map of predicted prevalence. Number of infectious individuals (asymptomatic plus presymptomatic plus symptomatic) today in each municipality.



7 Mobility between municipalities

Number of trips out from each municipality during each day, based on Telenor mobility data. We have observed a large reduction in inter-municipality mobility in week 11 (around March 11), with a minimum reached on Tuesday 17 March. The reduction with respect to the weeks before (for example week 10) is on average 50%. Thereafter, we observe a slight increasing trend: in Oslo, for example, out-mobility has increased of roughly 2% per day in the last three weeks. Weekends still have a lower mobility, indicating that there is still some commuting-to-job during weekdays.




Percentage reduction in total mobility out from each municipality: Monday April 6th is compared to Monday March 2, which is the last Monday before interventions (March 12), etc.



	Man_06	Tirs_07	Ons_08	Tors_09	Fre_10
Hele Norge	34%	32%	40%	44%	44%
Oslo	55%	53%	57%	60%	63%
Bergen	30%	28%	39%	41%	38%
Trondheim	37%	33%	49%	66%	69%
Stavanger	41%	37%	47%	47%	48%
Bærum	53%	52%	56%	56%	55%
Kristiansand	34%	32%	40%	45%	58%
Drammen	40%	39%	43%	51%	50%
Asker	39%	37%	42%	41%	39%
Lillestrøm	44%	47%	51%	62%	59%
Fredrikstad	28%	23%	34%	29%	31%
Sandnes	41%	38%	48%	53%	51%
Tromsø	40%	49%	55%	63%	77%
Ålesund	23%	21%	37%	48%	41%
Sandefjord	34%	29%	39%	40%	45%
Nordre Follo	42%	41%	47%	49%	49%
Sarpsborg	28%	30%	36%	42%	42%
Tønsberg	35%	34%	40%	48%	48%
Skien	33%	29%	38%	41%	42%
Bodø	50%	52%	60%	53%	71%
Moss	33%	29%	39%	40%	37%

8 Long-term prediction results

Predicted daily number of COVID-19 patients in hospital and receiving ventilator treatment in Norway until the end April, 2021, in addition to prevalence. The figures are made using 10 runs of each of the 100 candidate models, where the R_{eff} , from the March 15 and onward ranges between 0.50 and 0.85, with interquartile range 0.65-0.70 and median 0.65.

Note that the confidence intervals reflected on the plots are two tailed around the median, and therefore the upper 95 % level shows the 97.5 % boundary.






None of the simulations exceeded the surge capacity need of **500 ventilator treatments**.

9 Long-term scenario results

Here we show how the epidemics will develop, from April 13th if the reproductive number would increase to 1.15, 1.3 and 1.5 respectively. We show the daily number of covid-19 patients with ventilator treatment.




Model

We use a metapopulation model to simulate the spread of COVID-19 in Norway in space and time. The model consists of three layers: the population structure in each municipality, information about how people move between different municipalities, and local transmission within each municipality. In this way, the model can simulate the spread of COVID-19 within each municipality, and how the virus is transported around in Norway.

Transmission model

We use an SEIR (Susceptible-Exposed-Infected-Recovered) model without age structure to simulate the local transmission within each area. Mixing between individuals is assumed random. Demographic changes due to births, immigration, emigration and deaths are not considered. The model distinguishes between asymptomatic and symptomatic infection, and we consider presymptomatic infectiousness among those who develop symptomatic infection. In total, the model consists of 6 disease states: Susceptibles (S), Exposed, infected, but not infectious (E), Presymptomatic infected (E2), Symptomatic infected (I), Asymptomatic infected (Ia), and Recovered, either immune or dead (R). A schematic overview of the model is shown below:



Movements between municipalities:

We use 6-hourly mobility matrices from Telenor to capture the movements between municipalities. The matrices are scaled according to the overall Telenor market share in Norway, estimated at 48%. Since week 8, we use the actual daily mobility matrices to simulate the past. In this way, alterations in the mobility pattern will be incorporated in our model predictions. To predict future movements, we use the latest weekday measured by Telenor. We follow closely the development in the mobility matrices, and weekend patterns will be introduced if needed.

Healthcare utilization

Based on the daily incidence data from the model and the population age structure in each municipality, we calculated the hospitalization using a weighted average. The hospitalization is assumed delayed relative to symptom onset. We calculate the number of patients admitted to ventilator treatment from the patients in hospital using age adjusted probabilities and an assumed delay.

Seeding

At the start of each simulation, we locate 5.367.580 people in the municipalities of Norway according to data from SSB per January 1. 2020. All confirmed Norwegian imported cases with information about residence municipality and test dates are used to seed the model, until 18th March. For each case, we add an additional random number of infectious individuals, in the same area and on the same day, to account for asymptomatic imported cases who were not tested or others missed.

Reproduction number and calibration

We estimate the reproduction number R_0 so that the predicted mean number of hospitalized individuals is closest to the true number of hospitalized individuals, from March 10 until March 20. Restrictions introduced on March 12 require a change in the effective reproduction number. The effective reproductive number R_{eff} is active from March 15 until today, and in the future when we predict. R_0 is calibrated to hospitalisation data (number of occupied beds) until March 20; R_{eff} from March 21 until today. We weight the last days (from March 28) more than the first days. We estimate R_0 by selecting the 100 best R_0 s out of 10000 random candidates (each simulated 10 times). For these 100 best, we find by grid search the R_{eff} that fit best the hospitalisation data. Then we run 10 simulations of the future for each of these 100 best (R_0, R_{eff}) .

Parameters used today

Table 7: Assumptions I

Assumptions	Mean	Distribution	Reference
Seeding			
Scaling factor on imported cases	Min. 1.49 1st Qu. 2.11 Median 2.33 Mean 2.32 3rd Qu. 2.53 Max. 3.14	random	Calibrated to hospitalizations together with R_0
Telenor coverage	48%		https://ekomstatistikken.nkom.no//statistics/
Model parameters			
Exposed period ($1/\lambda_1$)	3 days	Exponential	Fraser et al. Not published
Pre-symptomatic period ($1/\lambda_2$)	2 days	Exponential	Fraser et al. Not published
Symptomatic infectious period ($1/\mu$)	5 days	Exponential	Fraser et al. Not published
Asymptomatic, infectious period ($1/\mu$)	5 days	Exponential	Fraser et al. Not published
Infectiousness asympt. (r_{I_a})	0.1	Fixed	Fraser et al. Not published
Infectiousness presymp (r_{E_2})	1.25	Fixed	Fraser et al. Not published
Prob. asymptomatic infection (p_a)	0.4		Fraser et al. Not published
R_0 (until March 14)	Min. 2.46 1st Qu. 2.88 Median 3.06 Mean 3.06 3rd Qu. 3.22 Max. 3.86	random	Calibrated to hospitalizations
R_{eff} , from 15 March until today	Min. 0.50 1st Qu. 0.65 Median 0.65 Mean 0.68 3rd Qu. 0.70 Max. 0.85	random	Calibrated to hospitalizations


Table 8: Assumptions II

Assumptions	Mean	Distribution	Reference
Healthcare			
Time sympt. onset to hospitalisation	8 days	Poisson	
Fraction asymptomatic infections	40%	Fixed	Mizumoto et al 2020 20% for the old population, Diamond Princess
% symptomatic and asymptomatic infections requiring hospitalization:			Verity et al 2020 corrected for: % of elderly living in of elderly living in Norway (last two age groups). Also corrected by 1/3 to account for severity in comparison with the expected fatality rate.
0-9 years	0.00%	Fixed	
10 - 19 years	0.013%		
20 - 29 years	0.37%		
30 - 39 years	1.13%		
40 - 49 years	1.43%		
50 - 59 years	2.73%		
60 - 69 years	3.93%		
70 - 79 years	5.53%		
80+ years	5.33%		
% hospitalized patients requiring ICU			
0-9 years	5%	Fixed	Verity et al 2020
10 - 19 years	5%		
20 - 29 years	5%		
30 - 39 years	5%		
40 - 49 years	6.3%		
50 - 59 years	12.2%		
60 - 69 years	27.4%		
70 - 79 years	43.2%		
80+ years	70.9%		
Overall hospitalization risk	1.9%	Fixed	Verity et al 2020 (adapted to Norwegian population)
Normal hospitalization length	8 days	Poisson	Ferguson et al 2020
Time in hospital before ICU	4 days	Poisson	Ferguson et al 2020, Expert opinion
Time in ICU	12 days	Poisson	Ferguson et al 2020, Expert opinion
Mobile phone mobility			
Until April 13	Measured Telenor mobility		
Data used in the predictions	April 7	Fixed	Corrected to preserve population

Supplementary analysis: Instantaneous Reproduction Number based on lab-confirmed cases only, EpiEstim

These following results are based on confirmed cases and should be interpreted very carefully due to the multiple changes in testing criteria during this period. During the early part of the period, testing was mainly based on travel to areas with an ongoing outbreak, while the last two weeks have changed to testing people with an acute respiratory infection. The fact that the reproduction number is close to or below one corresponds to the break in the exponential growth in confirmed cases. The last two weeks have seen a fairly constant rate of new confirmed cases.

Using the date of onset of the confirmed cases from the outbreak registry we estimate the instantaneous reproduction number using the procedure outlined in Thompson et Al (2019). This method, implemented in the EpiEstim R-package uses a Bayesian method to estimate the instantaneous reproduction number smoothed over a sliding window of 7 days. For data without onset date, we estimate an onset date by subtracting 7 days, which is the mean delay between the date of onset and reporting date in the data, from the reporting date. The serial interval was assumed at 5 days with uncertainty; the serial interval refers to the time between symptom onset between successive cases in a chain of transmission (see <https://www.medrxiv.org/content/10.1101/2020.02.03.20019497v2>). To account for censoring of observations with onset dates in the last few days we correct the observed data by the mean of a negative binomial distribution with observation probability given by the empirical cumulative distribution of the onset to reporting date distributions. Due to this correction, the results from the last few days should be interpreted with caution. These dates are indicated by the weaker colours.



FHI COVID-19 modelling team:

- **Birgitte Freiesleben De Blasio** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.
- **Francesco Di Ruscio** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.
- **Gunnar Øyvind Isaksson Rø** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.
- **Solveig Engebretsen** - Norsk Regnesentral.
- **Arnoldo Frigessi** - Oslo Centre for Biostatistics and Epidemiology, University of Oslo and Oslo University Hospital
- **Alfonso Diz-Lois Palomares** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.
- **Kenth Engø-Monsen** - Telenor Research.
- **Richard White** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.
- **Gry Marysol Grøneng** - Department of Infectious Disease Epidemiology and Modelling. Folkehelseinstituttet.