


EDS 212: Day 2, Lecture 2

Derivatives

August 4th, 2025


Introduction to Derivatives

Recall average and instantaneous rate of change



Big idea: Taking the average rate of change to a set limit will eventually converge to the instantaneous.

Average slope




Start with:

Nearby point:

Average slope:

Average slope



Start with a point $(x, f(x))$ on the graph.


A nearby point can be $(x + \Delta x, f(x + \Delta x))$, where Δx is a short distance from x .

The average slope between $(x, f(x))$ and $(x + \Delta x, f(x + \Delta x))$ is given by


$$\text{average slope} = \frac{f(x + \Delta x) - f(x)}{(x + \Delta x) - x} = \frac{f(x + \Delta x) - f(x)}{\Delta x}.$$

What happens when Δx becomes smaller and smaller?


What happens when $\Delta x \rightarrow 0$?




What happens when $\Delta x \rightarrow 0$?



What happens when $\Delta x \rightarrow 0$?




What happens when $\Delta x \rightarrow 0$?



Tangent lines

What if we let $\Delta x = 0$?

Then we would have a slope line that only touches the graph at exactly x .



These very special types of lines are called **tangent lines**.

Derivative definition

Remember the slope of the line that passes between two points (x_1, y_1) and (x_2, y_2) is:

$$\text{slope} = \frac{y_2 - y_1}{x_2 - x_1}.$$

If we have a function $f(x)$, then $(x, f(x))$ represents a point on its graph.

Choose a different point on the graph that is Δx away: $(x + \Delta x, f(x + \Delta x))$.

Plug these points on the graph of $f(x)$ into the slope equation:

$$\text{slope} = \frac{f(x + \Delta x) - f(x)}{\Delta x}.$$

When we let $\Delta x \rightarrow 0$ we get **the slope of the tangent line at x**

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

This is the derivative of f at x . Also denoted $\frac{df}{dx}$.

The expression for the instantaneous slope at any point on a function, aka the **derivative**

IS FOUND BY:

- ① Finding an expression for the **slope** between 2 points separated by Δx ...

$$\frac{df}{dx} = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x}$$

- ② evaluating that slope as the points get infinitely close together.




Not all functions are differentiable

- All differentiable functions are continuous, but not all continuous functions are differentiable.
- The absolute value function $|x|$ is one example, this is defined by

$$|x| = \begin{cases} x & , x \geq 0 \\ -x & , x < 0 \end{cases} .$$

It's graph is:



Let's take a 5 minute break




image: Flaticon.com

Why derivatives in an EDS program?

Describing rates of change is common in environmental science
(rate of pollutant concentration change, rate of population
growth, rate of energy consumption)


Calculus and derivatives are used to study change



Environmental Science studies change



Environmental Science studies change



Environmental Science studies change



🤔 Talk with a partner, what are 2 fields of environmental science where studying the rate of change and derivatives would be important?

Rules for differentiation (1)

Constant Rule 

Power Rule 

Rules for differentiation (1)

Constant Rule

Let c be a constant (so, a number). If $f(x) = c$, then $f'(x) = 0$. Equivalently

$$\frac{d}{dx}c = 0$$

Power Rule

Let n be a (real) number, if $f(x) = x^n$, then $f'(x) = nx^{n-1}$. Equivalently

$$\frac{d}{dx}[x^n] = nx^{n-1}.$$

Examples

$$y = 100 \qquad f(x) = x^5 \qquad f(x) = \frac{1}{x^2}$$

Rules for differentiation (2)

Sum and Difference Rules



Rules for differentiation (2)

Sum and Difference Rules

Let $f(x)$ and $g(x)$ be differentiable functions. Then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$
$$\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

😵 All this says is: if the function has pieces that are added or subtracted you can take the derivative of each individual piece and add those derivatives.

Example 

$$f(x) = x^3 - x^2 - 15$$

Rules for differentiation (3)

Constant Multiplier Rule



Rules for differentiation (3)

Constant Multiplier Rule

The derivative of a constant c multiplied by a function $f(x)$ is the same as the constant multiplied by the derivative:

$$\frac{d}{dx}[kf(x)] = k\frac{d}{dx}f(x)$$

Example

$$y = x^2 + \sqrt{2}x - \frac{8}{x} + 4$$

Derivative of logs & exponents

Two special derivatives:

- $\frac{d}{dx}(e^x) = e^x$

- $\frac{d}{dx}\ln(x) = \frac{1}{x}$

Summary of differentiation rules

Differentiation Rule	$f'(x)$ notation	$\frac{d}{dx}$ notation
Constant Rule	$f(x) = c \Rightarrow f'(x) = 0$	$\frac{d}{dx}[c] = 0$
Power Rule	$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$	$\frac{d}{dx}[x^n] = nx^{n-1}$
Sum Rule	$(g(x) + h(x))' = g'(x) + h'(x)$	$\frac{d}{dx}[g(x) + h(x)] = \frac{d}{dx}g(x) + \frac{d}{dx}h(x)$
Difference Rule	$(g(x) - h(x))' = g'(x) - h'(x)$	$\frac{d}{dx}[g(x) - h(x)] = \frac{d}{dx}g(x) - \frac{d}{dx}h(x)$
Constant Multiplier Rule	$(c \cdot g(x))' = c g'(x)$	$\frac{d}{dx}[c g(x)] = c \frac{d}{dx}g(x)$

And two special derivatives: $\frac{d}{dx}(e^x) = e^x$ and $\frac{d}{dx}\ln(x) = \frac{1}{x}$.

There's more differentiation rules (product, quotient, chain rule).

They're fun, but we won't be reviewing them.

Higher order derivatives

- We can continue to take derivatives of derivatives.
- The second derivative becomes the rate of change of the rate of change.
- If we interpret the first derivative as velocity, then the second derivative is acceleration.
- Notation:
 - Function: $f(x) = y$
 - 1st derivative: $f'(x)$ or $\frac{dy}{dx}$
 - 2nd derivative: $f''(x)$ or $\frac{d^2y}{dx^2}$
 - nth derivative: $f^{(n)}(x)$ or $\frac{d^ny}{dx^n}$

Derivatives practice

Solve individually and then check with a partner.

1. Which rules should you use to take these derivatives?

$$\text{A) } f(x) = 3x^4 \qquad \text{B) } y = 4x^2 + 3x - 16$$

2. Find the derivatives of these functions

$$\text{A) } y = 3x^2 \qquad \text{B) } h(x) = 2x(x^2 + 3x) \qquad \text{C) } g(y) = 2\sqrt{y} - e^y + 20 \ln(y)$$

3. Find the 3rd derivative of:

$$G(z) = 3z^4 - 8z^3 + 2z - 19$$

Derivatives practice

 Let's see a solution!

2.B) Find the derivative

$$h(x) = 2x(x^2 + 3x)$$

Derivatives practice

 Let's see a solution!

2.C) Find the derivative

$$g(y) = 2\sqrt{y} - e^y + 20 \ln(y)$$

Derivatives practice

 Let's see a solution!

3. Find the 3rd derivative of $G(z) = 3z^4 - 8z^3 + 2z - 19$.

Partial derivatives

Partial derivatives

When we find a partial derivative, we find an expression for the slope with respect to *one variable* in a multivariate function.

Notation: the partials of $f(x, y, z)$ are $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, and $\frac{\partial f}{\partial z}$

We read $\frac{\partial f}{\partial x}$ as “the partial derivative of f with respect to x ”

How to take partial derivatives? Find the derivative with respect to a single variable, *treating all others as constants*.

Partial derivatives example



Find all partial derivatives of

$$B(x, y, z) = 0.4x^3y - 3.6y^2 + 4zx + 8$$

Partial derivatives example

Find all partial derivatives of

$$B(x, y, z) = 0.4x^3y - 3.6y^2 + 4zx + 8$$

$$\frac{\partial B}{\partial x} = 1.2x^2y + 4z$$

$$\frac{\partial B}{\partial y} = 0.4x^3 - 7.2y$$

$$\frac{\partial B}{\partial z} = 4x$$

Partial derivatives example

Find all partial derivatives of

$$B(x, y, z) = 0.4x^3y - 3.6y^2 + 4zx + 8$$

$$\frac{\partial B}{\partial x} = 1.2x^2y + 4z$$

$$\frac{\partial B}{\partial y} = 0.4x^3 - 7.2y$$


$$\frac{\partial B}{\partial z} = 4x$$

 What is the value of the partial derivative of B with respect to z at the point $(1, 2, 3)$?

OK, but what do partials actually mean?

The slope with respect to one variable if other variables are held constant.

We have a function $f(x,y)$. Its graph is a surface in 3D :



Partial derivatives practice

a. The temperature (in Celsius) across a surface (where x and y are in meters) is described by:

$$T(x, y) = x^2y - 2x + y - 1$$

At what “rate” is temperature changing (with respect to distance):

- In the x direction, at the point $(1,3)$ on the surface?
- In the y direction, at the point $(0,2)$ on the surface?

b. A dragon’s breath temperature (T , in degrees Celsius) is modeled as a function of its wingspan (W , in meters) and length (L , also in meters):

$$T(W, L) = 0.41WL + 2.6W^2$$

- At what rate is breath temperature changing with respect to length for a dragon that is 4.1m long, with a wingspan of 4.5m?
- At what rate is breath temperature changing with respect to wingspan, for the same dragon?

Partial derivatives practice

 Let's see a solution.

- a. The temperature (in Celsius) across a surface (where x and y are in meters) is described by:

$$T(x, y) = x^2y - 2x + y - 1$$

At what “rate” is temperature changing (with respect to distance):

- In the x direction, at the point $(1,3)$ on the surface?
- In the y direction, at the point $(0,2)$ on the surface?

Partial derivatives practice

 Let's see a solution.

- b. A dragon's breath temperature (T , in degrees Celsius) is modeled as a function of its wingspan (W , in meters) and length (L , also in meters):

$$T(W, L) = 0.41WL + 2.6W^2$$

- At what rate is breath temperature changing with respect to length for a dragon that is 4.1m long, with a wingspan of 4.5m?
- At what rate is breath temperature changing with respect to wingspan, for the same dragon?

Example: higher order & partial derivatives in environmental data science

The **Advection-Dispersion-Reaction Equation** for solute transport models the change in a solute concentration C over time t and x, y, z space, where groundwater is flowing in direction x :

$$\frac{\partial C}{\partial t} = D_x \frac{\partial^2 C}{\partial x^2} + D_y \frac{\partial^2 C}{\partial y^2} + D_z \frac{\partial^2 C}{\partial z^2} - v \frac{\partial C}{\partial x} - \lambda RC$$

D_x, D_y, D_z are hydrodynamic diffusion coefficients in the x, y , and z direction.

Let's break it down.

$$\frac{\partial C}{\partial t} = D_x \frac{\partial^2 C}{\partial x^2} + D_y \frac{\partial^2 C}{\partial y^2} + D_z \frac{\partial^2 C}{\partial z^2} - v \frac{\partial C}{\partial x} - \lambda RC$$

- **Left-hand-side:** Rate of concentration change (over time)
- **Right-hand-side first 3 terms:** Concentration change due to dispersion in x , y , and z directions
- **Right-hand-side fourth term:** Concentration change due to groundwater transport (in groundwater flow direction, x)
- **Right-hand-side final term:** Reaction term (e.g. biodegradation / abiotic degradation)

What we covered today

- 🌱 The number e and $\exp()$
- 🐰 Logistic growth
- 📊 Natural logarithms
- 🔍 Limits
- ✍️ Derivatives and rate of change
- 🏔️ Tangent lines
- 📝 Rules for differentiation
- 💠 Constant rule
- 💪 Power rule
- + Sum and difference rule
- ✖️ Product rule
- 📈 Higher order derivatives
- 🌐 Partial derivatives