

# Homework 1-Corrections-Math 5733

Evan Shapiro

Master's of Integrated Science, University of Colorado Denver

February 19, 2017

1.4

We start with the the following Cauchy differential equation:

$$v'(t) = -(\alpha + \epsilon)v(t)$$

$$v(0) = v_0,$$

where  $\alpha > 0$  is a measured parameter of the differential equation, and  $\epsilon$  is a perturbation in the measurement.

a) For a solution at time  $t = 1$ , do small perturbations in  $\alpha$  imply small changes in the solution,  $u(t)$ ? In other words, is the solution at time  $t=1$  stable for small perturbations in  $\alpha$ ?

First, we solve the Cauchy problem:

$$\frac{dv}{dt} = -(\alpha + \epsilon)v(t)$$

$$\frac{dv}{v} = -(\alpha + \epsilon)dt$$

$$\ln(v) = -(\alpha + \epsilon)t + C$$

$$v(t) = e^{-(\alpha + \epsilon)t + C}$$

$$v(t) = v_0 e^{-(\alpha + \epsilon)t}$$

Expand this function with a Taylor series around the parameter value  $\alpha$  for small values of  $\epsilon$ , at time  $t = 1$ .

$$v(-(\alpha + \epsilon)) = u_0(e^{-\alpha} - \epsilon * e^{-c}),$$

where  $\epsilon * e^{-c}$  is the truncation error, and  $c \in (\alpha, \alpha + \epsilon)$ , such that it maximizes the error term. We can use this Taylor expansion to bound the error of the perturbed function about the point  $t = 1$ . If we define the error to be bounded by some value  $\eta$ , the difference between the exact and perturbed solution is:

$$|e^{-(\alpha)t} - e^{-(\alpha + \epsilon)t}| \leq |\epsilon * e^{-c}|,$$

which is of order  $\epsilon$ , and is thus finitely bounded. Limiting the bound we see that

$$\lim_{\epsilon \rightarrow 0} |e^{-(\alpha)t} - e^{-(\alpha+\epsilon)t}| \leq \lim_{\epsilon \rightarrow 0} |\epsilon * e^{-c}| = 0$$

This means that a perturbation in the parameter  $\alpha$  of  $O(\epsilon)$  lead to perturbations in the data of  $O(\epsilon)$  making this a stable system.

b) Next we assume that both  $u_0$  and  $\alpha$  are measured. Discuss the stability of the problem in this context. The solution for the Cauchy problem with both perturbed  $\alpha$  and  $\epsilon$  is:

$$v(t) = (u_0 + \epsilon_0)e^{-(\alpha+\epsilon_1)t}.$$

This has a linear dependence on  $u_0$  and an exponential dependence on  $\alpha$ , so it is stable with respect to small perturbations of  $u_0$ , regardless of the size of  $u_0$ , and is stable with respect to small perturbation of  $\alpha$

1.5(c) Solve the following Cauchy problem

$$\begin{aligned} u_t + xu_x &= x & x \in \mathbb{R}, t > 0 \\ u(x, 0) &= \cos(90x). \end{aligned}$$

This is an inhomogeneous Cauchy ODE. So first we solve for homogeneous component,  $a(x,t)$ , and then solve the inhomogeneous component. In the previous problem, part (b), we saw that this type of initial condition yields the solution.

$$u_h = \cos(90xe^{-t}) \quad x_0 = xe^{-t}$$

The solution to the characteristic equation is:

$$u_c = \int_0^t x_0 e^\tau d\tau$$

$$u_c = x_0(e^t - 1)$$

Substituting in  $x_0$ :

$$u_c = x(1 - e^{-t})$$

The general solution is thus:

$$u(x, t) = \cos(90xe^{-t}) + x(1 - e^{-t})$$

Checking that this is the solution. The initial condition returns:

$$u(x, 0) = \cos(90x)$$

While substituting the general solution into the ODE yields:

$$u_t = 90xe^{-t} \sin(90xe^{-t}) + e^{-t}$$

$$xu_x = -90xe^{-t}\sin(90xe^{-t}) + x(1 - e^{-t})$$

$$u_t + xu_x = x$$

Thus, the solution is verified.

1.14

Consider the following Cauchy problem:

$$u_{tt} = c^2 u_{xx}$$

$$u(x, 0) = \phi(x)$$

$$u_t(x, 0) = \psi(x)$$

Let:

$$v = u_t + cu_x.$$

Part (a)

Show that

$$v_t - cv_x = 0$$

Solution:

Given that  $v = u_t + cu_x$ ,

$$v_t = u_{tt} + cu_{xt}, \quad v_x = u_{tx} + cu_{xx}.$$

This problem has  $u_{tt} = c^2 u_{xx}$ . Regarding the cross terms, according to Clairut if a function  $u$  is defined on an open set  $D \in \mathbb{R}^2$ , and both  $u_{xy}, u_{yx}$  are continuous throught  $D$ , then  $u_{xy} = u_{yx}$ . Thus

$$v_t - cv_x = c^2 u_{xx} + cu_{xt} - cu_{xt} - c^2 u_{xx} = 0$$

Part (b):

Find  $v(x, t)$  expressed by  $\phi$  and  $\psi$ .

Solution:

First recognize that:

$$\frac{\partial v(x, t)}{\partial t} = v_t - cv_x,$$

which motivates us to use the method of characteristics to solve for  $v(x, t)$  :

$$v_t - cv_x = 0$$

$$\frac{dx}{dt} = -c$$

$$x = x_0 - ct$$

$$\begin{aligned}
x_0 &= x + ct \\
v(x, 0) &= v(x_0, 0) = u_t(x, 0) + cu_x(x, 0) \\
v(x, t) &= c\phi'(x + ct) + \psi(x + ct)
\end{aligned}$$

Part (c)

Explain why

$$u(x, t) = \phi(x - ct) + \int_0^t v[x - c(t - \tau), \tau] d\tau$$

Solution:

Use the solution for  $x_0$  from the method of characteristics to solve the inhomogeneous component of this problem.

$$\begin{aligned}
\frac{dx}{dt} &= c \\
x &= x_0 + ct \\
x_0 &= x - ct \\
u(x, 0) &= u(x, t) = \phi(x - ct) + \int_0^t v(x_0 + c\tau, \tau) d\tau
\end{aligned}$$

Substituting  $x_0$  into the integral

$$\begin{aligned}
u(x, t) &= \phi(x - ct) + \int_0^t v(x - ct + c\tau, \tau) d\tau \\
u(x, t) &= \phi(x - ct) + \int_0^t v(x - c(t - \tau), \tau) d\tau
\end{aligned}$$

Part (d)

Derive the expression

$$u(x, t) = \frac{1}{2}(\phi(x + ct) \pm \phi(x - ct)) + \frac{1}{2c} \int_{x-ct}^{x+ct} \psi(\theta) d\theta$$

Solution:

Substitute the solution from part b

$$v(x, t) = c\phi'(x + ct) + \psi(x + ct)$$

into the solution for part c

$$u(x, t) = \phi(x - ct) + \int_0^t v(x - c(t - \tau), \tau) d\tau.$$

This becomes:

$$u(x, t) = \phi(x - ct) + \int_0^t c\phi'(x - c(t - \tau) + c\tau) d\tau + \int_0^t \psi(x - c(t - \tau) + \tau) d\tau,$$

which simplifies to

$$u(x, t) = \phi(x - ct) + \int_0^t c\phi'(x - ct + 2c\tau)d\tau + \int_0^t \psi(x - ct + 2c\tau)d\tau.$$

Lets solve the first integral

$$\int_0^t c\phi'(x - ct + 2c\tau)d\tau$$

by substituting in

$$u(\tau) = x - ct + 2c\tau, \quad du(\tau) = 2cd\tau.$$

The new limits of integration are:

$$u(0) = x - ct, \quad u(t) = x + ct$$

The new integral is:

$$\frac{1}{2} \int_{x-ct}^{x+ct} \phi'(u)du$$

In part b we took the derivative of  $\phi$  with respect to  $x$  to yield  $\phi'$ . Using calculus we see

$$\frac{d\phi}{dx} = \frac{du}{dx} \frac{d\phi}{du}$$

with

$$\frac{du}{dx} = 1$$

So the integral becomes:

$$\frac{1}{2} \int_{x-ct}^{x+ct} d\phi(u),$$

which according to the Fundamental Theorem of Calculus yields

$$\int_0^t c\phi'(x - ct + 2c\tau)d\tau = \frac{1}{2}(\phi(x + ct) - \phi(x - ct)).$$

Using the same  $u$ -substitution in the second integral as we did for the first yields the following solution for  $u(x, t)$ :

$$\int_0^t \psi(x - ct + 2c\tau)d\tau = \frac{1}{2c} \int_{x-ct}^{x+ct} \psi(u)du$$