

# Pesticides in small agricultural streams in Germany

Eduard Szöcs,<sup>\*,†</sup> Marvin Brinke,<sup>‡</sup> Bilgin Karaoglan,<sup>¶</sup> and Ralf B. Schäfer<sup>†</sup>

*Institute for Environmental Sciences, University of Koblenz-Landau, Germany, German Federal Institute of Hydrology (BfG), Koblenz, Germany, and Federal Environmental Agency (UBA), Dessau-Roßlau, Germany*

E-mail: szoechs@uni-landau.de

## Abstract

Fehlt noch...

## Introduction

More than 50% of the total land area in Germany are used by agriculture<sup>1</sup>. In the year 2014 more than 45,000 tonnes of 766 authorized pesticides were sold for application on this area<sup>2</sup>. The applied pesticides may enter surface waters via spray-drift, edge-of-field run-off or drainage, with run-off being one of the major input routes<sup>3,4</sup>. Once entered the surface waters pesticides are frequently detected in environmental monitoring<sup>5</sup> and may have adverse effects on biota and ecosystem functioning<sup>6,7</sup>.

---

<sup>\*</sup>To whom correspondence should be addressed

<sup>†</sup>Institute for Environmental Sciences

<sup>‡</sup>German Federal Institute of Hydrology

<sup>¶</sup>German Federal Environmental Agency

Malaj et al.<sup>5</sup> analyzed data supplied to the European Union (EU) in the context of the Water Framework Directive (WFD) and showed that pesticides jeopardize the health of freshwater ecosystems. However, this study reflected only a small part of data that is available from national monitoring programs. These programs are setup for determination and surveillance of the chemical and ecological status of surface, ground and drinking water. Large amounts of data are generated, which possibly can also be used to answer other questions. In Germany monitoring programs are setup independently by the federal states in compliance with the WFD<sup>8</sup> and additional state specific needs. However, currently there is no curated national-wide compilation of this data available.

Stehle and Schulz<sup>9</sup> compiled 1566 measured concentrations of 23 insecticides in the EU from scientific publications. They found that insecticides that many of these measurements exceed regulatory acceptable concentrations (RAC), especially in very small catchments  $<1\ km^2$ . Small water bodies are important refuges of biodiversity<sup>10</sup> and enable downstream colonisation of polluted streams<sup>11</sup>. At the same time they may be exposed to a high risk of pesticide contamination from adjacent agricultural areas and low dilution effects<sup>4</sup>. Although small streams comprise a major fraction of streams<sup>12</sup>, relatively little is known about their chemical and ecological status.

The aim of this study was to compile all available chemical monitoring data in Germany and to answer the questions:

- (i) Can the currently available monitoring data used for a representative description of the pollution situation?
- (ii) Are small agricultural waters more polluted compared to bigger streams? Are there thresholds in these relationships?
- (iii) How polluted are small streams and which pesticides are most important?

# Methods

## Data compilation

We queried chemical monitoring data of pesticides from sampling sites with catchment sizes < 100km<sup>2</sup> for the years 2005 to 2015 from all 13 non-city federal states of Germany. Additionally, we compiled data available from previous studies and searched online databases. This yielded to a total of more than 30 datasets of different formats. In the following we will use the ISO 3166-2:DE standard abbreviations for federal states.

We homogenized and unified these datasets into a common database. We implemented a robust and reproducible data cleaning work flow<sup>13</sup>, though parts of the dataset are proprietary. An overview of the data cleaning process is provided in the supplemental materials. To assess whether samples were taken during potential rainfall events we performed spatio-temporal intersection of sampling events with daily precipitation data from the sampling date and the day before.<sup>14</sup>.

## Characterization of chemical pollution

We characterized chemical pollution (excluding sum parameters) using three indicators:

1. National and international Environmental Quality Standards (EQS)<sup>15,16</sup>: We used only Maximum Annual Concentration EQS (MAC-EQS) for characterization. These were available for 29 compounds (Supplement, Table xxx). ref
2. Regulatory Acceptable Concentrations (RAC)<sup>17</sup>: This is the lowest concentration at which no acceptable biological effects are expected. These are derived during authorization process of pesticides and incorporate also an uncertainty factor. The German Federal Environmental Agency provided RACs for 105 compounds (Supplement, Table xxx). We expressed RAC as Risk Quotient (RQ); ref

$$RQ_i = \frac{C_i}{RAC_i} \quad (1)$$

Where  $C_i$  is the concentration of a compound  $i$  in a sample.

3. Maximum Toxic Units ( $TU_{max}$ )<sup>18</sup>:

$$TU_{max} = \max\left(\frac{C_i}{EC_{50,D.magna,i}}\right) \quad (2)$$

Where  $C_i$  is the concentration of compound  $i$  in a sample and  $EC_{50,D.magna,i}$  is the concentration of this compound where 50% of the exposed animals showed after 48 hours an effect in a laboratory study. We compiled  $EC_{50,D.magna}$  values from literature<sup>5</sup>, databases<sup>19,20</sup> or model predictions<sup>21</sup>, where experimental data had priority. We could compile  $EC_{50,D.magna}$  values for 394 compounds (Supplement, Table xxx)). We used the maximum TU per sample, as it is independent of the number of measured compounds and makes no assumptions on the mode of action. Additionally, we also calculated the sum of toxic units ( $TU_{sum}$ ). A table of all included compounds can be found in the supplement.

## Characterization of catchments

We delineated catchments upstream of the sampling sites using a digital elevation model<sup>22</sup> and a multiple flow direction algorithm<sup>23</sup> as implemented in GRASS GIS 7<sup>24</sup>. Catchment delineation has been manually checked for accuracy. In areas with low relief energy the delineation algorithm did not produce accurate results and we used river catchments provided by federal state authorities in these cases. For each catchment we calculated the relative coverage (%) with agricultural areas based on Official Topographical Cartographic Information System (ATKIS) of the land survey authorities.

## Statistical analyses

All data-processing and analyses have been performed using R<sup>25</sup>. To display differences in the spectra of analyzed compounds between federal states we used Multidimensional Scaling (MDS) based on Jaccard dissimilarity in conjunction with hierarchical clustering using the vegan package<sup>26</sup>. We expected non-linear responses to agriculture and catchment size and therefore, used generalized additive models (GAM) to identify relationships<sup>27</sup>.

We modeled the number of RAC exceedances as:


$$\begin{aligned} No_i &\sim NB(\mu_i, k) \\ E(No_i) &= \mu_i \text{ and } Var(No_i) = \mu_i + \frac{\mu_i^2}{k} \\ log(\mu_i) &= \beta_0 + f_1(Agri_i) + f_2(Size_i) + log(n_i) \end{aligned} \quad (3)$$

where  $No_i$  is the observed number of exceedances at site  $i$ ,  $Agri_i$  the proportion of agriculture within the catchment and  $Size_i$  the catchment size of the site. We modeled  $No_i$  as resulting from a negative binomial distribution ( $NB$ ) and used the number of sampling events per site ( $n_i$ ) as an offset to account different sampling efforts.  $f_1$  and  $f_2$  are smoothing functions using thin plate regression splines<sup>28</sup>. The degree of smoothness was estimated using restricted maximum likelihood (REML) during model fitting process<sup>29</sup>. Similar models were fitted to the number of EQS-exceedances and the 95th percentile of  $TU_{max}$  (see Supplement for details). We used point-wise 95% Confidence Intervals of the first derivative of the fitted smooth to check if there are regions of statistically significant changes. GAMs were fitted using the mgcv package<sup>29</sup>.

# Results

## Overview and representativeness of compiled data

The compiled dataset comprised only few standing waters (58 sites) and the majority (90%) of samples were taken via grab sampling. Therefore, we report only results of grab samples from streams. The analyzed dataset comprised 2,918,604 measurements from 42,236 samples from 3,049 sampling sites. We found big differences in the number of sampling sites between federal states (Figure 1 and Supplement, Table ).



Set  
ref to  
table

Figure 1: Spatial distribution of the 3109 sampling sites. Colour codes different federal states.

In total 484 different compounds used as pesticides and their metabolites were measured at least once (Supplement, Table ). Most of the compounds were herbicides (179), followed by insecticides (117) and fungicides (109). We found substantial differences of the spectra of analyzed compounds between federal states (Figure 2). Hierarchical clustering revealed three groups: i) with less than 100 compounds (SL, ST and TH), ii) with a medium sized spectra and

Set  
ref to  
table

iii) with a big and distinct spectra (RP and NI).

Only 5.5% (160,800) of all measurements were detected above the limit of quantification (LOQ). The spatio-temporal intersection revealed that 5% of the samples were taken at or after days with rainfall events greater than 10mm / day. .




Figure 2: Compound spectra of the different federal states. Left: Barcode plot - Each vertical line is an analysed compound. Right: MDS ordination. Colors according to three groups determined by hierarchical clustering (see Supplement Figure xxx).

We were able to derive for 2376 sites catchment sizes and the proportion of agriculture within catchments. The distribution of sampling sites across catchment area and agricultural area in the catchment revealed a sharp decline in the distribution of catchment-sizes below  $10 \text{ km}^2$ , with most sampling sites with catchments between  $10$  and  $25 \text{ km}^2$  (Figure 3). The proportion of agriculture in the catchments decreased with increasing catchment size.

## Are small agricultural waters more polluted compared to bigger streams?

Modeling the number of RAC exceedances as function of agriculture within catchment and catchment size revealed that there is a strong and statistically significant increase up to 25%




Figure 3: Distribution of catchment area and agriculture within the catchment area across the sampling sites. Only sampling sites with catchment area  $< 150 \text{ km}^2$  are displayed. Colour codes the 2-dimensional density of points.

agriculture. Above this threshold the exceedances level off followed by a increase above 75% (Figure 4, left).

We could no detect any effect of catchment size on the number of RAC exceedances (Figure 4, right) and no interaction between these two predictors.

The number of EQS-exceedances and the 95th percentile of  $TU_{max}$  showed similar patterns and thresholds (see Supplement Figure Sxxx - Sxxx). references

## Pollution of small agricultural streams

Based on the results previous results and given the low amount of sample sites with catchment sizes below  $10 \text{ km}^2$  we explored the pollution of small agricultural streams, defined as streams with catchment size less than  $30 \text{ km}^2$  and more than 25% agriculture within the catchment




Figure 4: Effect of agriculture within the catchment (left) and catchment size (right) on the number of RAC exceedances. Red line marks statistically significant changes. Dashed lines denote 95% pointwise Confidence Intervals.

( $n = 1030$  sites with 10591 samples and 36550 measurements). Out of the 29 compounds with EQS, the most frequent EQS exceedances were recorded for the herbicides Nicosulfuron (2.1%,  $n = 1769$ ), Flufenacet (1.1%,  $n = 6301$ ) and Isoproturon (0.7%,  $n = 8380$ ). Other compounds show exceedances only at less than 0.5% of all samples (Figure 5A). Neonicotinoid Insecticides and Chlropyrifos showed highest risk quotients. For Thiacloprid, Imidacloprid and Chlropyrifos RAC was less than LOQ, therefore, all detections have a  $RQ > 1$  (Figure 5B). In 15% of all samples ( $n = 6377$ ) risk quotients higher than 1 were observed. Highest  $RQ$  were observed for Chlropyrifos (244), Dimoxystrobin(117) and Isoproturon (80). In 33% of the samples no pesticides were detected. The mean  $\log(TU_{max})$  for detects was -4.6. 2.6% of the samples showed  $\log(TU_{max})$  values greater than -2 (Figure 5C). We found a high correlation of  $\log(TU_{max})$  and  $TU_{sum}$ , with a maximum higher value of  $TU_{sum}$  xx units. In most samples (55.5%) more than one compound was detected, with a maximum of 54

ausrechnen  
und r  
angeben

different compounds (Figure 5D).




Figure 5: Overview on pesticide pollution of small agricultural streams. **A:** 10 compounds with most frequent EQS exceedances. See B for color legend. **B:** Distribution of  $TU_{\max}$  values. Values with  $TU_{\max} = 0$  are not displayed (3501 out of 10591 samples). **C:** 15 compounds with highest risk quotients. **D:** Distribution of the number of detected compounds in a sample.

# Discussion

## Representativeness of the dataset

The compiled dataset of governmental monitoring data represents currently the most comprehensive one available for Germany. Similar nationwide datasets have been compiled for the Netherlands<sup>30</sup>, Switzerland<sup>31</sup> and the United States (Water Quality Portal (WQP) [www.waterqualitydata.us](http://www.waterqualitydata.us)).

Nevertheless such nationwide compilations, may not only be useful for governmental surveillance, but also to answer other questions, like validation of exposure modeling,<sup>32</sup> retrospective evaluation of regulatory risk assessment<sup>9,33</sup> or investigation of pesticide mixtures.<sup>34</sup>

A nationwide assessment is hampered by the inhomogeneity of monitoring data between federal states: There are not only big differences in the spatial distribution and quantity of sampling sites (Figure 1), but also the spectrum of analyzed compounds (Figure 2) and differences in the quality of chemical analyses. Future monitoring programs should aim at resolving these differences in order to enable a nationwide assessment.

## Influence of catchment area and agriculture

We found a strong influence of agriculture on the pollution of streams. If there is more than 25% agriculture within a catchment pesticides can be detected in streams and may have biological effects .

We did not find support for our hypothesis that small streams are more polluted than big streams, though previous studies have shown such a relationship<sup>6,9</sup>. This could be explained by the relatively small gradient of catchment sizes in our dataset, with most of the streams being  $<100 \text{ km}^2$  (Figure 3, top). For example the gradient of Schulz<sup>6</sup> covered 6 orders of magnitude. Another reason might be the unequal distribution of catchment sizes, with less sites  $<10 \text{ km}^2$  and  $>100 \text{ km}^2$  (Figure 3, top).

compare  
num-  
bers  
of dif-  
ferent  
at-  
lases,  
num-  
ber of  
sites /  
com-  
pounds

citation  
needed

## Pollution of streams

Stehle and Schulz<sup>9</sup> found the highest percentage of RAC exceedances organophosphate insecticides. Our results revealed that neonicotinoid insecticides show high exceedances, followed by the organophosphate chlorpyrifos. This difference can be attributed to the low sample size for neonicotinoid insecticides in their study ( $n = 33$ ) compared to the dataset presented here and shows that this class may pose a high risk to freshwaters . Our results show lower proportions of exceedances compared to Stehle and Schulz<sup>9</sup>, which can be attributed to different aims of the data sources: scientific research aims at finding pollutants, whereas monitoring aims mainly at surveillance of water quality. This is also reflected in the different sample sizes (1,566 vs. 2,918,604 measurements ) and the high number of non-detects in the monitoring data.

Contrary, Knauer<sup>33</sup> found exceedances from monitoring data mainly for herbicides and fungicides and only one insecticide Chlorpyrifos. This might reflect differences in use between countries and RACs.

xxxxsamples showed  $TU_{max}$  values greater than -3, a threshold above biological effects have been shown . The high correlation of TU and the relatively low exceedance of  $TU_{sum}$  reveals that in compound mixtures occurring in the field there is a skewed distribution of toxicities with one dominant compound. Nevertheless, most pesticides did not occur individually but in mixtures. This highlights that

Exceedances of RACs are undesirable ...

## Acknowledgement

The authors thank the authorities for providing chemical monitoring data and the German Federal Environmental Protection Agency (UBA) for funding this project.

zahlen  
neon-  
ics  
anderes  
Wort  
use  
num-  
ber for  
sub-  
stances  
with  
RAC  
add  
num-  
ber  
add  
cita-  
tion  
schae-  
fer

## **Supporting Information Available**

The following files are available free of charge.

- Supplemental \_ Materials.pdf : Supplemental Materials (Figures, Tables, Models).

This material is available free of charge via the Internet at <http://pubs.acs.org/>.

## **References**

- (1) Statistisches Bundesamt, *Bodenfläche nach Art der tatsächlichen Nutzung*; Fachserie 3 Reihe 5.1; 2014.
- (2) Bundesamt für Verbraucherschutz und Lebensmittelsicherheit, *Absatz an Pflanzenschutzmitteln in der Bundesrepublik Deutschland - Ergebnisse der Meldungen gemäß § 64 Pflanzenschutzgesetz für das Jahr 2014*; 2015.
- (3) Schulz, R. Comparison of spray drift-and runoff-related input of azinphos-methyl and endosulfan from fruit orchards into the Lourens River, South Africa. *Chemosphere* **2001**, *45*, 543–551.
- (4) Liess, M.; Schulz, R.; Liess, M.-D.; Rother, B.; Kreuzig, R. Determination of insecticide contamination in agricultural headwater streams. *Water Research* **1999**, *33*, 239–247.
- (5) Malaj, E.; Ohe, P. C. v. d.; Grote, M.; Kühne, R.; Mondy, C. P.; Usseglio-Polatera, P.; Brack, W.; Schäfer, R. B. Organic chemicals jeopardize the health of freshwater ecosystems on the continental scale. *Proceedings of the National Academy of Sciences* **2014**, *111*, 9549–9554.
- (6) Schulz, R. Field Studies on Exposure, Effects, and Risk Mitigation of Aquatic Nonpoint-Source Insecticide Pollution: A Review. *Journal of Environmental Quality* **2004**, *33*, 419–448.

- (7) Schäfer, R. B.; Caquet, T.; Siimes, K.; Mueller, R.; Lagadic, L.; Liess, M. Effects of pesticides on community structure and ecosystem functions in agricultural streams of three biogeographical regions in Europe. *Science of the Total Environment* **2007**, *382*, 272–285.
- (8) Quevauviller, P.; Borchers, U.; Thompson, C.; Simonart, T. *The Water Framework Directive: Ecological and Chemical Status Monitoring*; John Wiley & Sons, 2008.
- (9) Stehle, S.; Schulz, R. Pesticide authorization in the EU—environment unprotected? *Environmental Science and Pollution Research* **2015**,
- (10) Davies, B. R.; Biggs, J.; Williams, P. J.; Lee, J. T.; Thompson, S. A comparison of the catchment sizes of rivers, streams, ponds, ditches and lakes: implications for protecting aquatic biodiversity in an agricultural landscape. *Hydrobiologia* **2008**, *597*, 7–17.
- (11) Liess, M.; von der Ohe, P. C. Analyzing effects of pesticides on invertebrate communities in streams. *Environmental Toxicology and Chemistry* **2005**, *24*, 954–965.
- (12) Nadeau, T.-L.; Rains, M. C. Hydrological Connectivity Between Headwater Streams and Downstream Waters: How Science Can Inform Policy1: Hydrological Connectivity Between Headwater Streams and Downstream Waters: How Science Can Inform Policy. *JAWRA Journal of the American Water Resources Association* **2007**, *43*, 118–133.
- (13) Poisot, T. Best publishing practices to improve user confidence in scientific software. *Ideas in Ecology and Evolution* **2015**, *8*.
- (14) Rauthe, M.; Steiner, H.; Riediger, U.; Mazurkiewicz, A.; Gratzki, A. A Central European precipitation climatology – Part I: Generation and validation of a high-resolution gridded daily data set (HYRAS). *Meteorologische Zeitschrift* **2013**, *22*, 235–256.
- (15) OGewV, Verordnung zum Schutz der Oberflächengewässer (Oberflächengewässerverordnung). 2011; BGBl. I S. 1429.

- (16) European Union, DIRECTIVE 2013/39/EU OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. 2013.
- (17) Brock, T. C. M., Alix, A., Brown, C. D., Capri, E., Gottesbüren, B. E., Heimbach, F., Lythgo, C. M., Schulz, R., Strelöke, M., Eds. *Linking aquatic exposure and effects: risk assessment of pesticides: EU and SETAC Europe workshop ELINK, Bari, Italy, and Wageningen, Netherlands*; CRC Press: Boca Raton, 2010.
- (18) Sprague, J. Measurement of pollutant toxicity to fish, II-Utilizing and applying bioassay results. *Water Research* **1970**, *4*, 3–32.
- (19) Lewis, K. A.; Tzilivakis, J.; Warner, D. J.; Green, A. An international database for pesticide risk assessments and management. *Human and Ecological Risk Assessment: An International Journal* **2016**, *22*, 1050–1064.
- (20) U.S. EPA, The ECOTOXicology knowledgebase (ECOTOX). 2015; <http://cfpub.epa.gov/ecotox/>.
- (21) Schüürmann, G.; Ebert, R.-U.; Kühne, R. Quantitative read-across for predicting the acute fish toxicity of organic compounds. *Environmental Science & Technology* **2011**, *45*, 4616–4622.
- (22) EEA, Digital Elevation Model over Europe (EU-DEM). 2013; <http://www.eea.europa.eu/data-and-maps/data/eu-dem#tab-metadata>.
- (23) Holmgren, P. Multiple flow direction algorithms for runoff modelling in grid based elevation models: An empirical evaluation. *Hydrological Processes* **1994**, *8*, 327–334.
- (24) Neteler, M.; Bowman, M. H.; Landa, M.; Metz, M. GRASS GIS: A multi-purpose open source GIS. *Environmental Modelling & Software* **2012**, *31*, 124–130.

- (25) R Core Team, *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2016.
- (26) Oksanen, J.; Blanchet, F. G.; Kindt, R.; Legendre, P.; Minchin, P. R.; O'Hara, R. B.; Simpson, G. L.; Solymos, P.; Stevens, M. H. H.; Wagner, H. *vegan: Community Ecology Package*; 2016; R package version 2.3-5.
- (27) Fewster, R. M.; Buckland, S. T.; Siriwardena, G. M.; Baillie, S. R.; Wilson, J. D. Analysis of population trends for farmland birds using generalized additive models. *Ecology* **2000**, *81*, 1970–1984.
- (28) Wood, S. N. Thin Plate Regression Splines. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)* **2003**, *65*, 95–114.
- (29) Wood, S. N. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **2011**, *73*, 3–36.
- (30) Vijver, M. G.; Van 'T Zelfde, M.; Tamis, W. L.; Musters, K. J.; De Snoo, G. R. Spatial and temporal analysis of pesticides concentrations in surface water: Pesticides atlas. *Journal of Environmental Science and Health, Part B* **2008**, *43*, 665–674.
- (31) Munz, N.; Leu, C. Pestizidmessungen in Fliessgewässern - schweizweite Auswertung. *Aqua & Gas* **2011**, *11*, 32–41.
- (32) Knäbel, A.; Meyer, K.; Rapp, J.; Schulz, R. Fungicide Field Concentrations Exceed FOCUS Surface Water Predictions: Urgent Need of Model Improvement. *Environmental Science & Technology* **2014**, *48*, 455–463.
- (33) Knauer, K. Pesticides in surface waters: a comparison with regulatory acceptable concentrations (RACs) determined in the authorization process and consideration for regulation. *Environmental Sciences Europe* **2016**, *28*.

- (34) Schreiner, V. C.; Szöcs, E.; Bhowmik, A. K.; Vijver, M. G.; Schäfer, R. B. Pesticide mixtures in streams of several European countries and the USA. *Science Of The Total Environment* **2016**, under Review.

# Graphical TOC Entry

