

# Pesticides pollution of small streams in Germany

Eduard Szöcs,<sup>\*,†</sup> Marvin Brinke,<sup>‡</sup> Bilgin Karaoglan,<sup>¶</sup> and Ralf B. Schäfer<sup>†</sup>

*Institute for Environmental Sciences, University of Koblenz-Landau, Germany, German  
Federal Institute of Hydrology (BfG), Koblenz, Germany, and Federal Environmental  
Agency (UBA), Dessau-Roßlau, Germany*

E-mail: szoecs@uni-landau.de

## Abstract

Fehlt noch...

## Introduction

More than 50% of the total land area in Germany are used by agriculture<sup>1</sup>. In the year 2014 more the 45,000 tonnes of 766 authorized pesticides were sold for application on this area<sup>2</sup>. The applied pesticides may enter surface waters via spray-drift, edge-off-field run-off or drainage, with run-off being one of the major input routes<sup>3,4</sup>. Once entered the surface waters pesticides are frequently detected in environmental monitoring<sup>5</sup> and may have adverse effects on biota and ecosystem functioning<sup>6,7</sup>.

---

<sup>\*</sup>To whom correspondence should be addressed

<sup>†</sup>Institute for Environmental Sciences

<sup>‡</sup>German Federal Institute of Hydrology

<sup>¶</sup>German Federal Environmental Agency

National monitoring programs are setup for determination and surveillance of the chemical and ecological status of surface, ground and drinking water. These monitoring programs produce huge amounts of data, which possibly can also be used to answer other questions. In Germany monitoring programs are setup by the federal states in compliance with the Water Framework Directive<sup>8</sup> and further state specific needs. However, currently there is no curated national-wide compilation of this data.

Small water bodies are important refuges of biodiversity enabling downstream colonisation. At the same time they may be exposed to a high risk of pesticide contamination from adjacent agricultural areas and low dilution effects<sup>4</sup>.

Although small waterbodies make up most  
running water with at least 10km<sup>2</sup> catchment area

However, very little is known about the pollution of such streams. The

The aim of this study was to compile monitoring data on a national scale and to answer the questions:

- (i) Is the currently available monitoring data a representative description of the pollution situation?
- (ii) Are small agricultural waters more polluted compared to bigger streams? Are there thresholds?
- (iii) How polluted are small streams and which pesticides are responsible?

## Methods

### Data compilation

We queried chemical monitoring data of pesticides from sampling sites with catchment size  $< 100\text{km}^2$  for the years 2005 to 2015 from all 13 non-city federal states of Germany. Additionally, we compiled data available from previous studies and searched online databases. This yielded to a total of more than 30 datasets of different formats.

We homogenized and unified these datasets into a common database. We implemented a robust and transparent data cleaning work flow<sup>9</sup>, though parts of the dataset are proprietary. An overview of the data cleaning process is provided in the supplemental materials. To assess whether samples were taken during potential rainfall events we intersected sampling coordinates with daily precipitation data<sup>10</sup> from the sampling date and the day before.

## Characterization of chemical pollution

We characterized chemical pollution using three indicators:

1. National and international Environmental Quality Standards (EQS)<sup>11,12</sup>: We used only Maximum Annual Concentration EQS (MAC-EQS) for characterization.
2. Regulatory Acceptable Concentrations (RAC)<sup>13</sup>: This is the lowest concentration at which no acceptable biological effects are expected. These are derived during authorization process of pesticides and contain an uncertainty factor. The German Federal Environmental Agency provided RACs for this study. We expressed RAC as Risk Quotient (RQ)

$$RQ = \frac{C}{RAC} \tag{1}$$

Where  $C$  is the concentration of a compound in a sample.

3. Maximum Toxic Units ( $TU_{max}$ )<sup>14</sup>:

$$TU_{max} = \max\left(\frac{C_i}{EC_{50,D.magna,i}}\right) \tag{2}$$

Where  $C_i$  is the concentration of compound  $i$  in a sample and  $EC_{50,D.magna,i}$  is the concentration of this compound where 50% of the exposed animals showed after 48

hours an effect in a laboratory study. We compiled  $EC_{50,D.magna}$  values from literature<sup>5</sup>, databases<sup>15,16</sup> or model predictions<sup>17</sup>, where experimental data had priority. We used the maximum TU per sample, as it is independent of the number of measured compounds and makes no assumptions on the mode of action. A table of all included compounds can be found in the supplement.

## Characterization of catchments

We delineated catchments upstream of the sampling sites using a digital elevation model<sup>18</sup> and a multiple flow direction algorithm<sup>19</sup> as implemented in GRASS GIS 7<sup>20</sup>. Catchment delineation has been manually checked for accuracy. In areas with low relief energy the delineation algorithm did not produce accurate results and we used river catchments provide by federal state authorities in these cases. For each catchment we calculated the relative coverage (%) with agricultural areas based on Official Topographical Cartographic Information System (ATKIS) of the land survey authorities.

## Statistical analyses

We used Multidimensional Scaling (MDS) based on jaccard dissimilarity in conjunction with hierarchical clustering to display differences in the spectra of analysed compounds per federal state.

# Results

## Overview and representativeness of compiled data

The compiled dataset comprised only a small fraction of standing waters and most of the samples were sampled via grab sampling. Therefore, we report only results for grab sam- pling from streams.

zahlen

The analysed dataset comprised 42236 samples from 3049 sampling sites (Figure 1 and Supplement). We found big differences in the number of sampling sites between federal states.

The dataset include 484




Figure 1: Spatial distribution of the 3109 sampling sites. Colour codes different federal states.




Figure 2: Distribution of catchment area and agriculture within the catchment area across the sampling sites. Only sampling sites with catchment area  $< 150 \text{ km}^2$  are displayed. Colour codes the 2-dimensional density of points.

Are small agricultural waters more polluted compared to bigger streams?

Pesticide pollution of small streams

## Discussion

Vergleich mit der Schweiz.....

## Subsection

## Acknowledgement

The authors thank the authorities for providing chemical monitoring data and the German Federal Environmental Protection Agency (UBA) for funding this project.

## Supporting Information Available

The following files are available free of charge.

- Supplemental\_Materials.pdf : Supplemental Materials

This material is available free of charge via the Internet at <http://pubs.acs.org/>.

## References

- (1) Statistisches Bundesamt, *Bodenfläche nach Art der tatsächlichen Nutzung*; Fachserie 3 Reihe 5.1; 2014.
- (2) Bundesamt für Verbraucherschutz und Lebensmittelsicherheit, *Absatz an Pflanzenschutzmitteln in der Bundesrepublik Deutschland - Ergebnisse der Meldungen gemäß § 64 Pflanzenschutzgesetz für das Jahr 2014*; 2015.

- (3) Schulz, R. Comparison of spray drift-and runoff-related input of azinphos-methyl and endosulfan from fruit orchards into the Lourens River, South Africa. *Chemosphere* **2001**, *45*, 543–551.
- (4) Liess, M.; Schulz, R.; Liess, M.-D.; Rother, B.; Kreuzig, R. Determination of insecticide contamination in agricultural headwater streams. *Water Research* **1999**, *33*, 239–247, Anfragen 38 VG Wort 1/99 Paper Sabine PDF.
- (5) Malaj, E.; Ohe, P. C. v. d.; Grote, M.; Kühne, R.; Mondy, C. P.; Usseglio-Polatera, P.; Brack, W.; Schäfer, R. B. Organic chemicals jeopardize the health of freshwater ecosystems on the continental scale. *Proceedings of the National Academy of Sciences* **2014**, *111*, 9549–9554.
- (6) Schulz, R. Field Studies on Exposure, Effects, and Risk Mitigation of Aquatic Nonpoint-Source Insecticide Pollution: A Review. *Journal of Environmental Quality* **2004**, *33*, 419–448, PDF Carola.
- (7) Schäfer, R. B.; Caquet, T.; Siimes, K.; Mueller, R.; Lagadic, L.; Liess, M. Effects of pesticides on community structure and ecosystem functions in agricultural streams of three biogeographical regions in Europe. *Science of the Total Environment* **2007**, *382*, 272–285.
- (8) Quevauviller, P.; Borchers, U.; Thompson, C.; Simonart, T. *The Water Framework Directive: Ecological and Chemical Status Monitoring*; John Wiley & Sons, 2008.
- (9) Poisot, T. Best publishing practices to improve user confidence in scientific software. *Ideas in Ecology and Evolution* **2015**, *8*.
- (10) Rauthe, M.; Steiner, H.; Riediger, U.; Mazurkiewicz, A.; Gratzki, A. A Central European precipitation climatology – Part I: Generation and validation of a high-resolution gridded daily data set (HYRAS). *Meteorologische Zeitschrift* **2013**, *22*, 235–256.



- (11) OGEwV, Verordnung zum Schutz der Oberflächengewässer (Oberflächengewässerverordnung). 2011; BGBl. I S. 1429.
- (12) European Union, DIRECTIVE 2013/39/EU OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. 2013.
- (13) Brock, T. C. M., Alix, A., Brown, C. D., Capri, E., Gottesbüren, B. E., Heimbach, F., Lythgo, C. M., Schulz, R., Streloke, M., Eds. *Linking aquatic exposure and effects: risk assessment of pesticides: EU and SETAC Europe workshop ELINK, Bari, Italy, and Wageningen, Netherlands*; CRC Press: Boca Raton, 2010.
- (14) Sprague, J. Measurement of pollutant toxicity to fish, II-Utilizing and applying bioassay results. *Water Research* **1970**, 4, 3–32.
- (15) Pesticide Action Network, PAN Pesticide Database. 2015; <http://www.pesticideinfo.org/>.
- (16) U.S. EPA, The ECOTOXicology knowledgebase (ECOTOX). 2015; <http://cfpub.epa.gov/ecotox/>.
- (17) Schüürmann, G.; Ebert, R.-U.; Kühne, R. Quantitative read-across for predicting the acute fish toxicity of organic compounds. *Environmental Science & Technology* **2011**, 45, 4616–4622.
- (18) EEA, Digital Elevation Model over Europe (EU-DEM). 2013; <http://www.eea.europa.eu/data-and-maps/data/eu-dem#tab-metadata>.
- (19) Holmgren, P. Multiple flow direction algorithms for runoff modelling in grid based elevation models: An empirical evaluation. *Hydrological Processes* **1994**, 8, 327–334.
- (20) Neteler, M.; Bowman, M. H.; Landa, M.; Metz, M. GRASS GIS: A multi-purpose open source GIS. *Environmental Modelling & Software* **2012**, 31, 124–130, pdf RS.

## Graphical TOC Entry

Some journals require a graphical entry for the Table of Contents. This should be laid out "print ready" so that the sizing of the text is correct. Inside the `tocentry` environment, the font used is Helvetica 8 pt, as required by *Journal of the American Chemical Society*. The surrounding frame is 9 cm by 3.5 cm, which is the maximum permitted for *Journal of the American Chemical Society* graphical table of content entries. The box will not resize if the content is too big: instead it will overflow the edge of the box. This box and the associated title will always be printed on a separate page at the end of the document.