

EDUARD SZÖCS

QUANTITATIVE ECOTOXICOLOGY

WITH R!

This document was created using L^AT_EX, knitr and the tufte book class.

Copyright © 2013 Eduard Szöcs



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

First draft, October 2013

Contents

1	<i>Introduction</i>	9
2	<i>The Measurement Process</i>	11
2.1	<i>Winsorized Mean and Standard Deviation</i>	11
2.2	<i>Probability Plotting</i>	12
3	<i>Bioaccumulation</i>	13
4	<i>Tests for Detection of Chronic Lethal and Sublethal Stress</i>	15
5	<i>Lethal and Other Quantal Responses to Stress</i>	17
5.1	<i>Fitting dose-response models</i>	17
6	<i>Population and Metapopulation Effects</i>	19
7	<i>Community Effects</i>	21
7.1	<i>Species Richness</i>	21
7.2	<i>Analyzing mesocosm data</i>	21
7.3	<i>Species Sensitivity Distributions</i>	21
	<i>R Session Info</i>	23

Bibliography 25

List of Figures

2.1	A histogramm of the so ₄ data.	11
-----	---	----

List of Tables

1

Introduction

```
require(devtools)  
install_github("qetx", "EDiLD")
```

```
require(qetx)
```


2

The Measurement Process

2.1 Winsorized Mean and Standard Deviation

The following sulfate concentrations (mg/L) were measured during a routine water quality survey of the Savannah River (South Carolina). The data is available in the `qetx` package ¹:

```
data(so4)
```

```
so4
## [1] 1.3 2.3 2.6 3.3 3.5 3.5 3.6 4.0 4.1 4.5 5.2 5.6
## [13] 5.7 6.1 6.2 6.5 6.9 7.1 7.7 7.9 9.9
```

```
length(so4)
```

```
## [1] 21
```

```
mean(so4)
```

```
## [1] 5.119
```

```
sd(so4)
```

```
## [1] 2.137
```

So there are 21 measurements with a mean of 5.12 mg/L and a standard deviation of 2.14 mg/L.

Suppose we have a detection limit of 2.5 mg/L and want to win-sorize values below LOD, i.e. replace the two lowest values by 2.6 mg/L and the two highest values by 7.7 mg/L.

¹ Note that in this case you do not have to assign the data to a name.




Figure 2.1: A histogram of the `so4` data.

Happily there is function in the `qetx` to do this for us: `winsor()`. This function takes a vector of values and a second argument specifying how many values should be winsorized (either by giving a LOD-value or the number of values on each side) ².

```
so4_w <- winsor(so4, lod = 2.5)
so4_w

## [1] 2.6 2.6 2.6 3.3 3.5 3.5 3.6 4.0 4.1 4.5 5.2 5.6
## [13] 5.7 6.1 6.2 6.9 6.5 7.1 7.7 7.7 7.7
## attr("width")
## [1] 2
```

² Look at the source of this function - type the function name into the console - to see which computations are done.

This give the expected results, moreover we see that on each end two observations where modified ³.

```
mean(so4_w)

## [1] 5.081

sd(so4_w)

## [1] 1.792

sd_winsor(so4_w)

## [1] 2.24
```

³ stored within the attribute 'width' of the resulting vector. **TODO: verbatim within sidenote.**

The Winsorized mean (\bar{x}_w) now is 5.08 mg/L, the standard deviation of the modified data set (s) is 1.79 mg/L and the Winsorized standard deviation (s_w) 2.24 mg/L.

2.2 Probability Plotting

3

Bioaccumulation

4

Tests for Detection of Chronic Lethal and Sub-lethal Stress

5

Lethal and Other Quantal Responses to Stress

5.1 Fitting dose-response models

6

Population and Metapopulation Effects

7

Community Effects

7.1 Species Richness

7.2 Analyzing mesocosm data

Principle Response Curves (PRC)¹ are commonly used for analyzing ecotoxicological mesocosm experiments. PRC is just a special form of Redundancy Analysis (RDA) and analyses the multivariate response to a treatment over time ².

Here we will analyze the pyrifos data set from the publication³ which is shipped with the vegan package.

¹ Van den Brink, P. and Ter Braak, C. (1999). Principal response curves: Analysis of time-dependent multivariate responses of biological community to stress. *Environmental Toxicology and Chemistry*, 18(2):138–148

² Legendre and Legendre, 2013

³ Van den Brink and Ter Braak, 1999

```
require(vegan)
data(pyrifos)
head(pyrifos[, c(1:10)])

##           Simve Daplo Cerpu Alogu Aloco Alore Aloaf Copsp
## w.4.c1 3.951      0      0      0      0      0      0 2.773
## w.4.c2 2.303      0      0      0      0      0      0 2.079
## w.4.c3 4.595      0      0      0      0      0      0 3.761
## w.4.c4 2.398      0      0      0      0      0      0 3.296
## w.4.c5 4.025      0      0      0      0      0      0 3.466
## w.4.c6 2.303      0      0      0      0      0      0 2.197
##           Ostsp Slyla
## w.4.c1 0.000 1.386
## w.4.c2 0.000 0.000
## w.4.c3 0.000 0.693
## w.4.c4 0.693 0.000
## w.4.c5 0.000 0.000
## w.4.c6 0.000 0.000
```

7.3 Species Sensitivity Distributions

R Session Info

```
sessionInfo()

## R version 3.0.2 (2013-09-25)
## Platform: x86_64-pc-linux-gnu (64-bit)
##
## locale:
##  [1] LC_CTYPE=en_US.UTF-8
##  [2] LC_NUMERIC=C
##  [3] LC_TIME=en_US.UTF-8
##  [4] LC_COLLATE=en_US.UTF-8
##  [5] LC_MONETARY=en_US.UTF-8
##  [6] LC_MESSAGES=en_US.UTF-8
##  [7] LC_PAPER=en_US.UTF-8
##  [8] LC_NAME=C
##  [9] LC_ADDRESS=C
## [10] LC_TELEPHONE=C
## [11] LC_MEASUREMENT=en_US.UTF-8
## [12] LC_IDENTIFICATION=C
##
## attached base packages:
## [1] stats      graphics  grDevices  utils      datasets
## [6] methods    base
##
## other attached packages:
## [1] vegan_2.1-35    lattice_0.20-23 permute_0.7-4
## [4] getx_0.0.1      knitr_1.5
##
## loaded via a namespace (and not attached):
## [1] evaluate_0.5.1 formatR_0.9    grid_3.0.2
## [4] highr_0.2.1    stringr_0.6.2 tools_3.0.2
```


Bibliography

Legendre, P. and Legendre, L. (2013). *Numerical ecology*. Elsevier, Amsterdam; Boston.

Van den Brink, P. and Ter Braak, C. (1999). Principal response curves: Analysis of time-dependent multivariate responses of biological community to stress. *Environmental Toxicology and Chemistry*, 18(2):138–148.