

# Statistical Ecotoxicology

## - Improving the utilization of data for ecological risk assessment

Eduard Szöcs

Institute for Environmental Sciences, University of Koblenz-Landau

Landau, 22.09.2016

# My field of research is somewhere between...




... Eco(-toxico)logy, Data Analysis & Programming

# Statistical Ecotoxicology

# Current use in ecotoxicology

- Ecological risk assessment (ERA) relies on statistics




# Current use in ecotoxicology


- ▶ Ecological risk assessment (ERA) relies on statistics
- ▶ Experiments with low replication

# Current use in ecotoxicology


- ▶ Ecological risk assessment (ERA) relies on statistics
- ▶ Experiments with low replication
- ▶ Usually analysed using Linear Models of transformed data
- ▶ Null Hypothesis Significance Testing ( $\Rightarrow$  NOEC)



# Statistical Power in current experimental designs in ecotoxicology is unacceptably low




# Generalized Linear Models can do better





# Generalized Linear Models can do better



Better abandon NOEC and use a regression design <sup>1</sup>...

<sup>1</sup> debated since 30 years.

# Monitoring Data

# Monitoring data...


- ▶ ... provides an opportunity to study large-scale dynamics of pesticides

# Monitoring data...

- ▶ ... provides an opportunity to study large-scale dynamics of pesticides
- ▶ ... provides the biggest amount of data available


# Monitoring data...

- ▶ ... provides an opportunity to study large-scale dynamics of pesticides
- ▶ ... provides the biggest amount of data available
- ▶ ... is really messy




# The biggest currently available dataset on

## pesticides



3,000 sites, 45,000 samples,  
500 pesticides

## invertebrates



14,000 sites, 27,000 samples,  
3000 taxa

Statistical Ecotoxicology



Monitoring Data



Software



Outlook



# Additional data on

## Sites

- ▶ catchment size
- ▶ agriculture within catchment

# Additional data on

## Sites

- ▶ catchment size
- ▶ agriculture within catchment

## Samples

- ▶ daily precipitation



# Additional data on

## Sites

- ▶ catchment size
- ▶ agriculture within catchment

## Samples

- ▶ daily precipitation

## Compounds

- ▶ RAC, LC50, EQS
- ▶ chemical group
- ▶ identifiers
- ▶ properties

# Results - Thresholds

Statistical Ecotoxicology



Monitoring Data




Software



Outlook



# Results - Thresholds



# Results - Precipitation & Seasonality

- Used a mixture model

$$RQ_i \sim ZAGA(\mu_i, \sigma, \pi_i) = \begin{cases} (1 - \pi_i) & \text{if } y < LOQ \\ \pi_i \times f_{\text{Gamma}}(\mu_i, \sigma) & \text{if } y \geq LOQ \end{cases}$$

# Results - Precipitation & Seasonality

- ▶ Used a mixture model

$$RQ_i \sim ZAGA(\mu_i, \sigma, \pi_i) = \begin{cases} (1 - \pi_i) & \text{if } y < LOQ \\ \pi_i \times f_{Gamma}(\mu_i, \sigma) & \text{if } y \geq LOQ \end{cases}$$


- ▶ Precipitation and Quarter as predictors
- ▶ Site within state as random intercept

# Results - Precipitation & Seasonality

- ▶ Used a mixture model

$$RQ_i \sim ZAGA(\mu_i, \sigma, \pi_i) = \begin{cases} (1 - \pi_i) & \text{if } y < LOQ \\ \pi_i \times f_{\text{Gamma}}(\mu_i, \sigma) & \text{if } y \geq LOQ \end{cases}$$

- ▶ Precipitation and Quarter as predictors
- ▶ Site within state as random intercept
- ▶ Precipitation before sampling increases RQ




# Results - Precipitation & Seasonality

- ▶ Used a mixture model

$$RQ_i \sim ZAGA(\mu_i, \sigma, \pi_i) = \begin{cases} (1 - \pi_i) & \text{if } y < LOQ \\ \pi_i \times f_{\text{Gamma}}(\mu_i, \sigma) & \text{if } y \geq LOQ \end{cases}$$

- ▶ Precipitation and Quarter as predictors
- ▶ Site within state as random intercept
- ▶ Precipitation before sampling increases RQ
- ▶ Summer higher RQ, but compound specific




# Results - Small Water Bodies (SWB)

- ▶ most streams are *small*
- ▶ refuge of biodiversity
- ▶ High risk of pollution
  - ▶ adjacency to fields
  - ▶ low dilution



# Results - Small Water Bodies (SWB)

- ▶ most streams are *small*
- ▶ refuge of biodiversity
- ▶ High risk of pollution
  - ▶ adjacency to fields
  - ▶ low dilution
- ▶ **Neonicotinoids**
- ▶ **up to 244x RAC**
- ▶ **ecological effects likely**



Software

# Biologists and Chemists face similar problems...

## Names

*Osmia rufa*, *Osmia bicornis*,  
*Osmia ruffa*, *Osmia unilandaui*,  
*Osmia spec.*

Chlorpyrifos, Chlorpyrifos,  
Chlorpyrifos, Chlorpyrifos-ethyl,  
Chlorpyrifos

# Biologists and Chemists face similar problems...

## Names

*Osmia rufa*, *Osmia bicornis*,  
*Osmia ruffa*, *Osmia unilandaui*,  
*Osmia spec.*

Chlorpyrifos, Chlorpyrifos,  
Chlorpyrifos, Chlorpyrifos-ethyl,  
Chlorpyrifot

## Hierarchies

Hymenoptera/ Apoidea/  
Megachilidae/ *Osmia*/ *rufa*

organophosphate, ester,  
insecticide

# Biologists and Chemists face similar problems...

## Names

*Osmia rufa*, *Osmia bicornis*,  
*Osmia ruffa*, *Osmia unilandaui*,  
*Osmia spec.*

Chlorpyrifos, Chlorpyrifos,  
Chlorpyrifos, Chlorpyrifos-ethyl,  
Chlorpyrifos

## Hierarchies

Hymenoptera/ Apoidea/  
Megachilidae/ *Osmia*/ *rufa*

organophosphate, ester,  
insecticide

## Attributes

Wing length, Mass, Season

Mass,  $K_{OW}$ ,  $LC_{50}$

# Biologists and Chemists face similar problems...

## Names

*Osmia rufa*, *Osmia bicornis*,  
*Osmia ruffa*, *Osmia unilandaui*,  
*Osmia spec.*

Chlorpyrifos, Chlorpyrifos,  
Chlorpyrifos, Chlorpyrifos-ethyl,  
Chlorpyrifos

## Hierarchies

Hymenoptera/ Apoidea/  
Megachilidae/ *Osmia*/ *rufa*

organophosphate, ester,  
insecticide

## Attributes

Wing length, Mass, Season

Mass,  $K_{OW}$ ,  $LC_{50}$

## Identifiers

NCBI, ITIS, EOL, ...

2921-88-2, Clc1c(OP(=S)[...],  
InChI=1S/C9H11C[...],  
SBPBAQFW[...], CSID,...

# Biologists and Chemists face similar problems...

## Names

*Osmia rufa*, *Osmia bicornis*,  
*Osmia ruffa*, *Osmia unilandaui*,  
*Osmia spec.*

Chlorpyrifos, Chlorpyrifos,  
Chlorpyrifos, Chlorpyrifos-ethyl,  
Chlorpyrifot

## Hierarchies

Hymenoptera/ Apoidea/  
Megachilidae/ *Osmia*/ *rufa*

organophosphate, ester,  
insecticide

## Attributes

Wing length, Mass, Season

Mass,  $K_{OW}$ ,  $LC_{50}$

## Identifiers

NCBI, ITIS, EOL, ...

2921-88-2, Clc1c(OP(=S)[...],  
InChI=1S/C9H11C[...],  
SBPBAQFW[...], CSID,...

---

## Amount of data

2993 taxa

489 pesticides  
(+ 590 other organics)

# Instead of wasting time...

Statistical Ecotoxicology



Monitoring Data



Software



Outlook






# Instead of wasting time...

## taxize - taxonomic search and retrieval in R



# Instead of wasting time...

## taxize - taxonomic search and retrieval in R



Statistical Ecotoxicology



Monitoring Data



Software



Outlook



# Instead of wasting time...

Statistical Ecotoxicology



Monitoring Data



Software



Outlook



# Instead of wasting time...

*"webchem ...likely saved hundreds of working hours"*

Münch (2016)

Statistical Ecotoxicology

○  
○○○○

Monitoring Data

○  
○○○○○○

Software


○  
○○●

Outlook

○○  
○○○○

# Instead of wasting time...

*"webchem ...likely saved hundreds of working hours"*



Statistical Ecotoxicology



Monitoring Data



Software



Outlook



# Conclusions from my PhD

- ▶ Change your model, not your data

# Conclusions from my PhD

- ▶ Change your model, not your data
- ▶ Ultimately ban NOEC

# Conclusions from my PhD

- ▶ Change your model, not your data
- ▶ Ultimately ban NOEC
- ▶ Monitoring data can be used to
  - ▶ study pesticide dynamics
  - ▶ inform ERA



# Conclusions from my PhD

- ▶ Change your model, not your data
- ▶ Ultimately ban NOEC
- ▶ Monitoring data can be used to
  - ▶ study pesticide dynamics
  - ▶ inform ERA
- ▶ Agricultural SWB at risk from pesticides


# Conclusions from my PhD

- ▶ Change your model, not your data
- ▶ Ultimately ban NOEC
- ▶ Monitoring data can be used to
  - ▶ study pesticide dynamics
  - ▶ inform ERA
- ▶ Agricultural SWB at risk from pesticides
- ▶ Handling big eco(toxico-)logical data not easy
  - ▶ now easier

# Outlook


# Analysing chemical concentrations is not easy, because of

- ▶ continuous distribution in  $\mathbb{R}_0^+$
- ▶ censoring ( $x < \text{LOQ}$ )




# Analysing chemical concentrations is not easy, because of


- ▶ continuous distribution in  $\mathbb{R}_0^+$
- ▶ censoring ( $x < \text{LOQ}$ )
- ▶ non-linearity (season, trends)
- ▶ dependency (spatial, temporal)
- ▶ missing data




# Dealing with censored, non-normal data



# Dealing with censored, non-normal data




# Dealing with censored, non-normal data






# Dealing with censored, non-normal data



# Dealing with censored, non-normal data



Guidance how to model environmental concentrations is missing


# Temporal dynamics of pesticide occurrence

- ▶ Pesticides show compound specific dynamics
- ▶ Mixture dynamics? - Multivariate response.

# Temporal dynamics of pesticide occurrence

- ▶ Pesticides show compound specific dynamics
- ▶ Mixture dynamics? - Multivariate response.
- ▶ Seasonality, Trends (Fade out...)?

$$y = \beta_0 + f_{\text{seasonal}}(x_1) + f_{\text{trend}}(x_2) + \epsilon; \epsilon \sim ???$$



# Statistical Ecotoxicology

- Improving the utilization of data for ecological risk assessment

Eduard Szöcs

Institute for Environmental Sciences, University of Koblenz-Landau



<http://edild.github.io/>



@EduardSzoecs



[szoecs@uni-landau.de](mailto:szoecs@uni-landau.de)



[https://github.com/edild/talk\\_work2](https://github.com/edild/talk_work2)

