1

ASSIGNMENT 1

YENIGALLA SAMYUKTHA EE20MTECH14019

Abstract—This document illustrates the ratio in which a line divides another line joining two points

Download all python codes from

https://github.com/EE20MTECH14019/EE5609/ tree/master/Assignment _1/Codes

and latex-tikz codes from

https://github.com/EE20MTECH14019/EE5609/ tree/master/Assignment 1

1 Problem

In what ratio is the line joining $\begin{pmatrix} -1\\1 \end{pmatrix}$ and $\begin{pmatrix} 5\\7 \end{pmatrix}$ is divided by the line

$$\begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = 4 \tag{1.0.1}$$

2 EXPLANATION

The point **X** divides the line segment joining the two points $\mathbf{A} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 5 \\ 7 \end{pmatrix}$ in ratio k : 1. Then,

$$\mathbf{X} = \frac{(k\mathbf{B} + \mathbf{A})}{(k+1)} \tag{2.0.1}$$

3 solution

From the equation (2.0.1)

$$(k+1)\mathbf{X} = k\mathbf{B} + \mathbf{A} \tag{3.0.1}$$

Let
$$\mathbf{n} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$\implies (k+1)\mathbf{n}^T\mathbf{X} = \mathbf{n}^T(k\mathbf{B} + \mathbf{A}) \qquad (3.0.2)$$

$$\implies k(\mathbf{n}^T \mathbf{X} - \mathbf{n}^T \mathbf{B}) = \mathbf{n}^T \mathbf{A} - \mathbf{n}^T \mathbf{X}$$
 (3.0.3)

$$\implies k = \frac{\mathbf{n}^T \mathbf{A} - \mathbf{n}^T \mathbf{X}}{\mathbf{n}^T \mathbf{X} - \mathbf{n}^T \mathbf{B}}$$
 (3.0.4)

Hence on solving the equation (3.0.4) using

$$\mathbf{n}^T \mathbf{X} = 4 \tag{3.0.5}$$

The line $\begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = 4$ divides the line joining points $\mathbf{A} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 5 \\ 7 \end{pmatrix}$ in the ratio k = 1/2




Fig. 0: Line as $(1\ 1)x=4$ intersecting the line joining points A and B