1

Assignment 16

Yenigalla Samyuktha EE20MTECH14019

Abstract—This document solves a problem on Jordan form of a complex matrix.

Download all latex-tikz codes from

https://github.com/EE20MTECH14019/EE5609/ tree/master/Assignment 16

1 Problem

How many possible Jordan forms are there for a 6×6 complex matrix with characteristic polynomial $(x + 2)^4 (x - 1)^2$?

2 EXPLANATION

From the characteristic polynomial,

$$(x+2)^4 (x-1)^2$$
 (2.0.1)

We get the eigen values of the 6×6 complex matrix as,

$$\lambda_i = \{-2, -2, -2, -2, 1, 1\}$$
 (2.0.2)

The minimal polynomial for a matrix with characteristic polynomial must have both (x + 2) and (x - 1) as factors and it must divide (2.0.1). Hence the minimal polynomial will be of the form

$$p = (x+2)^a (x-1)^b$$
, $a \le 4, b \le 2$ (2.0.3)

So, there are 8 different possibilities for a minimal polynomial. Let us note that one minimal polynomial may correspond to more than one Jordan forms. The possible Jordan blocks associated with the eigen values -2 and 1 are given in the Table 1. From Table 1, we can say that there are $5 \times 2 = 10$ possible Jordan forms. For example, the Jordan forms corresponding to minimal polynomial

 $p = (x + 2)^2 (x - 1)$ built from the Jordan blocks can be,

$$\mathbf{J} = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$
 (2.0.4)

and

$$\mathbf{J} = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$
 (2.0.5)

3 Answer

Therefore 10 different Jordan forms are possible for a 6×6 complex matrix with characteristic polynomial $(x + 2)^4 (x - 1)^2$.

Factor	Possible Jordan blocks	
(x+2)	$ \begin{pmatrix} -2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} $	(2.0.6)
	$ \begin{pmatrix} -2 & 1 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} $	(2.0.7)
	$\begin{pmatrix} -2 & 1 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & -2 \end{pmatrix}, \begin{pmatrix} -2 & 1 & 0 & 0 \\ 0 & -2 & 1 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}$	(2.0.8)
	$ \begin{pmatrix} -2 & 1 & 0 & 0 \\ 0 & -2 & 1 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & -2 \end{pmatrix} $	(2.0.9)
(x - 1)	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	(2.0.10)
	$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$	(2.0.11)

TABLE 1: Possible Jordan Blocks