

EE5609: Matrix Theory

Assignment-2

M Pavan Manesh
EE20MTECH14017

Abstract—This document contains a solution for proving the determinant of the given matrix is zero.

Download the python codes from

<https://github.com/pavanmanesh/EE5609/blob/master/Assignment2/codes>

and latex-tikz codes from

<https://github.com/pavanmanesh/EE5609/tree/master/Assignment2>

Let A be the given matrix and x be a vector where x is

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \quad (3.0.4)$$

From (2.0.1), we can write the augmented form in this way:

$$\begin{pmatrix} -a & 0 & -c & 0 \\ 0 & a & -b & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (3.0.5)$$

upon solving the above , the value of x:

$$\mathbf{x} = \begin{pmatrix} -\frac{c}{a}x_3 \\ \frac{b}{a}x_3 \\ x_3 \end{pmatrix} \quad (3.0.6)$$

$$\mathbf{x} = \begin{pmatrix} -c \\ b \\ a \end{pmatrix} \quad (3.0.7)$$

1 PROBLEM

$$\begin{vmatrix} 0 & a & -b \\ -a & 0 & c \\ b & c & 0 \end{vmatrix} = 0 \quad (1.0.1)$$

2 PROPERTIES

Properties used for solving this problem:

$$A\mathbf{x} = \mathbf{0} \quad (2.0.1)$$

A has a zero eigen value if x has a nontrivial solution.

From (2.0.1), one of the eigen value is equal to 0. We know that the determinant of a matrix is the product of the eigen values. so, if one of the values is zero , $|A| = 0$ (i.e., A is singular). So,

$$\begin{vmatrix} 0 & a & -b \\ -a & 0 & c \\ b & c & 0 \end{vmatrix} = 0 \quad (3.0.8)$$

3 SOLUTION

Converting A into reduced echelon form:

$$\begin{pmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{pmatrix} \xrightarrow{R_1 \leftrightarrow R_2} \begin{pmatrix} -a & 0 & -c \\ 0 & a & -b \\ b & c & 0 \end{pmatrix} \quad (3.0.1)$$

$$\xrightarrow{R_3 \leftarrow a \times R_3 + b \times R_1} \begin{pmatrix} -a & 0 & -c \\ 0 & a & -b \\ 0 & ac & -bc \end{pmatrix} \quad (3.0.2)$$

$$\xrightarrow{R_3 \leftarrow R_3 - c \times R_2} \begin{pmatrix} -a & 0 & -c \\ 0 & a & -b \\ 0 & 0 & 0 \end{pmatrix} \quad (3.0.3)$$