

EE5609: Matrix Theory

Assignment-8

M Pavan Manesh
EE20MTECH14017

Abstract—This document explains how to find a row-reduced matrix which is row equivalent to equation (1.0.1),

Download all latex-tikz codes from

<https://github.com/pavanmanesh/EE5609/tree/master/Assignment8>

and all python codes from

<https://github.com/pavanmanesh/EE5609/tree/master/Assignment8/codes>

$$\mathbf{E}_{31}\mathbf{E}_{21} = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -i & 0 & 1 \end{pmatrix} \quad (3.0.1)$$

$$\mathbf{E}_{31}\mathbf{E}_{21}\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -i & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ 2 & 2 \\ i & 1+i \end{pmatrix} \quad (3.0.2)$$

$$\Rightarrow \mathbf{A}_1 = \mathbf{E}_{31}\mathbf{E}_{21}\mathbf{A} = \begin{pmatrix} 1 & -i \\ 0 & 2+2i \\ 0 & -i \end{pmatrix} \quad (3.0.3)$$

Step 2: Performing $R_2 \leftarrow \frac{1}{2+2i}R_2$ given by \mathbf{A}_1 on equation (3.0.3),

$$\mathbf{D}_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{4}(1-i) & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (3.0.4)$$

$$\mathbf{D}_1\mathbf{A}_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{4}(1-i) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ 0 & 2+2i \\ 0 & -i \end{pmatrix} \quad (3.0.5)$$

$$\Rightarrow \mathbf{A}_2 = \mathbf{D}_1\mathbf{A}_1 = \begin{pmatrix} 1 & -i \\ 0 & 1 \\ 0 & -i \end{pmatrix} \quad (3.0.6)$$

Step 3: Performing $R_3 \leftarrow R_3 + iR_2$ given by \mathbf{E}_{32} on equation (3.0.6),

$$\mathbf{E}_{32} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & i & 1 \end{pmatrix} \quad (3.0.7)$$

$$\mathbf{E}_{32}\mathbf{A}_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & i & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ 0 & 1 \\ 0 & -i \end{pmatrix} \quad (3.0.8)$$

$$\Rightarrow \mathbf{A}_3 = \mathbf{E}_{32}\mathbf{A}_2 = \begin{pmatrix} 1 & -i \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (3.0.9)$$

1 PROBLEM

Find a row-reduced matrix which is row equivalent to,

$$\mathbf{A} = \begin{pmatrix} 1 & -i \\ 2 & 2 \\ i & 1+i \end{pmatrix} \quad (1.0.1)$$

What are the solutions of $\mathbf{A}\mathbf{x}=\mathbf{0}$?

2 THEOREM

Let \mathbf{R} be a row-reduced echelon matrix which is row equivalent to \mathbf{A} . Then the systems

$$\mathbf{A}\mathbf{x} = \mathbf{0}, \mathbf{R}\mathbf{x} = \mathbf{0} \quad (2.0.1)$$

have the same solutions.

3 SOLUTION

Step 1: Performing $R_2 \leftarrow R_2 - 2R_1$ and $R_3 \leftarrow R_3 - iR_1$ given by elementary matrix $\mathbf{E}_{31}\mathbf{E}_{21}$ on the

Step 4: Performing $R_1 \leftarrow R_1 + iR_2$ given by \mathbf{E}_{12} on equation (3.0.9),

$$\mathbf{E}_{12} = \begin{pmatrix} 1 & i & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (3.0.10)$$

$$\mathbf{E}_{12}\mathbf{A}_3 = \begin{pmatrix} 1 & i & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -i \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (3.0.11)$$

$$\Rightarrow \mathbf{A}_4 = \mathbf{E}_{12}\mathbf{A}_3 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (3.0.12)$$

\therefore Row-reduced matrix of \mathbf{A} given by equation (1.0.1) is,

$$\mathbf{A} = \begin{pmatrix} 1 & -i \\ 2 & 2 \\ i & 1+i \end{pmatrix} \xrightarrow{RREF} \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (3.0.13)$$

From (2.0.1) and (3.0.13), the only solution of $\mathbf{A}\mathbf{x} = \mathbf{0}$ is,

$$\mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (3.0.14)$$