#### 1

# Matrix theory Assignment 1

## M Pavan Manesh

Abstract—This document explains the equation of the line passing through the point of intersection of the lines that has equal intercepts on the axes

Download all python codes from

https://github.com/pavanmanesh/EE5609/tree/master/codes

and latex-tikz codes from

https://github.com/pavanmanesh/EE5609

#### 1 Problem

Find the equation of the line passing through the point of intersection of the lines

$$(4 7)\mathbf{x} = 3$$
$$(2 -3)\mathbf{x} = -1$$

that has equal intercepts on the axes

## 2 Solution

The above two line equations can be expressed as the matrix equation

$$\begin{pmatrix} 4 & 7 \\ 2 & -3 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 3 \\ -1 \end{pmatrix} \tag{2.0.1}$$

Constructing the augmented matrix

$$\begin{pmatrix} 4 & 7 & 3 \\ 2 & -3 & -1 \end{pmatrix} \tag{2.0.2}$$

Transforming the matrix into row-echelon form

$$\begin{pmatrix}
4 & 7 & 3 \\
2 & -3 & -1
\end{pmatrix} \xrightarrow{R2 \leftarrow 2R2 - R1}$$

$$\begin{pmatrix}
4 & 7 & 3 \\
0 & -13 & -5
\end{pmatrix} \xrightarrow{R2 \leftarrow -R2/13, R1 \leftarrow R1/4}$$

$$\begin{pmatrix}
1 & 7/4 & 3/4 \\
0 & 1 & 5/13
\end{pmatrix} \xrightarrow{R1 \leftarrow R1 - 7/4R1/4}$$

$$\begin{pmatrix}
1 & 0 & 2/26 \\
0 & 1 & 5/13
\end{pmatrix}$$
(2.0.3)

The solution for x can be written as

$$\mathbf{x} = \begin{pmatrix} 2/26 \\ 5/13 \end{pmatrix} \tag{2.0.4}$$

Thus, The point of intersection is at point (2/26, 5/13) i.e. (0.07, 0.38)

Let the equation of the line be

$$\mathbf{n}^T \mathbf{x} = c \implies \mathbf{x}^T \mathbf{n} = c \tag{2.0.5}$$

Let the intercepts be a,b on the x and y axis respectively. Then,

$$\begin{pmatrix} a & 0 \end{pmatrix} \mathbf{n} = c \tag{2.0.6}$$

$$\begin{pmatrix} 0 & b \end{pmatrix} \mathbf{n} = c \tag{2.0.7}$$

resulting in the matrix equation

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \mathbf{n} = c \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$
 (2.0.8)

As the intercepts are equal, Let a=b

$$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \mathbf{n} = c \begin{pmatrix} 1 \\ 1 \end{pmatrix} \tag{2.0.9}$$

$$\mathbf{n} = \frac{c}{a} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \tag{2.0.10}$$

As the line passes through point of intersection, We can use the equation (2.0.4) in equation (2.0.5) to find the value of c

$$c = \begin{pmatrix} 1 & 1 \end{pmatrix} \begin{pmatrix} \frac{2}{26} \\ \frac{2}{26} \end{pmatrix}$$
 (2.0.11)

$$\implies c = \frac{2}{26} + \frac{2}{26} \implies c = \frac{6}{13}$$
 (2.0.12)

So, the equation of line can be written as

$$\implies \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = \frac{6}{13} \tag{2.0.13}$$




Fig. 0: The intercepts of the required line are equal