

8.3.12

EE24BTECH11012 - Bhavanisankar G S

QUESTION :

Find the area bounded by the curves $\{(x, y) : y \geq x^2 \text{ and } y = |x|\}$

SOLUTION :

Theoretical :

1) FINDING THE POINT OF INTERSECTION :

$$y = x^2 \quad (1.1)$$

$$y = |x| \quad (1.2)$$

By the symmetry of the equations, the required area is double the area of

$$y = x^2 \text{ for } x \geq 0 \quad (1.3)$$

$$y = x \text{ for } x \geq 0 \quad (1.4)$$

Clearly, the points of intersection are

$$x = 0 \quad (1.5)$$

$$x = 1 \quad (1.6)$$

2) EVALUATING THE INTEGRAL :

From the graph, it can be seen that

$$x \geq x^2 \text{ for } 0 \leq x \leq 1 \quad (2.1)$$

Hence, the integral becomes

$$A = 2 \left(\int_0^1 (x - x^2) dx \right) \quad (2.2)$$

$$A = 2 \left(\left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 \right) \quad (2.3)$$

$$A = 2 \left(\frac{1}{2} - \frac{1}{3} \right) \quad (2.4)$$

$$A = 2 \left(\frac{1}{6} \right) \quad (2.5)$$

$$A = \frac{1}{3} \quad (2.6)$$

Hence, the area bounded by the given curves is $\frac{1}{3}$.

Simulation :

- 1) For a general interval, say $[a, b]$, split up the intervals into n parts such that

$$h = \frac{b - a}{n} \quad (1.1)$$

- 2) Consider the points

$$x_0 = a \quad (2.1)$$

$$x_n = b \quad (2.2)$$

$$x_{i+1} = x_i + h \quad (2.3)$$

- 3) **Trapezoid rule :**

Summing the areas of the trapezoids formed, we have

$$f(x) = x - x^2 \quad (3.1)$$

$$A \approx \frac{h}{2} ((f(x_0) + f(x_1)) + (f(x_1) + f(x_2)) + \cdots + (f(x_{n-1}) + f(x_n))) \quad (3.2)$$

$$A \approx \frac{h}{2} \left(f(x_0) + 2 \sum_{i=1}^{n-1} f(x_i) + f(x_n) \right) \quad (3.3)$$

In the given question,

$$a = 0 \quad (3.4)$$

$$b = 1 \quad (3.5)$$

Clearly,

$$f(a) = f(b) = 0 \quad (3.6)$$

since both the curves have $(0, 0)$ and $(1, 1)$ as their common points. Simplifying from (1.1) and (3.3), we have

$$A \approx \frac{1}{n} \left(\sum_{i=1}^{n-1} x_i - x_i^2 \right) \quad (3.7)$$

$$A \approx \frac{1}{n} \left(\sum_{i=1}^{n-1} \frac{i}{n} - \left(\frac{i}{n} \right)^2 \right) \quad (3.8)$$

$$A \approx \frac{1}{n^2} \left(\sum_{i=1}^{n-1} \left(i - \frac{i^2}{n} \right) \right) \quad (3.9)$$

Consider

$$A_{n+1} = A_n + \frac{h}{2} (y_n + y_{n+1}) \quad (3.10)$$

$$A_{n+1} = A_n + \frac{h}{2} (y_n + (y_n + h y'_n)) \quad (3.11)$$

$$A_{n+1} = A_n + \frac{h}{2} (y_n + (y_n + h(1 - 2x_n))) \quad (3.12)$$

$$A_{n+1} = A_n + \frac{h}{2} (2y_n + h(1 - 2x_n)) \quad (3.13)$$

which is the required difference equation.

- 4) The above equation can be coded to obtain the area bounded by the two curves.
- 5) It can be seen that the approximate solution turns out to be 0.3333333299999998.




Fig. 5.1: Plot of the given question.