

11.16.3.8.5

EE24BTECH11012 - Bhavanisankar G S

QUESTION :

Three coins are tossed once. Find the probability of getting no head.

SOLUTION :

Variable name	Description
S	Sample space
X	Random variable corresponding to the number of heads
p	Toss corresponding to head
$F_{\mathbf{X}}(x)$	Cumulative distribution function (CDF)
$p_{\mathbf{X}}(x)$	Probability Mass function (PMF)

Let us assume the random variable to be the sum of three Bernoulli Random Variables.

$$\mathbf{X} = \mathbf{X}_1 + \mathbf{X}_2 + \mathbf{X}_3 \quad (0.1)$$

$$\mathbf{X}_i = \begin{cases} 1 & , \text{ Outcome - head} \\ 0 & , \text{ Outcome - tail} \end{cases} \quad (0.2)$$

$$\Rightarrow p_{X_i}(k) = \begin{cases} 1 - p & , k = 0 \\ p & , k = 1 \end{cases} \quad (0.3)$$

Considering all the outcomes as equally likely, we have

$$p = \frac{1}{2} \quad (0.4)$$

For the given question, let \mathbf{X} denote the number of heads. The sample space corresponding to the given scenario is tabulated below.

Event	Sample space
$p_{\mathbf{X}}(0)$	$\{TTT\}$
$p_{\mathbf{X}}(1)$	$\{TTH, THT, HTT\}$
$p_{\mathbf{X}}(2)$	$\{HHT, HTH, THH\}$
$p_{\mathbf{X}}(3)$	$\{HHH\}$

By the properties of Z-transform of **Probability Mass Function**, we have

$$M_{\mathbf{X}}(z) = M_{\mathbf{X}_1}(z)M_{\mathbf{X}_2}(z)M_{\mathbf{X}_3}(z) \quad (0.5)$$

$$M_{\mathbf{X}_1} = \sum_{n \rightarrow -\infty}^{\infty} p_{\mathbf{X}_1}(n)z^{-n} = p + (1-p)z^{-1} \quad (0.6)$$

$$M_{\mathbf{X}_2} = \sum_{n \rightarrow -\infty}^{\infty} p_{\mathbf{X}_2}(n)z^{-n} = p + (1-p)z^{-1} \quad (0.7)$$

$$M_{\mathbf{X}_3} = \sum_{n \rightarrow -\infty}^{\infty} p_{\mathbf{X}_3}(n)z^{-n} = p + (1-p)z^{-1} \quad (0.8)$$

$$M_{\mathbf{X}}(z) = \left(p + (1-p)z^{-1}\right)^3 \quad (0.9)$$

$$= \sum_{k \rightarrow -\infty}^{\infty} {}^3C_k p^{3-k} (1-p)^k z^{-k} \quad (0.10)$$

$$p_{\mathbf{X}}(k) = {}^3C_k p^{3-k} (1-p)^k \quad (0.11)$$

Substituting (0.4) in (0.11), we have

$$p_{\mathbf{X}}(k) = \frac{{}^3C_k}{8} \quad (0.12)$$

The PMF is then given by -

$$p_{\mathbf{X}}(k) = \begin{cases} \frac{{}^3C_k}{8} & 0 \leq k \leq 3, k \in \mathbb{W} \\ 0 & \text{otherwise} \end{cases} \quad (0.13)$$

$$\Rightarrow p_{\mathbf{X}}(k) = \begin{cases} \frac{1}{8} & k = 0 \\ \frac{3}{8} & k = 1 \\ \frac{3}{8} & k = 2 \\ \frac{1}{8} & k = 3 \\ 0 & \text{otherwise} \end{cases} \quad (0.14)$$

The corresponding **Cumulative Distribution Function** can then be written as -

$$F_{\mathbf{X}}(k) = Pr(\mathbf{X} \leq k) \quad (0.15)$$

$$= \sum_{x=0}^k p_{\mathbf{X}}(x) \quad (0.16)$$

$$\Rightarrow F_{\mathbf{X}}(k) = Pr(\mathbf{X} \leq k) = \begin{cases} 0 & x < 0 \\ \frac{1}{8} & 0 \leq x < 1 \\ \frac{1}{2} & 1 \leq x < 2 \\ \frac{7}{8} & 2 \leq x < 3 \\ 1 & x \geq 3 \end{cases} \quad (0.17)$$

$$F_X(0) = P(\mathbf{X} \leq 0) \quad (0.18)$$

$$= \frac{1}{8} \quad (0.19)$$




Fig. 0.1: Probability Mass Funtion




Fig. 0.2: Cumulative Distribution Function




Fig. 0.3: Simulation