

# Uncertain Lines: Explaining phenotype of *Apistogramma species*

Jessica Bullock  
EEB313


*Apistogramma:*  
'uncertain line'








Wide geographic range  
Morphology varies  
Degree of dimorphism varies  
Complexity of pattern varies



*A. atahualpa*




*A. bitaeniata*



My hypothesis is that the visual conditions where the fish live are responsible for the phenotype of the species that evolve.

I predict that analysis will cluster species best by water type rather than evolutionary history.






# Tempo and rates of diversification in the South American cichlid genus *Apistogramma* (Teleostei: Perciformes: Cichlidae)

Christelle Tougard, Carmen R. García Dávila, Uwe Römer, Fabrice Duponchelle, Frédérique Cerqueira, Emmanuel Paradis, Bruno Guinand, Carlos Angulo Chávez, Vanessa Salas, Sophie Quérrouil, Susana Sirvas, Jean-François Renno

Published: September 5, 2017 • <https://doi.org/10.1371/journal.pone.0182618>

## The Sources



# The Data

Eco data - (Römer 2006) categorical ecological data, including water body type, water type, habitat type, and river system, for 102 species of *Apistogramma*

Character data - (Römer 2006) 51 phenotype traits including colour pattern, body shape, and dentition for 102 species of *Apistogramma*

Lineage data - (Tougard *et al*) – lineage 1-4 as a categorical trait for 31 species of *Apistogramma*



# Data preparation

Species names – updated taxonomic changes since 2006 (many)

Character data – omitted context-dependent behavioural traits

Joined datasets by species


“?” -> NA

“occasionally” -> yes




Filtered out rows with missing data



Filtered out columns with only one level

MCA factor map











| Dimension 1          | r2        |
|----------------------|-----------|
| Lineage              | 0.9304433 |
| Body form            | 0.7619608 |
| Infraorbital pores   | 0.7619608 |
| Longitudinal band    | 0.7172129 |
| Caudal spot          | 0.7093105 |
|                      |           |
| Dimension 2          | r2        |
| Head-body proportion | 0.9139043 |
| Lineage              | 0.9136878 |
| Jaw spot             | 0.7130203 |
| Chin spot            | 0.6938044 |
| Chest spot           | 0.6105135 |






Strength of relationships between Apistogramma species



# Combined data tree with Tougard et al lineages, UPGMA method



## References:

Römer U. 2006. Cichlid Atlas Volume 2. 1<sup>st</sup> Ed. Melle, Germany: Mergus.

Tougard C, García Dávila CR, Römer U, Duponchelle F, Cerqueira F, Paradis E, Guinand B, Angulo Chávez C, Salas V, Quérrouil S, Sirvas S, Renno JF. 2017. Tempo and rates of diversification in the South American cichlid genus *Apistogramma* (Teleostei: Perciformes: Cichlidae). PLoS One. Sep 5;12(9):e0182618.