EIGSEP Memo 008

Marjum Pass 2026-07: image-domain modeling of the sky, the ground and the Sun

EIGSEP Team

Living document. This version: 2026-11-06

Abstract

How calibrated Marjum spectra are modelled in the image domain: as the beam-weighted sum of a Galactic sky map, a terrain at one uniform temperature below the DEM horizon, a receiver offset, and the Sun. The memo covers the model and the degeneracies its geometry leaves, the night fit to the calibrated 07-17/18 data, and the Sun. Box-air records solar radio bursts at the times independent spectrographs recorded them, which identifies solar emission in the data. The quiet Sun is harder: the ridges hide it for hours, so it appears only at a handful of terrain crossings. The one calibrated crossing, 15:53 UTC on 07-17, shows an excess 2.0 times the concurrent Sun's prediction (1.5 scaled by the template amplitude), of the size the raw data give at the other crossings; none of them detects the Sun convincingly. Each measurement is shown against the raw data around it. Regressions on the rotating antenna give a repeatable ground term of 189–333 K at 125–180 MHz. The excess of thousands of kelvin below 85 MHz that earlier versions found there does not appear in the flagged raster on the present geometry, though it remains in unflagged data and in the 30 m window. Over phase C, raw power at 62–82 MHz improves with a slow additive daily term; the real sky beats skies shifted in sidereal time in each half of phase C but not over the whole, so it is not shown to be sky. With 20-min gain knots the phase-C ground term at 110–205 MHz agrees with the 3-h fit's to within a factor of two, but the fast gain takes up much of the sky's variation, and it follows neither a recorded stream nor the receiver's measured gain. The antenna's heights follow the rebuilt mode table: the 07-15 night and the 07-17 sunset were on the ground, and are left out until a horizon profile at 2 m exists. Every number, figure and table is computed by `make_products.py` in this directory from the products under `derived/ground_sky/` (Appendix A). The kelvin scale is memo 004's (`tcal@v0004`), the beam is HFSS rotated by the mount of memo 005's `empirical_raster_v0017`, and pointing is memo 003's `pointing@v5`.

1 Summary

- **The model** is linear at each frequency: sky pixels, one ground temperature, one offset per receiver regime and, optionally, the Sun's flux density. The columns are integrated through an assumed beam on the real geometry (Section 2). A uniform shift of sky and ground against the offset is an exact null direction.
- **The DEM stops short of the southern horizon.** Extending it 4 km south raises the horizon by up to 5.47° at 87.5 m and changes the predicted antenna temperature by up to 47 K, with the sky and beam fixed (Section 2, Fig. 1). The extended DEM and its horizon are now `derived/dem/v0004` and `horizon_profiles_v0005` (traced from geometry release `v0005`'s antenna), and every fit and product here uses them.
- **Heights.** The mode table rebuilt on 2026-10-18 (memo 001) puts the 07-15 night and the 07-17 sunset on the ground, not at 30 m, and part of the 07-18 night at unknown height. Box-air's lidar and IMU agree wherever they can test it (§4.1). Every fit here uses only rows at 30, 87.5 or 91 m; the ground stretches wait for a horizon profile at 2 m.
- **The ground temperature is not measured by the calibrated data.** They hold one pointing, one height and one regime at night (362 one-minute bins), where $T_{\rm gnd}$ and the offset are degenerate. Only their sum, the pedestal, is determined: 157 to 207 K at 110–170 MHz. A forecast on the full campaign geometry shows that height changes are what would break the degeneracy (Table 1). The nearest soil station shows no daily swing below 50 cm, and puts the soil a metre or more below the site at 288.2–297.9 K, depending on the depth; shallower soil is warmer, and in shale the emission may come from as little as 0.04 m down (§5.2).

- **The data vary about as much as GSM through HFSS predicts**: the template amplitude is $a \approx 1.1$ (0.9–1.7 over the band) on `tcal@v0004`'s kelvin scale; earlier versions found $a \approx 2$ on `tcal@v0003`, whose kelvin scale memo 004 has since revised. The model does not reach the noise: median χ^2 per bin is 94, and 245 on held-out blocks.
- **An independent night fit at the axle tilt the box's IMU measures.** On `fit_v0025`'s inputs, each frequency is fitted with one sky scale θ and one pedestal P ($y = \theta A + P$, no sky field), at the night's measured tilt ($+0.19 \pm 0.6°$), on the bins in which the IMU shows the box still. It gives $\theta = 1.169 \pm 0.012$ (weighted mean over frequency; `fit_v0025`'s a is 1.160 on the same weights); the night fixes θ and the mean level far better than P, the level extrapolated to zero sky. Both depend on the loss prior through the load term. On `tcal@v0004`'s kelvin scale, and if GSM2016 and the beam's normalisation are right, θ needs an antenna-side gain above 1; on the bench's noise-source scale with the switch path it needs 1.04 on average, still above the loss model's 0.47–0.77. The held-out χ^2 per bin is 7–28 below 200 MHz, 5.1 times lower than at the beam fit's $+4.74°$ tilt, and no polynomial trend in time passes the held-out rule (§5.1). A sky-shaped time column (a flat-mirror reflected sky) absorbs much of the residual, with unphysical amplitudes (diagnostic). Memo 005's raster beam v0018 in place of HFSS predicts the night worse (32.75 against 27.59), also where the raster measured the beam (diagnostic).
- **When the Sun is visible.** The ridges stand 39° high at 352 m to the east and 17° at 170 m to the west. The Sun clears the ridge at **15:53 UTC on 07-17** (87.5 m), 3.5 h after it reaches 0° altitude, and sets behind the ridge at **01:21 on 07-18** (91 m). The calibrated data therefore see the direct Sun only from 15:53 to 16:22 on 07-17. At these wavelengths the edge diffracts, so the Sun fades in over 10–30 min.
- **Box-air sees the Sun's radio bursts.** At the times e-CALLISTO spectrographs recorded type III bursts, raw box-air shows emission drifting from high to low frequency within seconds, bright at 45–85 MHz as those stations saw it. 5 of 12 bursts with data pass a test that 5.2% of random daytime times pass ($p = 2.2 \times 10^{-4}$). That identifies solar emission in our data physically (§6.5).
- **The quiet Sun across days.** All 4 terrain crossings of phase C are shown in their raw context; 3 of them, on three days and three heights, can be fitted. They allow a Sun term of 3–41 SFU at 121–219 MHz, the size of the concurrent Sun (RSTN at 245 MHz, steady at 13–17 SFU from 07-15 to 07-18). They do not detect it convincingly, since raw power drifts by as much for other reasons (§6.7).
- **The calibrated 07-17 sunrise excess is about twice the predicted Sun.** After 15:53 the calibrated spectrum rises by a smooth, additive 8–13 K at 148–199 MHz (Fig. 15). That is 2.0 times what concurrent RSTN flux predicts (1.5 times scaled by a), and 0.9–1.2 times the other crossings at 199 MHz. A local fit gives $S_\odot$ from 4 SFU at 109 MHz to 18 SFU at 199 MHz; the same fit at the 0° sunrise, with the Sun behind the ridge, gives -5 to 0 SFU. Something that changes as the box comes into sunlight would follow the same geometry, so the excess is not shown to be the Sun alone.
- **Impulsive events in daytime**: holding the height era or the local hour fixed, the rate is higher with the Sun visible, but at 30 m it shows no step where the Sun crosses the horizon and the longer events go the other way after the re-raise, so whether the events follow the Sun is open (§6.6).
- **Not yet possible**: separating the Sun from the lit ground by timing, or from the sky and ground by the beam's sweep during the raster (§6.8). Below 90 MHz the calibrated result is unusable.
- **The ground from the rotating antenna.** When the antenna moves, the ground fraction swings while the gain stays put, so the ground/GSM ratio of a gain-free regression gives the ground's brightness in GSM kelvin. Three 91 m windows agree on it at 7 frequencies: 189–333 K at 125–180 MHz. In the flagged raster it is 279±505 K at 64 MHz and 322±118 K at 82 MHz, not the thousands of kelvin earlier versions found on the previous mount and right-handed pointing; unflagged it is 1290 and 736 K, and the 30 m window still gives 2810±626 K at 64 MHz (§7). A joint fit across frequency and the sky-calibrated parked nights do not separate the ground, and low-order corrections to GSM are not separable from ground-fixed structure.
- **The low-frequency daily term over phase C.** On rows at confirmed heights, a slow additive term at 62–82 MHz improves the fit with 3-h gain knots over a constant ground by 0.14–0.27 points of held-out rms, more than mocks with the fast gain and no term give. In each half of phase C the real sky beats every shifted sky, but over all of phase C and on 07-16 alone shifted skies do as well, so the term is not shown to be sky. With 1-h knots it goes, as it does in mocks that contain one (§7.5).

- **The ground term with a fast gain.** Over phase C, gain knots every 20 min instead of 3 h lower the held-out rms by 0.2–0.6 points, and change $T_{\rm gnd}$ at 117–180 MHz by a factor 0.5–1.3. With 20-min knots $T_{\rm gnd}$ is 274–579 K over 110–205 MHz, with injection errors of 16–411 K. But the fast gain takes up much of the sky's variation there: the GSM amplitude falls to 0.27–1.48 of the 3-h fit's, and at 74, 125, 148, 172, 195, 211, 219, 234 MHz below 0.15 of it, where no ground term is quoted. The fast gain follows no recorded stream; where the receiver's gain after the RF switch is measured (07-17, 87.5 m) it moves by about 1.1% and does not follow the sky, while the fitted gain moves by up to 25.3%, so it is a gain change before the switch or the fit absorbing model mismatch. A fast gain shared across frequencies cannot be fitted jointly: every high-band training frequency but 141 MHz collapses when it helps fit the shared gain (§7.6).
- **A solar burst at the 07-17 sunset.** Box-air alone shows a broadband excess below 100 MHz during a C1.5 flare (23:51–00:09 UTC), with the antenna on the ground and tilted, not at 30 m. What ends it is open: from the ground there is no traced horizon, so the earlier reading as the ridge occulting the Sun is withdrawn (§6.7).

2 The model

At each frequency, every time bin t is modelled as

$$T(t) = \sum_p A_{\rm sky}[t,p]\,T_{\rm sky}[p] + A_{\rm gnd}[t]\,T_{\rm gnd} + T_{\rm off}[r(t)] + A_\odot[t]\,S_\odot, \tag{1}$$

with the sky in Galactic HEALPix pixels ($N_{\rm side} = 8$), the ground everything below the DEM horizon at one temperature, an offset per receiver regime r, and the Sun's flux density $S_\odot$ in SFU ($10^{-22}\,\mathrm{W\,m^{-2}\,Hz^{-1}}$). The sky is written $T_{\rm sky} = a \cdot \mathrm{GSM} + \delta$. The amplitude a is free; it absorbs the front-end loss (calibrated temperatures are at reference plane P, memo 004) and any GSM scale error. Only δ has a prior, a Gaussian of 10% of GSM.

Integration. The sky and ground columns are integrated on a topocentric grid ($N_{\rm side} = 64$). For each distinct pointing and height era, the beam (HFSS total power here) is rotated by the mount model of the reference beam fit `empirical_raster_v0017` (memo 005), the frame in which its empirical beam was fitted: $R = R_z(\psi)\,R_y(\tau)\,R_x(\mathrm{el} + \mathrm{el}_0)\,R_z(\mathrm{az} + \mathrm{az}_0)$, with $\mathrm{az}_0 = +15.93°$, $\mathrm{el}_0 = +0.05°$, the along-axle tilt $\tau = +4.74°$ and $\psi = 142.2°$ (`eigsep_data`'s `TxGeometryModel`). Pointing is `pointing@v5` (memo 003), whose azimuth is left-handed, as that fit assumes. It is cut at that era's DEM horizon (`horizon_profiles_v0005`, traced from the antenna of geometry release `v0005`, memo 002, on the DEM mosaic `derived/dem/v0004`, which extends v0003 6 km south). The transmitter comes from the same release. The beam fit's mount offsets were fitted on geometry release `v0004`; no fit on `v0005` exists yet (memo 005). Only the map from grid directions to Galactic pixels changes with time. That map is built with `eigsep_sim`'s `rot_gal2top_stack`, whose frame took the vertical from the geocentric radius and north from the J2000 pole, 0.19–0.27° from the true local frame; the fix is pending upstream (`EIGSEP/eigsep_sim` PR #7), and with the team's estimator on `fit_v0009`'s inputs it moves the sky amplitude by a median −0.17% (−7.9% to +4.1% over 44 frequencies) and lowers the median held-out χ^2 per bin from 234.3 to 225.5; the team's next rerun picks it up. Rows are normalized by the full-sphere beam integral, so sky and ground weights sum to one. That makes a uniform shift of sky and ground, against the offset, an exact null direction: breaking it needs a known offset or a known sky. The solver (`eigsep_sim.design_matrix.solve`) reports which columns move along a null direction and marks them unconstrained.

Bearings. The DEM, and so the horizon profiles, are in NAD83(2011)/UTM 12N. Grid north there is 1.52° from true north. The fit rotates the horizon to true bearings; the first trials of 2026-10-01 did not.

Terrain coverage. Earlier versions of this memo traced the horizon only out to the edge of the DEM mosaic `derived/dem/v0003`, which reaches $N = -1000$ m on its southern side. To see what that edge costs, we extended the mosaic 4 km south on the same grid, keeping every original pixel, the origin, the resolution and the vertical datum, and traced the horizon again from the 91 m-era antenna position of geometry release `v0004`, where this comparison was made (the products here use release `v0005`'s, $(E, N, U) = (1655.4, 2031.0, 1777.9)$ m in the DEM frame). The 30, 87.5 and 91 m height labels subtract 61, 3.5 and 0 m from U. Bearings are true

bearings, rotated from grid north by 1.52° as above, in 0.1° steps with 720 sub-samples of the mask. Only the DEM changes between the two traces: sky, beam, site, times, pointings and frequencies are the same. The southern terrain raises the horizon by up to 3.45°, 5.47° and 5.51° at 30, 87.5 and 91 m. At the eight observed night pointings and four frequencies (51–246 MHz), the predicted antenna temperature changes by at most 16, 47 and 48 K (Fig. 1). The 87.5 m case is the selected night (07-17, 04:22–11:23 UTC); the 30 and 91 m cases are synthetic height controls evaluated at those same times, not observations from those eras. These are differences between two model predictions, not measured errors in any calibrated spectrum. A third trace with 8 km of added terrain differs from the 4 km one by at most 4.1×10^{-12} K, at the nominal position and six ±3 mm displacements. Edge flags alone do not test coverage: the original DEM raised none at 30 m and still changed. The tracing is also not shown to be accurate to sub-mK: the finest bearing refinement tested changes the prediction by up to 7.6 mK, so the numbers above are diagnostic and not tied to any convergence bound. This paragraph and Fig. 1 compare that original DEM with the extension; every fit and product in this memo now uses the extended DEM v0004 below.

The extended terrain is now a product. `derived/dem/v0004` is v0003 plus 6 km to the south, on the same grid and keeping every v0003 pixel. Traced on it at the `v0004` antenna (`horizon_profiles_v0004`, for comparison), the horizon rises over v0003 by at most 3.45, 5.47 and 5.51° at 30, 87.5 and 91 m, on 209 bearings at 87.5 m, and is unchanged elsewhere. `curation/horizon_profiles_v0005`, the horizon this memo uses, is traced on the same DEM from the `v0005` antenna; it differs from v0004 by a median of at most 0.16° in any era and by up to 1.9°. Terrain outside v0004 cannot change it: `coverage_certificate.py` follows every bearing beyond the DEM edge on the USGS 1 arc-second DEM, out to the distance (at most 41 km) beyond which terrain below 4200 m would sit under the skyline, and finds none that rises above it, from the profile position at all three heights or from the eight displaced positions of the profile's sensitivity block (`curation/horizon_profiles_v0005_coverage.json`). The coarse terrain is raised by 8.7 m, the most by which it falls below the skylines traced inside the DEM; the smallest margin is then 0.99°, on southern bearings, and 2.62° in the other directions. The check covers this antenna position only.

The Sun column. The Sun is a point source:

$$A_\odot(t) = \frac{\lambda^2}{8\pi k}\, G(\hat{s}(t))\, V(t) \quad \text{(K per SFU)}, \tag{2}$$

for one linear polarization and an unpolarized source. Here G is the beam gain toward the Sun, with the same full-sphere normalization as the rows, and V is its terrain visibility. V is not read off the fixed horizon profile. The DEM is traced along the Sun's own bearing at each time, which gives the edge's elevation and its distance d. The ridges are 150–350 m away, so at these wavelengths the edge diffracts over $\sqrt{\lambda d/2} \approx 2$–6°, tens of minutes of solar motion. V is the Fresnel knife-edge power for the Sun's angle θ above the edge, $v = -\theta\sqrt{2d/\lambda}$: it is 0.25 at $\theta = 0$ and rises to 1 with lit-side ripples. The real ridge is rough and rounded, which smooths these ripples.

The knife-edge shape is an assumption. No measurement here tests it: its frequency-dependent onset, its value at the edge, its ripples. Every fit uses $A_\odot$ only as a template, with one fitted amplitude and, in the timing scans, one time shift. Figure 14 shows it against the data it was fitted to.

3 What the geometry can constrain

Before any data are used, `degeneracy_study.py` builds the design matrix on the campaign's real pointing and heights. It reports Fisher errors on $T_{\rm gnd}$ with radiometer noise in 120 s bins, with beam `empirical_raster_v0017`'s mount and `pointing@v5` (Table 1).

- With the sky free, $T_{\rm gnd}$ is never constrained. That holds even with a known offset: within one pointing and era the ground column is constant and trades against a uniform sky shift.
- With a 10% GSM prior and the offset free, $T_{\rm gnd}$ and the offset are anticorrelated at $\rho = -0.95$ to -1.00. Height changes, which move the horizon, break the degeneracy: all of phase C would give 0.7–19 K.
- The calibrated windows alone are weak unless the offset is known.

The study also simulates and recovers. With HFSS as the true beam the recovered $T_{\rm gnd}$ is unbiased. With the empirical beam as truth and HFSS assumed, it is biased by up to 9 K (12σ) at 199 MHz, while χ^2 per row moves only from 2.9 to 3.0. A good χ^2 cannot vouch for $T_{\rm gnd}$.




Figure 1: Effect of extending the DEM 4 km south, at the nominal antenna position, for the three height labels (rows). The horizon from the original DEM (`derived/dem/v0003`) and the extended DEM, with the affected southern sector enlarged, and the extended-minus-original predicted temperature at four frequencies, with the sky and beam fixed. Colours mark the eight observed night time/pointing pairs by UTC on 07-17; points are the evaluations and lines guide the eye. Within each pair only the DEM changes. The 30 and 91 m rows are synthetic height controls at the same times. The model azimuth is 0.865° for all eight pairs and the model elevations are 0.63–1.33° (`inputs/terrain_coverage/coordinates.csv`). Bearings are true; the horizon step is 0.1°.

slice	58.6	78.1	101.6	148.4	199.2
10% GSM sky prior, offset free					
parked, 30 m only	–	–	–	111.9	158.9
parked, all eras	21.5	10.1	5.3	2.3	0.8
all of phase C	18.6	8.5	4.1	1.8	0.7
tcal windows only	71.5	18.6	6.0	5.4	0.9
10% GSM sky prior, offset known to 1 K					
parked, 30 m only	9.1	4.3	2.4	2.1	1.8
parked, all eras	5.4	2.6	1.6	1.2	0.7
all of phase C	3.8	1.9	1.2	1.0	0.6
tcal windows only	9.9	4.2	1.9	1.8	0.7

Table 1: Forecast $\sigma(T_{\mathrm{gnd}})$ in K at five frequencies (MHz), from the degeneracy study (`summary.json` in `data-analysis`; radiometer noise, 120 s bins). "–": unconstrained. Its "calibrated windows" are `tcal` v0000's spans, the same as v0004's (07-17 04:11–16:22, 07-18 01:27–02:56).

4 Data

Box-air antenna-switch rows are binned in raw power first and calibrated afterwards, as two separate steps. `bin_raw.py` averages, at each HFSS beam frequency, the six channels 3–5 either side of it, midway between the transmitter's teeth. Samples flagged by `flags@v3` (memo 006) on any bit but the advisory bit 9 are dropped, as are box-air's self-EMI lines where memo 006 marks them on. Rows are averaged into bins of one era, regime and pointing, closed after 60 s or when the pointing moves 1° (`raw_binned_v0006`, all of phase C). The fits then calibrate each bin with `tcal@v0004` (memo 004): its scale and offset are evaluated at the bin's mean time for each channel and the calibrated channels averaged. 1147 bins fall where `tcal` covers, 07-17 04:11–16:22 (87.5 m, receiver regime rx-A) and 07-18 01:27–02:56 (91 m, rx-B). Bins with fewer than 10 rows (437) are dropped, which leaves 710 (609 rx-A, 83 rx-B; 73127 rows). Each bin's noise is the standard error of its rows, times the `tcal` scale (σ_k). Frequencies whose in-bin scatter exceeds ten times the radiometer expectation are not fitted, which removes FM and leaves 43.

4.1 Heights

Every fit here takes the antenna's height from the mode table as rebuilt on 2026-10-18 (memo 001, `derived/height_eras/v0001`), by source file, and uses only rows at a height with a horizon profile (30, 87.5 or 91 m). That table moves three stretches that earlier versions of this memo treated as at height. The 07-15 night and 07-16 23:27 to 07-17 01:15 UTC, called 30 m before, are on the ground (about 2 m). Around 07-18 00:00, called 91 m, the antenna was lowered and raised again, and its height is unknown. Of phase C's 17102 raw bins (`raw_binned_v0006`), 2812 are now at 30 m, 3009 on the ground and 1571 of unknown height; with the earlier labels (`raw_binned_v0001`, for comparison only), 6904 were at 30 m.

`validate_heights.py` checks the relabelled stretches against box-air's own recorded lidar range and IMU elevation, without the mode table, in all 3502 box-air files of 07-14 to 07-18 (Fig. 2, Table 2). The lidar returns a distance only in a few stretches; elsewhere every sample reports an error.

- **07-18, lowered and raised again.** From 07-17 23:30 the lidar reads a median of 1.1 m, the box on the ground. Over the next half hour it reads 31 m (28–39), going up, and then nothing within range.
- **The 07-17 sunset, on the ground.** From 07-16 23:27 to 07-17 01:00 the IMU puts the antenna at −66° elevation (43 files), against −2.2° hanging at 30 m on 07-16. That is the box tilted on the ground for tests, as memo 001 records, not hanging; the elevation returns near 0° at 01:00, when the raise begins.
- **Resting against hanging.** Hanging at 30, 87.5 or 91 m, the elevation moves within every file: 0% of those files have a spread under 0.1°, and the file medians take 99–166 distinct values. Before the 07-15 lift and on the afternoon of 07-15, 25–29% of files are that still, and the median stays at −3.2° (8 values in 1008 files): resting, not hanging. The 07-15 night itself has IMU data in only 5 of its 470 files and no lidar, so it is bracketed by these stretches, not tested.

The labels hold where these streams can test them. There is no horizon profile at 2 m, so the ground stretches cannot be modelled: that profile (`horizon_profiles_v0005` in memo 002's numbering) is the missing input.

Until it exists, the 07-15 night, the 07-17 sunset and the ground half of the 07-18 night are left out of every fit here.




Figure 2: Box-air's height from its own sensors, 07-14 to 07-18, one point per raw file (`validate_heights.npz`), against the 2026-10-18 mode table (shading). Top: the IMU's median elevation in each file (readings near ±180°, IMU wraps, fall off the axis). Middle: its spread within the file (max − min), log scale. Bottom: the lidar's median distance (points) and 10th–90th percentile (lines), log scale, where it returned any. Hanging at height the elevation always moves; on the ground before the lift it often does not, and at the 07-17 sunset it sits near −66°. The lidar sees the 07-18 lowering and re-raise.

5 The night fit

`fit_v0025` fits Eq. 1 without the Sun, to the 362 bins with the Sun below −10° (Fig. 3, Table 3). All of them are rx-A, at one pointing and one height.

- $T_{\rm gnd}$ **is not measured.** It is exactly degenerate with the offset, as Section 3 predicts. At 25 frequencies the ±1° elevation wobble formally separates them. The values there run from −6733 to 9536 K: fitted model error, not a temperature. The pedestal $\langle f_{\rm gnd}\rangle T_{\rm gnd} + T_{\rm off}$ is determined: 157 to 207 K at 110–170 MHz.
- **The sky amplitude is** $a \approx 1.1$, measured to ±0.02 at each frequency, so on `tcal@v0004`'s kelvin scale the sky's variation is close to GSM's through HFSS. It ranges from 0.9 to 1.7 across the band, far more than its error. How much of that is the kelvin scale, the beam's sky fraction or GSM, the fit cannot tell.

window (UTC)	mode table	lidar median (m)	IMU el. (deg)	IMU spread (deg)	files
07-14 00:00-07-15 03:30	∼2m	0.2 (0.2–0.2)	−3.2	0.25	1363
07-15 03:30-14:00, the 07-15 night	∼2m	–	−3.2	2.00	470
07-15 17:00-21:28	∼2m	–	−3.0	0.24	124
07-15 23:00-07-16 16:30	∼30m	–	−2.2	0.48	482
07-16 23:27-07-17 01:00, the 07-17 sunset	∼2m	–	−66.2	3.63	44
07-17 01:00-02:30, the raise	raising	–	0.4	2.91	42
07-17 04:00-16:00	∼87.5m	–	−0.0	0.46	336
07-17 16:00-21:28	∼91m	–	−0.2	0.92	153
07-17 23:30-07-18 00:00	unknown	1.1 (0.7–1.6)	−−	–	14
07-18 00:00-00:30	unknown	30.6 (27.8–39.4)	−−	–	14
07-18 00:30-03:10	∼91m-reraise	250.0 (250.0–250.0)	−176.0	0.71	71

Table 2: Box-air's lidar and IMU per window (`validate_heights.json`): the mode-table label, the lidar's median distance (10th–90th percentile, medians over files), the IMU's median elevation and the median within-file spread, and the number of raw files.

- **The model is not adequate at the noise level**: median χ^2 per bin is 94, and 245 on held-out alternating 30-min blocks.
- **The morning behind the ridge looks like night.** Adding the bins from 0° sunrise to the ridge crossing, `fit_v0026` (573 bins), raises the median χ^2 only from 94 to 125.

product	bins fitted	bins	freqs	median χ^2	median held-out χ^2
`fit_v0025`	night (Sun below −10°)	362 / 710	43	94	245
`fit_v0026`	Sun behind the ridge allowed	573 / 710	43	125	315
`fit_v0027`	Sun modelled, all bins	710 / 710	43	412	1490

Table 3: The fit products. Every fit evaluates the model on all bins; "bins fitted" are those used to solve. `fit_v0027` fits the Sun column on every bin, including those with the transmitter on; its held-out χ^2 reaches 1×10^5, and it is not used.




Figure 3: The night fit (`fit_v0025`). From the top: GSM amplitude; the rx-A pedestal; T_{gnd} where the solver calls it constrained (symlog axis; the dashed line is 300 K); χ^2 per bin on training and held-out blocks.

5.1 An independent night fit at the measured tilt

Draft, 2026-11-04; label: diagnostic.

This section refits `fit_v0025`'s night (07-17 04:12–11:24 UTC) at its 43 frequencies with an independent pipeline (the night fit of Christian Hellum Bye's EIGSEP D5 analysis, not shared; Appendix A). The inputs are `fit_v0025`'s: its calibrated vector (`raw_binned_v0006` on `tcal@v0004`), its 362 bins, its train/test split, beam `empirical_raster_v0017`'s az_0, el_0 and ψ, `pointing@v5` and `horizon_profiles_v0005`. One input is replaced: the axle tilt τ is the night's own, measured by box-air's IMU (below). One selection is added: the 44 bins in which the IMU shows the box moving are cut, which leaves 318 (below). Every other difference is one of method:

- the sky and ground columns are integrated harmonically (spherical-harmonic transforms) on the HFSS export on its own 1° grid, with the horizon as a fractional open-sky weight, instead of HEALPix pixel sums with a binary horizon; the local frame is tied to Galactic coordinates through geodetic coordinates;
- one sky scale and no δ field; the CMB enters as a term of its own;
- the weights include a multiplicative white term (below), and a load term from memo 004 is tested;
- the fit is judged by its held-out prediction, and a generic time trend is tested beside it.

The model. At the antenna terminals, bin t at one frequency sees

$$T_{\mathrm{ant}}(t) = S(t) + T_{\mathrm{CMB}}\,[1 - f(t)] + f(t)\,T_{\mathrm{gnd}}, \tag{3}$$

where S is the GSM2016 sky (CMB removed) weighted by the beam above the horizon, f is the beam's fraction below it, $T_{\mathrm{CMB}} = 2.725\,\mathrm{K}$ and T_{gnd} is one ground temperature. The passive path to plane P (balun, coax and switch path; available gain $G \le 1$) adds T_{add}, $T_P = G\,T_{\mathrm{ant}} + T_{\mathrm{add}}$. If the noise source's true excess differs from the one `tcal@v0004` assumes by a factor $1/k$, memo 004's calibration gives $y = k\,T_P + (1-k)\,c_L + T_{\mathrm{off}} + n$, with c_L the reference load's mismatch-weighted temperature, T_{off} an additive offset and n the noise. The fit has one coefficient per group the data can identify:

$$y(t) = \theta\,A(t) + P + n(t), \qquad A = S + T_{\mathrm{CMB}}(1-f), \qquad \theta = kG\gamma, \quad P = kG\langle f\rangle T_{\mathrm{gnd}} + k\,\bar{T}_{\mathrm{add}} + (1-k)\,\bar{c}_L + T_{\mathrm{off}}, \tag{4}$$

with γ the product of GSM2016's scale and the beam's sky normalisation, 1 if both are right. At one pointing f changes by at most 0.0017 over the night, so T_{gnd} and the offsets enter only through P (Section 3): no ground prior, no offset per regime, no sky freedom and no time-dependent parameter. Two time-dependent terms are left out: $f(t)$'s change, and $T_{\mathrm{add}}(t)$, whose outdoor part follows the air ($-5.3\,\mathrm{K}$ over the night, §5.2; the night-residual analysis's measured column for it changes by 0.8–2 K peak to peak across the band and is tested in Table 5). θ is the counterpart of a and P of the pedestal. The physical split comes after the fit: $k = \theta/(G\gamma)$ with a prior on the losses. The load term follows from the calibration equation: its time-varying part, $(1-k)\,[c_L(t) - \bar{c}_L]$, is a fixed column with $k = \theta/\bar{G}$, $\bar{G}$ the loss prior's median G at each frequency, iterated with the fit ($1-k$ from -1.10 to -0.46). θ and P therefore depend on the loss prior through this column. It is kept because it moves θ or P by more than σ_J (below) at 4 frequencies (at most $1.12\,\sigma_J$ in θ, $1.17\,\sigma_J$ in P), that is, it changes a reported result by more than its error, the analysis's criterion for keeping a term; it raises the held-out χ^2 per bin from 27.17 to 27.50 on the same test bins. P's parts are named but for one: $T_{\mathrm{off}} = k\,D + T_{\mathrm{rest}}$, where $k\,D$ is the difference between the receiver's noise waves on the antenna and on the load, and T_{rest} is whatever additive term is left after the named ones and $k\,D$. T_{rest} is unidentified, so no value of T_{gnd} or T_{off} is drawn from P here.

Noise, errors and scores. Each bin's error is $\sigma_{\mathrm{tot}}^2 = \sigma_k^2 + (\epsilon_w y)^2$: `fit_v0025`'s σ_k plus a white multiplicative excess, measured by maximum likelihood on triples of consecutive bins, $\epsilon_w = 0.10$–0.34% (weighted mean 0.130%). The night's raw integrations show what it is (the night-residual analysis; Fig. 10): single channels start at 1.1–1.2 times the radiometer level at 0.54 s, reach their minimum, 0.12–0.15%, at 9–17 s, and rise to 0.19–0.20% at 60 s, 7–8 times the radiometer level; the band's common mode rises to 0.11–0.14% at 60 s, the size of ϵ_w, and it rises with the antenna switched out too (the load, RFAMB: 0.025 to 0.042% from 1 to 17 s). The excess is a red part, common to the band and at least partly behind the antenna, not radiometer noise. The errors σ_J on θ and P are delete-one-block jackknife errors over the 15 blocks of 30 min. A mean over frequency (over the band, or over a part of it) takes its error from the same replicates: each replicate deletes the same

30 min at every frequency, the mean over frequency is formed for each replicate, and the jackknife spread of those means is the error. The frequencies' errors are strongly correlated, and treating them as independent would give θ's mean an error of 0.003 instead of 0.012. `fit_v0025`'s a and pedestal have no replicates, so means that involve them (a, θ/a, P less the pedestal, the method chain) are quoted without an error. The held-out χ^2 is per bin, as in Table 3: the fit on the 166 training bins scored on the 152 test bins, on $\sigma_{\rm tot}$. The quoted fit includes the load term (above). The generic-trend test starts from the same model with a ground column under the Gaussian prior $T_{\rm gnd} = 293 \pm 30$ K of §5.2 (R1g), adds polynomials in time of order 1–3 (R1P1–R1P3) or a linear drift in time of both the sky scale and the level (R1D), and adopts one only if it lowers the held-out χ^2, summed over frequency, in every held-out block or by more than 2.33 jackknife errors. At the IMU's tilt it adopts R1g: the linear trend wins 5/7 held-out blocks, a gain of 0.46 jackknife errors. With the beam fit's tilt it adopts R1D, and the linear trend alone wins 7/7 held-out blocks (2.44 errors). So no trend in time passes the rule at the measured tilt, and a drift does at the beam fit's.

The tilt. `fit_v0025` rotates the beam with the beam fit's $\tau = +4.74°$, which was fitted on the 07-17 raster in the `~91m` era. Box-air's IMU measures the axle's tilt in every era (memo 003, Sec. 7): over these bins of the `~87.5m` era it is $+0.19 \pm 0.6°$, the error mostly the IMU's slow drift, and the bins scatter about it by 0.54°. That is the IMU's value from the readings before the firmware calibration's accelerometer bias was removed; with it removed (memo 003, Sec. 7) the night's mean is +0.18°, 0.015° lower, far inside the quoted error, and the fit moves to it at its next rerun. Memo 003 shows the tilt per era and scans the night's held-out χ^2 against a constant τ (its Fig. 10); memo 005, Sec. 5.6 (the axle tilt), says what the measured tilt makes of the beam fit's. Like for like, the plain fit's (without the load term) held-out χ^2 is 139.8 at $\tau = +4.74°$, 27.2 at the IMU's τ and 27.4 at $\tau = 0$ (means over frequency; Fig. 6). The IMU's tilt predicts better than the beam fit's at 35 of 43 frequencies; the beam fit's is better only at the 8 frequencies from 203.12 to 234.38 MHz. The IMU's tilt and $\tau = 0$ differ by +0.19°, well inside the tilt's error, and are not separated (the IMU's is better at 28 frequencies). With the beam fit's tilt the model's Galaxy transit is mistimed, and the residual drifts across the night at every frequency (Figs. 4 and 5): its rms is 12.1 $\sigma_{\rm tot}$ averaged over frequency (worst 18.9), against 4.6 $\sigma_{\rm tot}$ (worst 12.4) at the IMU's tilt. The fit below uses the IMU's tilt. On the bins it keeps (below), the plain fit's held-out χ^2, scanned over a constant τ, is lowest at $+0.19 \pm 0.25°$ (the vertex of a parabola through the scan's lowest point and its two neighbours, with memo 003's jackknife error), at the IMU's tilt; on all 362 bins it is lowest at $0.00 \pm 0.41°$. The selection that keeps those bins is defined on the IMU, so the scan does not confirm the IMU's tilt independently. Pointing table v6 (memo 003, Sec. 7.1), released with this version, carries the tilt per sample; on the same bins it agrees with the IMU means used here (memo 003, Sec. 7), and the residual is the same with the per-bin tilt (memo 003), so the fit holds τ at the night's mean.

The selection. With its motors still, the box moves under wind load on its ropes. A moving box's accelerometer reading is not a static angle, and its beam is not where the model puts it. The fit therefore drops every bin in a 5-min UTC window in which the IMU's per-sample tilt (pointing table v6, memo 003, Sec. 7) has a standard deviation above 0.2°, or any sample is flagged as not quasi-static (flag bit 14). The rule is defined on the instrument, not on the fit or the wind, and was fixed before it was scored. It picks 7 windows (starting 04:10, 04:15, 08:15, 08:35, 08:40, 09:25, 10:35 UTC; standard deviations 0.24–0.48°, none flagged). They remove 44 of the 362 bins (39 training, 5 test), and every jackknife block keeps bins. Between the windows τ is held at the night's value. At the Tule Valley RAWS (`TVWU1`, 11.5 km north) and the SCAN station (`TULU1`, 5.0 km WSW on the valley floor; Table 6), the hourly wind falls from 19–22 mph at 01–02 UTC to at most 5.5 mph at 05–08 UTC (the site-weather analysis, §5.2). Its direction turns from S–SW to E–SE: between the two regimes' vector means, from 208° (02:36–04:36 UTC) to 120° (06:36–10:36) at the RAWS and from 186° to 87° at SCAN, with light wind from other directions between. In the strong wind before the fit's first bin (04:12 UTC), the same rule marks 28 windows from 01:00 to 04:05 disturbed, 21 of them flagged as not quasi-static. The cut windows at 04:10 and 04:15 UTC coincide with the end of the S–SW wind. Those from 08:15 to 10:35 fall in light E–SE wind: at the RAWS 4–9 mph, gusts 11–17 mph; at SCAN 2.8–5.1 mph in their hours, no more than the 3.3–5.5 mph of the calm hours. Hourly values 5.0–11.5 km away cannot resolve 5-min windows, so what moved the box then is open. On the test bins it leaves, the cut lowers the held-out χ^2 per bin from 30.02 to 27.50 (mean over frequency; lower at 34 of 43 frequencies, and worst at 70.31 MHz, +3.02). A threshold of 0.15° cuts 66 bins and scores 27.81; one of 0.30° cuts 28 and scores 28.44.

From `fit_v0025` to this fit. The night fit first reproduces `fit_v0025`'s sky amplitude, a_0, with its method, and then makes one change at a time, keeping each (Table 4). The steps interact, so the change at each step depends on the order they were run in: the table records one path, not each change's own effect. With every change made, the amplitude is 1.004 of a_0 (weighted mean over frequency). `fit_v0025` carries the beam fit's $+4.74°$ tilt, so the difference between its a and θ (black and blue in Fig. 7) mixes method and tilt.

Table 4: The method chain from `fit_v0025` to this fit (the night fit). Step 0 reproduces `fit_v0025`'s fit with its method and inputs; each later step makes one change and keeps the changes above it (steps in the order run; labels are the night fit's). θ/a_0: the sky amplitude over step 0's, weighted mean over frequency (no error: the steps have no jackknife replicates), and its range over the 43 frequencies. Steps 0–8 fit all 362 bins; step 9 cuts the 44 bins in which the IMU shows the box moving and is the headline fit of this section.

step	change (each added to the one above)	θ/a_0	range
0	`fit_v0025` reproduced (its method, $\tau = +4.74°$)	1	–
0t	tilt: $\tau = +4.74° \to +0.19°$ (the IMU's)	0.964	0.91–1.04
1	frame: the team's local frame → geodetic	0.955	0.84–1.08
2	beam: HEALPix → the HFSS export's 1° grid (harmonics)	0.954	0.84–1.08
5	δ sky field off	1.000	0.66–1.18
5b	free ground column off (P alone)	1.001	0.68–1.19
3	sky pixelisation: $N_{\rm side}$ $8 \to 64$	0.993	0.64–1.17
4	integration: pixel sums, binary horizon → harmonic, fractional horizon	0.992	0.64–1.17
6	CMB term $T_{\rm CMB}(1-f)$ on	0.992	0.64–1.17
7	weights: $\sigma_k \to \sigma_{\rm tot}$ (the plain fit)	0.980	0.70–1.14
8	load term as a fixed column	0.986	0.76–1.19
9	selection: the bins where the IMU shows the box moving cut (the headline fit)	1.004	0.76–1.25

Result. At the IMU's tilt (Fig. 7), θ has a weighted mean of 1.169 ± 0.012 and runs from 0.83 to 1.44; its jackknife error averages 3.3% (worst 17% at 226.56 MHz). Its shape across the band is a result: it rises to 1.35 at 82.0 MHz below the FM band, falls to 1.06 at 160.2 MHz and rises again to 1.44 at 210.9 MHz, well beyond its errors. `fit_v0025`'s a has a weighted mean of 1.183 on its own errors and 1.160 on θ's; θ/a has one of 0.999 on its propagated errors and 1.016 on θ's (range 0.76–1.25). Like P, θ depends on the loss prior through the load term. P has a weighted mean of 145.8 ± 3.6 K and runs from -32 to 210 K; on average it lies -18.8 K from `fit_v0025`'s rx-A pedestal (163.7 K), and furthest at 50.78 MHz ($+516$ K). Refitting on columns with $\mathrm{az}_0 \pm 0.7°$, $\mathrm{el}_0 \pm 0.93°$, the horizon raised or lowered by 0.26°, and $\tau \pm 0.5°$, each at the IMU's tilt, gives a systematic band of half-width 1.31% on θ (weighted mean; worst 2.32% at 164.06 MHz) and 3.85 K on P (worst 22.1 K at 50.78 MHz). The tilt steps change θ by at most 1.5% (at 203.1 MHz), inside the mount steps' envelope at all but 2 frequencies, where they widen θ's band by at most 0.0013% (at 203.1 MHz); they widen P's only at 199.2 MHz, by 0.16 K. The band's half-width exceeds σ_J at 14 of the 43 frequencies in θ (at most $1.72\,\sigma_J$, at 160.16 MHz) and at 10 in P (at most $1.41\,\sigma_J$, at 171.88 MHz), so there the mount matters as much as the noise. These steps are not measured mount errors: neither v0017 nor v0018 (memo 005) states an error for az_0 or el_0, and v0018 holds the tilt at the IMU's value, with that value's error. At the beam fit's tilt, θ would be $+2.4$% higher on average, and up to $+13$% at 203.12 MHz; that fit is not shown, since the IMU's tilt rules it out for this night.

What the night measures. Over the night A changes by a small fraction of its mean, so the jackknife correlation of θ and P is -0.9979 to -0.9936 across the band. This is not a degeneracy of the data. P is the level extrapolated to zero sky, $A = 0$, far from the night's values of A. The night measures θ, and the mean level $M = \theta\langle A\rangle + P$, with $\langle A\rangle$ the $\sigma_{\rm tot}$-weighted night mean of A, far better than P: $\sigma_J(M)$ is 0.2–3.0 K across the band, against 3–37 K for P. $P = M - \theta\langle A\rangle$ follows, and an error in θ moves it by $-\langle A\rangle$ times as much. While the residual is structured (below), θ and M are not independent: their jackknife correlation runs from -0.72 to $+0.61$ across the band. The degeneracies that remain are inside the groups: θ holds the kelvin scale, the losses, GSM's scale and the beam's sky fraction, and P the parts of Eq. 4. Both are developed in memo 009.

The kelvin scale. Everything here is on `tcal@v0004`'s kelvin scale, whose noise-source temperature is `noise_source@v0001`'s. The bench data behind it, reduced with the hot load's path through the RF switch (port RFAMB, as the bench was wired) included (the bench analysis of the EIGSEP D5 analysis), put the noise source's excess lower than `tcal@v0004` has it. On that bench scale every $T - T_L$ here, with T_L the reference load's temperature, scales by $r(\nu)$: 0.876 / 0.877 / 0.858 / 0.863 on average over 50–88 / 108–150 / 150–200 / 200–235 MHz, and 0.85–0.88 at the 34 frequencies where both scales exist (Fig. 7c, dashed, shows $1/r$). θ scales with it. This section's frame, the same as memo 009's, is therefore: on `tcal@v0004`; with the bench switch-path correction, $T - T_L$ scales by $r(\nu)$. That is a stated rescaling for a later `noise_source` release with the switch path, not a correction applied here.

In the terms of the loss model, $\theta = kG\gamma$. As `tcal@v0004` stands, $k = 1$, and if GSM2016's scale and the beam's normalisation are right ($\gamma = 1$) the fit needs $G = \theta \approx 1.169$: above 1 at 42 of 43 frequencies, which a passive path cannot give. On the bench scale, $k = 1/r$ and, with $\gamma = 1$, $G = r\,\theta$ is 1.04 on average (0.91–1.23), still above the loss prior's median G of 0.47–0.77 at every frequency. Taking G from the loss prior instead gives $k = \theta/G$ a weighted mean of 1.62 ± 0.03 (1.46–2.10, largest at 58.59 MHz), against 1.61 for `fit_v0025`'s a/G and the bench's $1/r$ of 1.15. If GSM2016, the HFSS beam's sky weighting and the passive front end are right, the noise source's excess in the field is smaller than either the bench or `tcal@v0004` says. k rests on the loss prior and is not measured here; memo 009 analyses the group.

What the model leaves. The model is still far from the noise: at the IMU's tilt the held-out χ^2 per bin is 7–28 below 200 MHz, and 32 to 165 above (Fig. 6). The residual has three features:

- a slow wave in time, common to the band from 100 MHz up and strongest above about 190 MHz (Fig. 4, bottom);
- an excess early in the night: over the 9 fitted bins before 04:30 UTC (the selection removes the earlier ones) the residual averages +1 to $+8\,\sigma_{\rm tot}$ at the three frequencies of Fig. 5, and reaches +5 to $+13\,\sigma_{\rm tot}$. The box settling after the raise, the noise source warming up and twilight are candidates;
- extra scatter at low frequency: from bin to bin the residual scatters by $1.9\,\sigma_{\rm tot}$ at 70.31 MHz, against 1.3 and 1.3 at 148.44 and 199.22 MHz (1 for white noise at $\sigma_{\rm tot}$).

The night-residual analysis of the EIGSEP D5 analysis adds candidate terms for them one at a time, each against a control with the same freedom, and scores them like the headline fit (Table 5; diagnostic). The flat-mirror reflected sky lowers the held-out χ^2 per bin to 26.30 with no free parameter (better at 22 of 43 frequencies). With one free amplitude α per frequency it falls to 13.93 (at 207–234 MHz, 24.7 against the headline fit's 106.9), against 23.40 for the same column reversed in time; but α runs from −646 to +88, where 1 is the Fresnel value, and θ and P move by up to 110 and $105\,\sigma_J$. A sky-shaped time column absorbs much of the residual, but the flat mirror is not the physics; a reflected-sky model on the terrain's echo geometry is in preparation. Even with $\alpha = 1$, θ and P move by up to $29\,\sigma_J$, so a reflected sky would matter for both. The measured $T_{\rm add}(t)$ column and the noise source's off-path term each beat their controls (25.01 against 25.54; 25.14 against 25.72).

The raster beam on the night. The night-residual analysis also puts memo 005's raster beam `empirical_raster_v0018` in place of the HFSS beam, at the IMU's tilt, both at $\ell \leq 64$ and on v0018's mount ($\mathrm{az}_0 = 15.970°$, $\mathrm{el}_0 = 0.094°$; Figs. 8 and 9). On held-out data the night prefers HFSS: the held-out χ^2 per bin, averaged over the 43 frequencies, is 27.59 with the HFSS beam (27.50 for the headline fit, which differs from it in ℓ cutoff and in mount, $\mathrm{az}_0 = 15.934°$, $\mathrm{el}_0 = 0.045°$) and 32.75 with v0018, which is better, by more than 0.05, at 5 frequencies. Two checks were declared before the run; their subsets share 24 frequencies. At the 24 frequencies on a tooth the beam fit used, v0018 loses by +3.52 on average (better at 3); at the 25 where both of the transmitter's polarizations were fitted (132.8–230.5 MHz), by +4.28 (better at 3). On the 19 frequencies where the raster beam is interpolated it loses by +7.23. The night therefore loses where the raster measured the beam too, though by less. v0018 does better at the top of the band, 218.8–234.4 MHz (by up to −47.1), and elsewhere only marginally, at 109.4 MHz (by −0.4). Splits chosen after the run locate the loss: with v0018's difference from HFSS kept only above 30° body elevation the held-out χ^2 is 33.67, and kept only below, 24.83, better than HFSS; kept only in the directions the raster sampled, 28.22, and only in the others, 29.05. At the three frequencies of Fig. 5 (Fig. 9), v0018's residual at 70.31 MHz swings from positive to negative across the peak of the night's sky, more than the HFSS beam's (held-out χ^2 42.2 against 26.2); at 148.44 and 199.22 MHz the two residuals follow the same slow wave (16.0 against 8.3, 19.3 against 15.3). Below 132.8 and above 230.5 MHz the raster measures one polarization projection of the field in each direction, and the other

Table 5: The night-residual analysis's candidate terms for the residual (§5.1), each added alone to the headline fit, with the control that has the same freedom (indented). Free: free parameters per frequency. Held-out χ^2 per bin, mean over the 43 frequencies, on the headline fit's bins and split. Better at: frequencies where it is lower than the headline fit's by more than 0.05. The last two columns: the largest shift of θ and of P from the headline fit over frequency, in its jackknife errors. Flat mirror: GSM2016 mirrored about the local horizontal, seen through the beam below the horizon, times a normal-incidence Fresnel reflectivity (0.20–0.21, $\varepsilon = 7$), with the ground's emission $(1 - R)\,T_{\mathrm{gnd}}$ at $T_{\mathrm{gnd}} = 293\,\mathrm{K}$; α scales it. $T_{\mathrm{add}}(t)$: the loss emission from the logged switch temperature and the site's outdoor air (§5.2) through the loss prior. $k(t)$'s off-path term: the noise-source board's temperature against the load's. Diagnostic.

term added to Eq. 4	free	held-out χ^2	better at	$\max \lvert\Delta\theta\rvert/\sigma_J$	$\max \lvert\Delta P\rvert/\sigma_J$
none: the headline fit	–	27.50	–	–	–
reflected sky, flat mirror, $\alpha = 1$	0	26.30	22	29.1	30.0
reflected sky, flat mirror, α free	1	13.93	25	110.1	105.3
control: the mirror column reversed in time	1	23.40	29	17.4	17.2
$T_{\mathrm{add}}(t)$, measured column	1	25.01	30	2.0	2.1
control: a linear ramp in time	1	25.54	26	1.7	1.8
$k(t)$, the noise source's off-path term	1	25.14	30	1.5	1.4
control: a linear $k(t)$	1	25.72	25	1.4	1.4

comes from the fit's prior toward HFSS (memo 005, Sec. 2). The comparison is diagnostic: the raster beam does not yet predict the night as well as HFSS, so HFSS stays this fit's beam. Understanding the difference needs the raster and the night to constrain the beam together, which is future work.




Figure 4: At the IMU's tilt the residual's rms falls from 12.1 to 4.6 σ_{tot} (means over frequency), and what remains is structured in time. The independent night fit (§5.1): `fit_v0025`'s calibrated vector (its 362 bins in time order down the page; 43 frequencies on a 3.9 MHz channel axis; unfitted channels, cells `fit_v0025` masks and the 44 bins the selection cuts blank), the model $\theta A + P$ with the load term, and the residual over σ_{tot}. Top row: the IMU's $\tau = +0.19°$; middle row: the beam fit's $\tau = +4.74°$, each fitted on all 318 bins the selection keeps. Data and model are drawn as T over the data's σ_{tot}-weighted night mean at each frequency, minus one, on one scale; the mean includes the pedestal P, so the fraction is smaller than the sky's own variation. Both residuals over σ_{tot} share one symmetric scale (±25); for scale, $1\,\sigma_{\mathrm{tot}} \approx \epsilon_w T$, 0.10–0.19% of T (0.32–0.34% at 101.6 and 105.5 MHz, inside the FM band). Bottom row: the top row's residual in kelvin, on a linear ±25 K scale; 68 of the 13430 unmasked cells (0.5%) lie beyond it, 68 of them below 108 MHz. Diagnostic.




Figure 5: With the beam fit's tilt the model mistimes the Galaxy's transit and the residual drifts across the night at all three frequencies. Fig. 4 at 70.31, 148.44 and 199.22 MHz. Top: T [K] in every panel, each on its own range: the data with $\pm\sigma_{\rm tot}$ (grey) and the model at the IMU's tilt (black) and at the beam fit's (orange). Bottom: the residuals over $\sigma_{\rm tot}$; grey band: ±1. The excess before 04:30 UTC and the extra scatter at 70.31 MHz are discussed in §5.1.




Figure 6: The misfit floor: held-out χ^2 per bin of the plain fit (no load term), trained on `fit_v0025`'s training blocks and scored on its test blocks, against frequency, at the beam fit's tilt and at the IMU's; legend: mean over frequency and the worst frequency. At the IMU's tilt the model misses the held-out bins by 7–28 times the noise variance below 200 MHz, rising to 165 at 226.56 MHz. The fit with the load term scores 27.5 on average at the IMU's tilt. $\tau = 0$ (mean 27.4) is not drawn; the tilt scan is memo 003, Fig. 10.




Figure 7: θ exceeds 1 at 42 of 43 frequencies, and the mount steps' band exceeds its jackknife error at 14; the k it implies over the loss prior lies above the bench's $1/r$ everywhere. The parameters of Eq. 4 against frequency, at the IMU's tilt. (a) θ, the multiplicative sky scale, and (b) P, the additive level, with jackknife errors; shaded: the envelope of the $\mathrm{az}_0 \pm 0.7°$, $\mathrm{el}_0 \pm 0.93°$, horizon $\pm 0.26°$ and $\tau \pm 0.5°$ refits. Black: the night fit of §5 (`fit_v0025`), its a and rx-A pedestal with their errors, at the beam fit's $+4.74°$ tilt (Table 4); pedestal values beyond the panel (-548 K at 50.78, 283 K at 58.59, 276 K at 62.50 MHz) are drawn as triangles at its edge. Red in (a): the θ the night would give on the bench's kelvin scale if GSM2016's scale and the beam's sky normalisation were right, $k_{\mathrm{bench}}G$ ($\gamma = 1$ in Eq. 4; 0.58–0.89), with the loss prior's 16–84% half-width and the bench k's error in quadrature; dotted, G alone (0.47–0.77), the θ if `tcal@v0004` were right ($k = 1$). (c) $k = \theta/G$ over the loss prior's draws, with θ drawn from its jackknife error (median and 16–84%). The prior is $G = G_{\mathrm{sw}}G_{\mathrm{cb}}$: the switch path measured, the balun from the ADT1.5-1+ datasheet's typical loss plus 0–0.2 dB for the balun board and the transformer's unit spread, and the coax 1.32 m of RG-401 at its datasheet loss (team memo 009 gives the evidence). Dotted: $k = 1$, `tcal@v0004` as it stands; dashed: the bench noise-source scale with the switch path, $1/r(\nu)$ (§5.1, the kelvin scale), with its total fractional error (green, 11% on average). Not drawn: the jackknife correlation of θ and P, between -0.9979 and -0.9936 at every frequency (memo 009 develops it). Diagnostic; (a) and (b) depend on the loss prior through the load term, (c) and (a)'s red curves on it directly. Drawn by this memo's `nightfit.py` from the night fit's `results.json`.




Figure 8: HFSS predicts the night better than memo 005's raster beam v0018 at most frequencies, also on the teeth the beam fit used; v0018 does better only above 200 MHz. Top: held-out χ^2 per bin against frequency of the headline fit's model (318 bins, `tcal@v0004`, `fit_v0025`'s split, the IMU's tilt) with the HFSS beam (black circles) and with v0018 (orange squares), both at $\ell \leq 64$ on v0018's mount (the night-residual analysis); legend: mean over frequency. Bottom: their difference, v0018 minus HFSS (above zero: HFSS better). Filled: the night frequency lies within 0.1 MHz of a tooth the beam fit used (64 teeth, 39 on arm 1 and 25 on arm 0); open: interpolated, lettered along the top by the nearest tooth's exclusion (I isolation, F FM, C coherence). Shaded: both arms fitted (132.8–230.5 MHz). Every night frequency is an arm-0 channel, so below 132.8 MHz each sits on an excluded arm-0 tooth between fitted arm-1 teeth. Below 64.5 MHz (dotted) the raster beam is the fit's HFSS prior ($\Delta \approx 0$). Drawn by this memo's `nightfit.py` from the night-residual analysis's `results.json`. Diagnostic.




Figure 9: With v0018 in place of the HFSS beam the residual at 70.31 MHz swings across the peak of the night's sky; at 148.44 and 199.22 MHz the two beams leave the same slow wave. The two fits of Fig. 8 at the three frequencies of Fig. 5, against UTC on 07-17. Top: the data with $\pm\sigma_{\mathrm{tot}}$ (grey) and the model $\theta A + P$ with the load term for the HFSS beam (black) and for v0018 (orange), each fitted on the 166 training bins. Bottom: data minus model over σ_{tot} (HFSS circles, v0018 squares), filled on the 152 held-out bins, whose blocks are shaded, and open on the training bins; grey band: ± 1. The bins the selection cuts are blank. Held-out χ^2 per bin, HFSS and v0018: 26.2 and 42.2 at 70.31 MHz, 8.3 and 16.0 at 148.44 MHz, 15.3 and 19.3 at 199.22 MHz. Drawn by this memo's `nightfit.py` from the night-residual analysis's fits. Diagnostic.




Figure 10: The multiplicative excess is red noise common to the band: single channels follow the radiometer line only over the first seconds. The noise against integration time (§5.1): overlapping Allan deviation of the fractional power of the 07-17 night's raw box-air integrations (0.54 s each; flags not applied, outlying channels screened; the night-residual analysis). The fractional power is taken about a quadratic in time fitted per RFANT dwell (a linear one per dwell, and per 60-s piece of RFANT, in the fourth panel), which biases the longest τ low. Three panels, the antenna (RFANT) in 3 bands: single raw channels (rms over the band's channels; shaded, the 16–84% range of the channels' Allan deviations), the band's common mode (its mean), and the radiometer level $1/\sqrt{\Delta\nu\,\tau}$ for one channel and for the band's channels (dashed); red: `fit_v0025`'s 60-s bin (vertical line) and the night fit's ϵ_w, rms over frequency (dotted, 0.149%). Fourth panel: common mode and single channels in the three switch states on 60-s dwells. Single channels' rise beyond 9–17 s is the red part that ϵ_w takes up in the 60-s bins. Drawn by this memo's `nightfit.py` from the night-residual analysis's `results.json`. Diagnostic.

5.2 Site weather and ground temperature

Draft, 2026-11-04; label: diagnostic.

The night fit's model (Eq. 3) has one ground temperature for the whole night, and T_{add} follows the temperature of the balun and coax outside the box. D5 logged neither. Public stations near the site (Table 6) bound both. The figures and numbers here are from the site-weather analysis of Christian Hellum Bye's EIGSEP D5 analysis, on the stations' published data, retrieved on 2026-11-04.

Table 6: The stations, from the antenna's ground point (1680 m): great-circle distance and bearing, elevation, and elevation less the site's ground.

station	network	distance, bearing	elevation [m]	− site ground [m]	used for
`TULU1`	NRCS SCAN	5.0 km, 256°	1402	−278	air; soil
`TVWU1`	BLM RAWS	11.5 km, 8°	1585	−95	outdoor air
`KCWU1`	UDOT RWIS	28.0 km, 227°	1923	+243	air (check)
`BKRN2`	NOAA USCRN	74.3 km, 250°	2017	+337	air (check)

The outdoor air. The Tule Valley RAWS (`TVWU1`) stands on the same west-flank bench as the site, 11.5 km north and 95 m below the site's ground. We use the position WRCC and NOAA HADS list; IEM lists one rounded to whole arc-minutes, 3.3 km away, and nothing here depends on the choice. Moved to the antenna's height (1770 m) at 6.5 K km^{-1}, its air temperature falls from 299.6 K at 02:00 to 294.3 K at 12:00 UTC on 07-17 (Fig. 11). The three nearer stations, moved the same way, spread by ±1.5 K at the start of the night and ±0.7 K at its end. We take that spread as the uncertainty of the transfer to the site. The lapse rate is a well-mixed daytime value: at night the flank can sit above the valley's cold pool, and the antenna, 90 m up, inside or above a surface inversion. A balun in free air also departs from screened air by radiation. This $T_{\mathrm{out}}(t)$ is a measured shape for T_{add}'s outdoor part, with no free amplitude; the night fit does not use it yet.

The ground. SCAN 2163 (Tule Valley, `TULU1`), 5.0 km WSW on the valley floor and 278 m below the site's ground, records the soil's temperature at 5, 10, 20, 51 and 102 cm in whole degrees Fahrenheit, steps of 0.56 K. Over the 7 days of D5 the daily swing averages 19.0, 18.3 and 5.7 K at 5, 10 and 20 cm. At 51 and 102 cm the reading moves by at most one step in a day (Fig. 12a, b). Over the 07-17 night (02:00–12:00 UTC) it changes by −8.9, −8.9 and −3.3 K at the three shallow depths, and not at all at 51 or 102 cm. If the 50–200 MHz emission comes from 50 cm or deeper, the ground's temperature is constant over the night to within a step, as Eq. 3 assumes.

Its level depends on that depth. In the D5 week the deep soil was near its seasonal maximum: 302.4 K at 51 cm and 299.7 K at 102 cm. The annual means are 289.4–290.8 K at both depths (July–June years 2024–25 and 2025–26; Fig. 12c). Moved to the site's ground with the air's lapse rate (−1.8 K), a metre down in July gives 297.9 K and the deep annual mean 288.2 K. The move assumes the soil follows the air's gradient. The ground-temperature prior (§5.1 and memo 009), 293 ± 30 K, is centred between the two, because the depth the emission comes from is not known (below); its width also covers rock against the valley's soil (Fig. 12d). It is a prior on the ground's physical temperature. The ground's brightness, $(1 - R)\,T_{\mathrm{gnd}}$ plus the sky it reflects, is not covered by its width: with the rock's reflectivities R below, $R\,T_{\mathrm{gnd}}$ alone is tens of kelvin, and the reflected sky is R times the sky the ground sees.

The ground's material. The soil station stands on deep valley alluvium, with no bedrock recorded. The beam's ground is the House Range flank and Dome Canyon: the NRCS soil survey (SSURGO) maps thin soil there over lithic bedrock at 24–46 cm (Table 7), and the canyon exposes Cambrian limestones with interbedded shales, summarised for this site in Reid Parsons' notes. Two U.S. Geological Survey maps cover the beam's ground: the antenna stands on Hintze (1974), *Preliminary geologic map of the Notch Peak Quadrangle, Millard County, Utah*, Miscellaneous Field Studies Map MF-636 (doi:10.3133/mf636), 0.26 km south of its 39°15′ boundary with Hintze (1981), *Preliminary geologic map of the Marjum Pass and Swasey Peak SW quadrangles, Millard County, Utah*, MF-1332 (doi:10.3133/mf1332), which covers the ground north of it, Marjum Pass included. Tabulated ground-penetrating-radar properties (Annan 2005, *GPR methods for hydrogeological studies*, in *Hydrogeophysics*, Springer, pp. 185–213, doi:10.1007/1-4020-3102-5_7; reproduced in Reid Parsons' notes),

limestone ε = 4–8 and 0.4–1 dB m^{-1} and shale ε = 5–15 and 1–100 dB m^{-1}, put the depth over which the emitted power falls by e at 4.3–10.9 m in limestone and 0.04–4.3 m in shale. How much of the temperature wave that depth sees is set by the damping depths, which the soil station measures: 0.12 m for the daily wave, fitted to the 5–20 cm swings (0.11–0.15 m day to day; it predicts 0.48 K at 51 cm, below the resolution as measured), and 4.2 m for the seasonal wave, from the amplitudes (4.4 m) and the 7-day lag (4.0 m) between 51 and 102 cm ($\sqrt{365}$ times the daily depth would give 2.3 m). With these for the rock too, limestone's emission sees under 2% of the daily swing and 23–44% of the seasonal one, so it emits near the annual mean; shale's sees up to 71% of the daily swing and 44–99% of the seasonal one, nearer the July value. The rock's normal-incidence power reflectivity is 11–23% (limestone) and 15–35% (shale), so the ground-reflected sky (§7) is not small, and the ground temperature here does not include it. These depths, fractions and reflectivities are estimates from tabulated properties and the valley soil's heat flow, not D5 measurements; rock conducts heat better than dry alluvium, which would lengthen the damping depths. The soil station measures the valley's alluvium, not the canyon's rock: the prior's centre comes from it together with this depth argument, and its 30 K width covers rock against soil, not the reflectivity's effect on the ground's brightness (above).

Table 7: The NRCS soil survey (SSURGO, Soil Data Access, queried 2026-11-04) at the soil station and at the antenna: the map unit, its major components, their taxonomic classes and the depth to a restrictive layer, as recorded.

	SCAN 2163 (valley floor)	antenna site (House Range flank)
map unit	Dera families assoication, 2 to 12 percent slopes	Amtoft-Amtoft, very shallow-Lodar families assoication, 5 to 60 percent slopes
components	Dera 60%, Dera 25%	Amtoft 35%, Amtoft 30%, Lodar 20%
class	Loamy-skeletal, carbonatic, mesic Typic Calciorthids	Loamy-skeletal, carbonatic, mesic Lithic Calcixerolls; Loamy-skeletal, carbonatic, mesic Lithic Xerollic Calciorthids
bedrock	none recorded	lithic, 24–46 cm

Two questions stay open: the beam-weighted mix of limestone and shale, with their moisture, which sets the emission depth; and whether the rock follows the valley soil's thermal regime. A temperature sensor in the ground at the site would settle the level.




Figure 11: Air temperature near the site (Table 6). (a) The four stations as published, 07-12 to 07-20 UTC; grey: the 07-17 night, 02:00–12:00 UTC. (b) The night, each station moved to the antenna's height (1770 m) at 6.5 K km^{-1}: the RAWS (`TVWU1`, the outdoor air temperature used here) with its transfer uncertainty σ_{tr} (half the three stations' spread) as its band, and the two other nearby stations. The RAWS reports hourly, at HH:36, in whole degrees Fahrenheit. Diagnostic.




Figure 12: Soil temperature at SCAN 2163 (TULU1, valley floor, 278 m below the site's ground). (a) Hourly at five depths through the D5 week; grey: the 07-17 night. (b) The daily range against depth (grey: each D5 day; black: their mean), the change over the night (red) and an exponential fit to the 5–20 cm ranges (dashed; damping depth 0.12 m); at 51 and 102 cm the range is below the 1 °F resolution (0.56 K), drawn as upper limits. (c) The values at local midnight at 51 and 102 cm, 2023-07 to 2026-10 (gaps: missing data), with July–June annual means (dashed; 2023–24 is left out, 65% complete) and D5 (red). (d) The T_{gnd} prior (§5.1 and memo 009), 293 ± 30 K, a prior on the physical temperature and not on the ground's brightness, against the D5-week values and the annual means, all moved to the site's ground (-1.8 K). Diagnostic.

6 The Sun

6.1 When it is visible

Table 8 lists every time the Sun's centre crosses the DEM horizon or 0° altitude while each height era was flown in phase C. Figure 13 shows the horizons and the Sun's tracks. At 87.5 m the Sun clears the eastern ridge at 15:53 UTC on 07-17, at 39° altitude and azimuth 93°, with the edge 352 m away. That is 3.5 h after it reaches 0° altitude at 12:26. At 91 m it sets behind the western ridge at 01:21 on 07-18 (17°, edge 170 m), 1.5 h before 0° (02:53).

Two consequences follow. The rx-B window starts at 01:27, after the terrain sunset, so of the calibrated data only 07-17 15:53–16:22 sees the direct Sun. And the 2026-10-01 trials' observation that the Sun "lifts 199 MHz within an hour of rising" cannot be the direct Sun: that hour is behind the ridge.




Figure 13: DEM horizon against true azimuth for the three heights (`horizon_profiles_v0005`, rotated from grid to true bearings), and the Sun's tracks on 07-17. Dots mark the terrain crossings in the calibrated eras.

era	event	horizon	UTC	Sun alt (deg)	Sun az (deg)	horizon (deg)	edge (m)
30m	set	terrain	07-16 00:00:37	32.2	272.4	32.2	150
30m	set	flat	07-16 02:54:17	-0.0	298.1	0	–
30m	rise	flat	07-16 12:25:27	0.0	62.0	0	–
30m	rise	terrain	07-16 16:24:37	44.8	98.7	44.8	335
87.5m	set	flat	07-17 02:53:44	-0.0	297.9	0	–
87.5m	rise	flat	07-17 12:26:14	0.0	62.3	0	–
87.5m	rise	terrain	07-17 15:53:44	38.8	93.4	38.8	352
91m-reraise	set	terrain	07-18 01:21:17	16.6	283.9	16.6	170
91m-reraise	set	flat	07-18 02:53:07	-0.0	297.6	0	–

Table 8: Sunrises and sunsets during phase C, by height era (`sun_events_v0007`). "terrain": the Sun's centre crosses the DEM horizon traced along its bearing; "flat": 0° altitude. Geometric, with no refraction.

6.2 The test around 15:53

The test reads the night fit, `fit_v0025`. Every bin with the Sun above 0° is out of sample there, so its residual (data minus fitted sky, ground and offset) is a prediction. That residual drifts by tens of kelvin over hours, as

the night model is carried into the morning. So each frequency is fitted locally, from 14:30 to 16:23 UTC, as

$$\mathrm{resid}(t) = c_0 + c_1(t - t_{\mathrm{ridge}}) + S_\odot\, A_\odot(t - \tau).$$

The transmitter of the beam-mapping raster is on 15:36–16:14, with short gaps (box-gnd transitions, memo 001). Its bins are never fitted. That leaves 60 bins, of which only 7 (16:14–16:22) have the Sun above the ridge. Errors on $S_\odot$ are scaled by $\sqrt{\chi^2/\mathrm{dof}}$.

Each of the following checks could have come out the other way:

1. **A null at 0° sunrise.** The same fit around 12:26 (96 transmitter-off bins), with a Sun column that ignores the terrain, should give $S_\odot \approx 0$ if the ridge hides the Sun.
2. **Timing.** χ^2 against a shift τ of the Sun column should be smallest near $\tau = 0$.
3. **Spectrum.** $S_\odot(\nu)$ should be positive and smooth, of the order of the quiet-to-active Sun.




Figure 14: Out-of-sample residuals of the night fit around the ridge crossing (dotted line), at six frequencies. Filled: transmitter off, fitted. Open: transmitter on, not fitted. Dashed: the fitted line alone. Blue: line plus Sun.

6.3 Result

- **Spectrum.** At 33 frequencies between 105 and 235 MHz, $S_\odot$ runs from 2 to 19 SFU (33 positive), rising to 199 MHz and lower above: 4±1 at 109 MHz, 8±1 at 148, 18±2 at 199 and 12±1 at 219 MHz (Fig. 16, Table 9).
- **Null.** At 0° sunrise the same fit gives −5 to 0 SFU over the same band. No step appears when the Sun rises behind the ridge.
- **Timing.** χ^2 is smallest at $\tau = -3$ min, and within 5 of that minimum from −7 to +7 min. A shift of 20 min earlier costs $\Delta\chi^2/(\chi^2/\mathrm{dof}) = 67$ on average.
- **Kelvin scale.** If a's departure from 1 lies in the kelvin scale or the beam, it scales $S_\odot$ too. $S_\odot/a$ is 3, 7 and 16 SFU at 109, 148 and 199 MHz.
- **Below 90 MHz** the result is not usable. The residual drifts by tens of kelvin across the window there, and the null fails (grey and orange points below 90 MHz in Fig. 16).

The spectra themselves. Figure 15 sets aside the local fit and compares spectra. It averages the calibrated spectrum (weighted means) over two windows of transmitter-off bins. "Before" is −40 to −15 min from the ridge crossing, with the Sun behind the ridge (16 bins). "After" is the 7 bins with the Sun above it. Their difference is shown raw, and with the night model's predicted sky and ground change between the windows subtracted. That change is at most 5 K above 109 MHz, but up to 43 K below 87 MHz, where the difference is dominated by drift the model does not capture. The difference is compared with a Sun predicted with nothing fitted to these data. The prediction takes RSTN's concurrent 245 MHz flux (§6.4), extrapolated down in frequency with the MWA quiet-Sun shape (scaled by 0.85), times the Sun column averaged over each window.

- Above 109 MHz the spectrum rises by a smooth 8 K at 148 MHz and 13±1 K at 199 MHz. The flux this implies, 7 and 18 SFU, agrees with the local fit.
- The prediction is 4 and 5 K. The measured jump is 1.1–3.5 times the prediction over 32 frequencies (weighted mean 2.0). Scaled by a, the prediction falls short by a factor of 2.2 below 200 MHz and 1.2 above 205 MHz.
- **Not a gain change.** A change in receiver gain would scale the whole spectrum, so $\Delta T/T$ would be constant. Instead it runs from 1 to 6% (coefficient of variation 0.38), while ΔT itself varies less (0.27). The jump is additive, like a new source, not multiplicative.

MHz	$S_\odot$ (SFU)	χ^2/dof	a	$S_\odot/a$	null at 12:26
109.4	4.3 ± 0.6	187	1.36	3.1	-0.6 ± 0.3
125.0	7.0 ± 1.1	340	1.32	5.3	-1.5 ± 0.4
148.4	7.7 ± 0.6	120	1.05	7.4	-1.6 ± 0.9
175.8	14.7 ± 1.2	51	1.06	13.9	-2.7 ± 0.8
199.2	17.7 ± 1.6	152	1.14	15.5	-3.0 ± 0.8
218.8	12.3 ± 0.9	117	1.48	8.3	-0.3 ± 0.6
234.4	13.1 ± 1.1	21	1.39	9.4	-0.5 ± 0.7

Table 9: $S_\odot$ at selected frequencies: the local fit's χ^2/dof, the GSM amplitude a of `fit_v0025`, $S_\odot/a$, and the flat-horizon null at 12:26.




Figure 15: Top: calibrated spectra before the ridge crossing (Sun behind the ridge) and after it (Sun above, transmitter off). FM is blank. Bottom: their difference, raw (open) and with the night model's sky/ground change removed (filled). Orange: the Sun predicted from concurrent RSTN 245 MHz flux with the MWA quiet-Sun spectral shape, through the beam and the ridge; dashed: the same times the GSM amplitude a. Nothing in the orange curves is fitted to these data.




Figure 16: Left: $S_\odot$ from the transmitter-off fit at the ridge crossing (filled), divided by the GSM amplitude a (open), and the flat-horizon null at 12:26 (squares). Diamonds are external solar monitors (Section 6.4). Right: mean $\Delta\chi^2$ over frequencies against a shift of the Sun column; $\tau > 0$ is later than the DEM predicts.

6.4 Concurrent solar monitoring

The USAF Radio Solar Telescope Network (RSTN) measures the whole-Sun flux at 245, 410 and 610 MHz every second. Two stations recorded continuously through 15:53–16:22 UTC on 07-17. At 245 MHz Sagamore Hill reads 14.9 SFU (rms 0.8, 99th percentile 18) and San Vito 14.1 SFU (rms 0.7, 99th percentile 16), over 1681 and 1680 one-second samples. Palehua's reading is below 5 SFU for 53% of the window, as if not yet tracking the Sun; over the transmitter-off bins with the Sun up it reads 25 SFU. The stations' noon values for 07-17 at 245 MHz are Sagamore Hill 13, San Vito 13, Palehua 21, Learmonth 23 SFU. The stations disagree by more than the quoted 2–3 SFU; we take the two continuous stations' mean, 14.5 SFU, as the concurrent reference at 245 MHz, and note that Palehua reads higher.

The Sun was quiet: SWPC lists no flare or radio burst between 13:18 and 20:16 UTC, the 10.7 cm flux was 104 sfu (Penticton, 17:00 UTC), and e-CALLISTO spectrographs show no burst at 100–250 MHz in the window. Sagamore Hill's spectrograph recorded a series of type III bursts at 25–180 MHz from 12:14 to 13:18, inside the null window, when the Sun was behind the ridge. We have not looked for them in our data.

Nothing calibrated was found below 245 MHz for that day. For the spectrum's shape, the MWA's quiet-Sun measurement (Oberoi, Sharma & Rogers 2017, on 2013-09-03) gives 1.7, 3.5, 5.1, 11.3 SFU at 103, 131, 167, 213 MHz, rising with frequency as ours does, with 10–60% systematic error. At 234 MHz, the top of our band, $S_\odot$ is 13±1 SFU and $S_\odot/a$ is 9 SFU. RSTN gives 14.5 SFU at 245 MHz, a ratio of 0.9 ($S_\odot$) or 0.6 ($S_\odot/a$). Between 105 and 200 MHz, $S_\odot$ is 1.3–3.4 times the MWA quiet-Sun curve, and $S_\odot/a$ is 1.0–2.9 times it. So $S_\odot$ is the quiet Sun's size to within a factor of 3.4. That agreement is not a test that separates the Sun from another effect of similar size at sunrise.

source	when	what it says
RSTN Sagamore Hill, 245 MHz	15:55–16:22	14.9 SFU mean, rms 0.8
RSTN San Vito, 245 MHz	15:55–16:22	14.1 SFU mean, rms 0.7
RSTN Palehua, 245 MHz	16:14–16:22	25.0 SFU mean; below 5 SFU (as if not tracking) for 53% of 15:55–16:22
RSTN noon values, 245 MHz	07-17 local noon	Sagamore Hill 13, San Vito 13, Palehua 21, Learmonth 23 SFU
Penticton F10.7	17:00	103.9 sfu observed
SWPC event list	all day	0007-0008 PAL RBR 245 820; 0150-0150 PAL RBR 245 260; 1214-1318 SAG RSP 025-180 VI/2; 2016-2053 SAG RSP 025-076 VI/2; 2025-2114 G19 XRA 1-8A B6.5 1.8E-03; 2030-2034 PAL RSP 025-075 III/2; 2214-2251 PAL RSP 025-055 VI/2; 2325-2325 PAL RSP 025-037 III/2
e-CALLISTO (6 stations)	15:45–16:30	no burst at 100–250 MHz (uncalibrated, visual)
MWA quiet Sun (Oberoi et al. 2017)	2013-09-03	103 MHz 1.71, 117 MHz 2.6, 131 MHz 3.46, 148 MHz 4.79, 167 MHz 5.08, 189 MHz 6.69, 213 MHz 11.3, 240 MHz 17.12 SFU

Table 10: External solar radio measurements for 2026-07-17 (`inputs/solar_reference.json`, with URLs).

6.5 Solar radio bursts in the raw data

A radio burst is a property of the Sun, recorded independently by other observatories. It is the most direct test of whether box-air receives solar emission. Type III bursts drift from high to low frequency within seconds and last tens of seconds; at 0.54 s per integration, box-air resolves them.

The reference list. `curation/solar_bursts_v0001.json` lists the 32 bursts that e-CALLISTO spectrographs recorded at two or more stations while the Sun was above the Marjum ridge (Table 8). It starts from e-CALLISTO's curated burst list for July 2026, a detector that requires three stations, with times to the minute. Spectrograms from 11 stations in daylight were then inspected to time each burst to about a second and to add bursts the list missed. All are type III or groups of type III. The stations cover 5–93 MHz, apart from one noisy one at 46–226 MHz, so the list cannot say whether a burst reached our higher frequencies.

Method. For each burst, `sun_bursts.py` reads raw box-air antenna rows from 90 s before to 90 s after the burst, with *no* RFI flags, since the flags mark positive excesses, which is what a burst is. It divides each channel by its median over that cutout. It then measures the excess in four bands during the burst, in robust standard deviations of the same band away from it, and finds when each 10 MHz sub-band peaks. Box-air antenna data exist at 12 of the 32 bursts; the rest fall in data gaps or calibration cycles (Table 11).

What the raw data show. Figure 17 shows three bursts as raw dynamic spectra, each channel divided by its median.

- **07-15 18:33 (on the ground).** A narrow ridge of emission appears inside the e-CALLISTO interval and drifts from the 70 MHz range down to the bottom of our band. The 10 MHz sub-bands peak in sequence, at −8 MHz/s. In the light curves, 45–65 MHz rises by 99% of the raw power (115σ), while 170–235 MHz does not move (1σ).
- **07-15 18:23 (on the ground).** Two components about 25 s apart, both low-frequency only, 119σ at 45–65 MHz. The brighter, later one drifts at −9 MHz/s.
- **07-17 20:33 (91 m, during the raster).** The brightest burst of the period (e-CALLISTO's "IIIG/3", a group). Several components sweep down to the bottom of our band. Sub-bands up to 120 MHz peak above 8σ and up to 220 MHz above 5σ, so the group reaches above the reference stations' bands. The raw power at 45–65 MHz rises by 120%. The raster's own modulation is visible behind it.

Test. A random time in daytime is often near one of the impulsive spikes of §6.6, so a big excess alone is not a detection. The bursts e-CALLISTO saw are bright at 45–85 MHz and fainter above, while the impulsive

spikes are broadband. The test therefore counts times that are bright at 45–65 MHz *and* quiet at 170–235 MHz ($z_{45–65} > 10$, $z_{170–235} < 5$), at the bursts and at 250 random times in the same Sun-up windows, at least 5 min from any listed burst (Fig. 18). 5.2% of the random times pass, against 5 of the 12 bursts with data; the chance of that many by accident is 2.2×10^{-4}. This test could have failed.

What this shows. Box-air records solar radio bursts. They come at the times independent stations recorded them, drift from high to low frequency as type III bursts do, and have the low-frequency-weighted spectrum those stations saw. That is a physical identification of solar emission in our data, not a coincidence in timing. It says nothing about the quiet Sun's flux, since the e-CALLISTO spectrograms are not calibrated.

burst start (UTC)	dur. (s)	e-C MHz	sta.	box-air excess (robust σ) 45–65	65–85	110–170	170–235	match
07-15 18:21:37	14	20–85	4	6	3	16	1	no
07-15 18:23:19	43	20–85	5	119	36	10	2	yes
07-15 18:33:02	19	10–85	6	115	4	1	1	yes
07-15 18:47:01	12	10–45	2	no box-air antenna data				–
07-15 22:45:10	5	10–45	2	no box-air antenna data				–
07-15 22:59:08	8	10–40	2	no box-air antenna data				–
07-15 23:49:00	10	15–50	2	11513	1796	3046	2184	no
07-16 16:23:55	19	20–85	6	19	1	2	4	yes
07-16 16:50:02	11	20–85	6	no box-air antenna data				–
07-16 16:54:58	13	20–90	7	61	11	72	19	no
07-16 17:16:06	26	20–85	6	no box-air antenna data				–
07-16 17:25:35	5	46–85	3	no box-air antenna data				–
07-16 17:26:22	10	20–85	6	no box-air antenna data				–
07-16 17:57:21	50	46–80	3	no box-air antenna data				–
07-16 18:33:13	31	20–85	6	no box-air antenna data				–
07-16 18:48:05	11	20–65	3	no box-air antenna data				–
07-16 19:17:07	16	20–85	5	no box-air antenna data				–
07-16 21:36:13	8	20–55	2	no box-air antenna data				–
07-16 22:05:44	10	15–45	2	no box-air antenna data				–
07-16 22:25:10	9	15–50	2	no box-air antenna data				–
07-16 22:28:10	16	15–60	3	no box-air antenna data				–
07-16 22:32:42	9	55–85	3	no box-air antenna data				–
07-16 23:58:59	34	20–75	2	17	9	16	20	no
07-17 20:16:26	14	30–85	4	44	9	3	0	yes
07-17 20:30:45	46	10–60	5	8	2	2	4	no
07-17 20:32:59	37	10–90	6	35	10	5	4	yes
07-17 20:34:14	23	10–85	5	5	2	0	1	no
07-17 21:21:22	99	10–60	4	4	4	2	2	no
07-17 22:26:42	49	15–45	2	no box-air antenna data				–
07-17 22:46:54	19	20–60	4	no box-air antenna data				–
07-17 22:51:02	21	15–40	2	no box-air antenna data				–
07-17 23:25:37	17	15–45	2	no box-air antenna data				–

Table 11: The 32 reference bursts (`curation/solar_bursts_v0001.json`): duration and frequency range at the e-CALLISTO stations, number of stations, and box-air's excess in four bands. "match": bright at 45–65 MHz and quiet at 170–235 MHz ($z_{45–65} > 10$, $z_{170–235} < 5$).




Figure 17: Three e-CALLISTO bursts in raw box-air data, time running down. Left: each channel divided by its median over the cutout, minus 1. Dotted lines are the burst's start and end at the reference stations, and the green bar along the top is their frequency range. Circles mark when each 10 MHz sub-band below 95 MHz peaks, where it is clear (above 8 robust σ) and within 8 s of the 45–65 MHz peak. Right: the same in four bands, on the same time axis.




Figure 18: The burst test. Each point is one 15–40 s window of raw box-air data. Its position says how much the raw power rose during the window, at 45–65 MHz (across) and at 170–235 MHz (up), in units of that band's usual scatter. Blue: the windows of the e-CALLISTO bursts, labelled by start time. Grey: 250 random windows in the same daytime hours. The bursts e-CALLISTO recorded are bright at low frequency and faint above 170 MHz, so a real one should land in the shaded region ($z_{45–65} > 10$, $z_{170–235} < 5$). The impulsive spikes of §6.6 rise in both bands and land at the upper right. 5 of 12 bursts fall in the shaded region, against 5.2% of the random windows. Axes are linear between −1 and 1 and logarithmic beyond.

6.6 Impulsive events in the raw data

Looked at without a reference list, raw box-air is full of impulsive events: single integrations that rise at every frequency at once, about 102 times an hour in daytime (Fig. 19, as horizontal lines). They do not drift. `impulsive_scan.py` finds them in phase C (35 h of antenna data at confirmed heights) and sorts them by whether the Sun is above the ridge at the antenna, up but behind the ridge, or down. Overall they come 102 per hour with the Sun visible (7.2 h), 75 with it behind the ridge (11.2 h), and 7 at night. Earlier versions found the visible and behind-the-ridge rates equal and called the events local; that comparison included the stretches on the ground, where they were most frequent.

These classes differ in height era and time of day as well as in the Sun, so `validate_impulsive.py` holds one variable at a time, on exposure rebuilt row by row in 1-minute cells (Table 12). Events cluster, so beside exact Poisson intervals it gives a bootstrap over 30-minute chunks, which is far wider and the one to use.

- **Within one era.** At 30 m the rate is 170 per hour with the Sun visible against 69 behind the ridge, but the latter is 154 on 07-15 and 12 on 07-16, so one day carries it; for events of three or more integrations the visible rate is higher on both days (22 and 33 against 9 and 6). After the 07-18 re-raise, one evening at 91 m, it is 490 visible against 272 behind the ridge, but 35 against 42 for the longer events. (The 07-17 raster day at 91 m has the Sun high throughout, 88 per hour, and no behind-the-ridge time to compare.)
- **The same hours of local time** with the Sun visible and behind the ridge overlap in only 2 hours (10:00, 19:00 MDT). Pooled over them, the visible rate is 3.6 times the other (1.8 for the longer events), but at 19:00 after the re-raise alone the ratio is 1.65 (0.57 for the longer events).
- **Against the Sun's height above the horizon**, at 30 m the rate shows no step where the Sun crosses it: 204 per hour with the Sun 5–10° behind the terrain, 46 within 2° below it and 103 within 2° above. After the re-raise it peaks within 2° of the horizon (608 and 795 per hour, against 185 with the Sun more than 10° behind it), but that is one sunset evening.

For all events, the Sun-visible rate is the higher one in every era and at the same hours; but at 30 m the rate does not step where the Sun crosses the horizon, the longer events go the other way after the re-raise, and several comparisons rest on one day or two chunks. The controls neither show the events following the Sun nor rule it out, so their origin is open. That they are broadband single integrations that do not drift, unlike the bursts of §6.5, is why a search for bursts must start from an independent list.

era	Sun	hours	events	rate, all (h^{-1})	rate, 3+ integrations (h^{-1})
30 m	visible	1.6	276	170 (151–192; 107–228)	27 (20–36; 16–36)
30 m	behind ridge	6.6	454	69 (63–75; 16–134)	6.8 (5.0–9.1; 2.8–11.1)
30 m	night	9.5	68	7.1 (5.5–9.1; 1.9–15.4)	0.4 (0.1–1.1; 0.0–1.1)
87.5 m	visible	2.8	38	14 (10–19; 1–38)	4.7 (2.5–8.0; 0.0–13.6)
87.5 m	behind ridge	3.1	24	7.7 (4.9–11.5; 1.9–15.1)	1.3 (0.3–3.3; 0.0–3.4)
87.5 m	night	7.5	40	5.4 (3.8–7.3; 1.8–10.3)	0.5 (0.1–1.4; 0.0–1.7)
91 m	visible	2.5	218	88 (77–100; 36–141)	24 (18–31; 3–54)
91 m, re-raised	visible	0.4	196	490 (424–564; 274–579)	35 (19–59; 34–35)
91 m, re-raised	behind ridge	1.3	363	272 (245–302; 185–424)	42 (32–55; 25–82)
91 m, re-raised	night	0.1	6	48 (18–104; 0–103)	0.0 (0.0–29.3; 0.0–0.0)

Table 12: Impulsive-event rates within each height era by the Sun's position (`validate_impulsive.json`; `impulsive_v0004`, rows at confirmed heights): rate per hour with its 95% exact Poisson interval and, after the semicolon, the 95% bootstrap interval over 30-minute chunks, which allows for clustering. "visible": the Sun above the traced horizon; "behind ridge": above 0° altitude but below it.




Figure 19: Two minutes of raw box-air data in daytime with the Sun behind the ridge, each channel divided by its median, time running down. The horizontal lines across all frequencies are the impulsive events.

6.7 Sunrise and sunset in the raw data

The calibrated data hold one terrain crossing. Phase C holds 4, the rest uncalibrated (Table 8). `sun_transitions.py` keeps the raw context of each, minute by minute, ±60 min (Figs. 21 and 22): the mean raw antenna spectrum, whether the transmitter was on, whether the antenna moved, and how many rows were antenna rows rather than calibration states.

It then fits raw power at each beam frequency as a line plus the Sun column at that crossing's pointing, over the parked stretch of data containing the crossing, without transmitter-on minutes unless the whole stretch has it on. Raw power has no kelvin scale. The fit gives $S_\odot/T_{\rm sys}$, and the 5th–95th percentile of the calibrated temperatures of 07-17 stands in for $T_{\rm sys}$. That stand-in comes from other hours and heights, so the fluxes in Table 13 are good to a factor of about 2.

The crossings in context. At these pointings the Sun's predicted signature in raw power is not a step but a slow bump. As the Sun sinks towards the ridge, the beam's gain toward it changes, and the ridge then cuts it off over 10–30 min. So it peaks some 20–30 min before sunset (the sunrise version is the blue curve of Fig. 14). It amounts to a few percent of the power for a Sun of a few SFU. Figures 21 (sunsets) and 22 (sunrises) show what the raw data do. Earlier versions also listed the 07-15 sunrise and the 07-17 sunset at 30 m; the antenna was on the ground for both (§4.1), there is no horizon to trace from 2 m, and they are left out.

- **07-16 sunset, 30 m, transmitter off**: the quietest crossing. Raw power rises steadily, by about 10% over the two hours, at all four frequencies. The fit puts a Sun of 12–15 SFU at 199 MHz on top of that rise, mostly from a bump 15–30 min before sunset. The bump is no larger than the unexplained wander elsewhere in the window.
- **07-16 sunrise, 30 m**: box-air's self-EMI comb (memo 006) is on until 16:51, and its lines cover much of the raw spectrum; the fitted fluxes are erratic.
- **07-17 sunrise, 87.5 m**: the transmitter is on for most of 15:36–16:14 (orange ticks), and its leakage dominates those minutes (the 10–30% excursions in the bottom panel). Calibration cycles also interrupt the antenna rows (grey). On transmitter-off minutes, including ones after the calibrated window ends at 16:22, the raw power is flat to about 1%, and the Sun term is 14–17 SFU at 199 MHz.
- **07-18 sunset, 91 m, transmitter on throughout**: fitted on channels midway between the comb teeth. The raw power wanders by several percent, and the Sun term is 16–20 SFU at 199 MHz.

A solar burst at the 07-17 sunset, on the ground. The antenna was on the ground, tilted to about −66°, from 07-16 23:27 to 07-17 01:15 (§4.1), not hanging at 30 m as earlier versions assumed. `validate_sunset_burst.py` reads the raw spectra of both boxes from 23:44 to 00:08 UTC and divides them by their median over the quiet minutes on either side (Fig. 20). From 23:53:45, box-air's power below 100 MHz rises far above that. Over 23:55:30–00:00:00 it is 1.09 times the reference at 50–60 MHz, 1.54 at 66–72 and 2.30 at 78–84 MHz, and 1.18 at 100–120 and 0.99 at 180–230 MHz.

- **It is not interference.** It is smooth across the band, and channels at whole megahertz rise like their neighbours (1.58 against 1.56), so it is not box-air's 1 MHz self-EMI comb (memo 006). Box-gnd, on the same correlator, stays at 0.95–1.05 times its reference below 90 MHz. The excess's lower edge drifts from about 80 to 50 MHz over ten minutes, with a fainter band near 160–175 MHz.
- **It is not pointing.** At 72–84 MHz it peaks at 2.5 times the reference at 23:58, at an elevation of −73°. At 23:44, at the same elevation, the power was 0.97 times the reference.
- **When it ends.** In 10 s means, the excess falls below 1.5 times the reference at 00:00:45 and below 1 at 00:02:35, while the elevation is still wandering about its park value of −73°. The elevation then steps at 00:02:45 and is within 1° of its final −66° from 00:03:25. With the step the power falls by a further 18%.
- **A flare was in progress.** GOES-18 recorded a C1.5 X-ray flare from 23:51 to 00:09 UTC, peaking at 00:02 (SWPC event 7990, `inputs/solar/20260716events.txt`). The same event has a 245 MHz burst of 820 SFU at 00:07–00:08 (Palehua); box-air at 180–230 MHz stays within 4% of its reference from a minute before it to two minutes after. e-CALLISTO stations (HAARP, COHOE) recorded a type III burst at 23:58:59 (34 s, 20–75 MHz) inside the excess (Table 11).

So the excess is solar emission during that flare: it is box-air's alone, broadband and smooth, and it coincides with the flare and with the type III burst. Its slow downward drift and the band near twice its frequency resemble a type II burst, but no observatory's sweep is listed to confirm the type. Why it ends is open. Earlier versions timed its end against the Sun's disk going behind the ridge as seen from 30 m and read it as occultation; from the ground the horizon is higher and not traced (there is no 2 m profile), so that timing is withdrawn. The flare's own decay, the Sun setting behind the terrain as seen from where the box stood, or the tilted antenna's response could each end it. Most of the fall still comes before the elevation moves; only the last 18% comes with the elevation step. That box-gnd misses it is not explained either: its position is not solved, and here both boxes were on the ground.

What the crossings say. Over the 3 fitted crossings without the self-EMI and 121–219 MHz (where above 2σ), the Sun term runs from 3 to 41 SFU. The concurrent prediction (RSTN at 245 MHz with the MWA quiet-Sun shape, Fig. 15) is 2–11 SFU over the same frequencies. RSTN's noon flux at 245 MHz stayed between 13 and 17 SFU from 07-15 to 07-18. So the raw data allow a Sun of the expected size. They do not detect it convincingly: the Sun term is a few-percent bump on raw power that drifts by as much for other reasons, and the kelvin stand-in is good only to a factor of 2. The calibrated 07-17 local fit, 18 SFU at 199 MHz, is 0.9–1.2 times the upper ends of their ranges there, and the same 07-17 crossing in raw data, with a longer stretch of transmitter-off minutes, gives 14–17 SFU. Earlier versions, on `tcal@v0003`, found the calibrated excess several times larger than any raw crossing.

The timing is weakly constrained. The best shifts τ^* in Table 13 range from −30 to −2 min, with broad χ^2 minima (0 at the ±60 min edge of the scan). The ground under the antenna goes into shadow about half an hour before the antenna at the 30 m sunset and nearly two hours before at the 91 m one, and comes into sunlight 15–50 min after it at the sunrises ("ground" column). A signal from the lit ground, or from heating there, would therefore lead at sunset and lag at sunrise. These data cannot tell those timings apart.




Figure 20: The 07-17 sunset in raw spectra, with box-air on the ground and tilted (`validate_sunset_burst.npz`; raw correlator files, all integrations on the antenna; 10 s means). Left and middle: box-air and box-gnd power divided by their median over quiet minutes before 23:53 and after 00:03 UTC, minus 1, with time running down. Right: band power over its pre-burst median for box-air at 72–84 MHz (solid orange) and 180–230 MHz (dashed), and box-gnd at 72–84 MHz (blue); the antenna's elevation in grey (top axis). Grey dashed: the elevation steps away from its park value. The excess is box-air's alone, drifts downward in frequency, and is mostly gone before the elevation steps.

crossing	status	fit (min)	τ^* (min)	ground (min)	121	148	199	219
					$S_\odot$ (SFU, range from T_{sys} stand-in)			
set 07-16 00:00 (30m)	fitted	-44 to +28	-4	-33	6–9	7–9	12–15	18–22
rise 07-16 16:24 (30m)	fitted; self-EMI on	-42 to +10	-16	+18	-9—13†	46–62	1–1†	29–36
rise 07-17 15:53 (87.5m)	fitted	-44 to +40	-2	+50	4–5	4–5	14–17	3–4
set 07-18 01:21 (91m-reraise)	fitted	-44 to +10	-30	-115	28–41	17–23†	16–20	23–28

Table 13: Raw-power fits at the terrain crossings (`transitions_v0007`): status, the fitted stretch and best timing shift τ^* (minutes from the crossing), when the ground under the antenna crosses ("ground", minutes), and $S_\odot$ at four frequencies (MHz) for the 5th–95th percentile of the 07-17 calibrated temperatures. †: below 2σ.




Figure 21: The three terrain sunsets in raw box-air data, one column each, time running down; the dotted line is the Sun's centre crossing the ridge at the antenna. Top: the raw spectrum, minute by minute, divided by its median over the two hours. Ticks at the right mark minutes with the transmitter on (orange), the antenna moving (black) or calibration rows (grey); the brown dashed line is when the ground under the antenna goes into shadow. Bottom: raw power at 148 and 199 MHz, divided by its median, on the same time axis. Where a fit was made, the dashed line is its baseline and the solid line baseline plus Sun. If the Sun is seen, the points follow the solid line, not the dashed one, through the half hour before the crossing.




Figure 22: The three terrain sunrises, as in Fig. 21; here the brown dashed line is when the ground under the antenna comes into sunlight, and the Sun's signature follows the crossing instead of preceding it.

6.8 The Sun's path through the beam

Figure 23 shows the Sun column, minute by minute over phase C (`sun_profile_v0007`, 2320 minutes with pointing). Parked, it changes slowly with the Sun's motion, between the terrain crossings. Moving, it changes fast: during the 07-17 raster the beam's gain toward the Sun swings by factors of several within seconds. On 07-15 and 07-16 much of the day has no valid pointing in the table, and outside the raster the table's azimuth zero inherits potentiometer slips (memo 003). So the predicted Sun there is uncertain.

The raster offers a test with no kelvin scale. Since the receiver gain multiplies everything, regressing raw power on the sky (GSM through the beam), the ground, a constant, the transmitter's leakage, a drift and the Sun column gives the Sun's flux on GSM's brightness scale (`raster_sun.py`). The null is the same fit with fake Suns at the same altitude in 11 other azimuths. The test fails (Table 14). The real Sun beats every fake at 3 of 13 frequencies with the HFSS beam and 10 of 25 with the empirical beam, and the fitted fluxes change sign (3 and 7 negative). The residual, 1.8% of the power with HFSS, depends on direction and is larger than the Sun's modulation. Any extra direction-dependent column absorbs it.




Figure 23: Simulated, not data: the predicted Sun column (antenna temperature per SFU of solar flux) through box-air's HFSS beam over phase C, at six frequencies, from the Sun's position, the antenna's recorded pointing and the knife-edge terrain visibility (`sun_profile_v0007`). No measured power is plotted.

	HFSS beam				empirical beam	
MHz	$S_\odot$ (SFU)	fakes $\geq$	resid.	MHz	$S_\odot$ (SFU)	fakes $\geq$
62.5	-2	9/11	1.4%	64.5	-5	2/11
74.2	-4	4/11	1.3%	70.3	-4	6/11
113.3	6	2/11	1.8%	76.2	-9	5/11
125.0	0	10/11	1.6%	82.0	-1	8/11
136.7	3	5/11	1.7%	113.3	7	0/11
148.4	5	2/11	1.4%	119.1	3	1/11
160.2	3	5/11	1.7%	125.0	1	8/11
171.9	4	5/11	2.5%	132.8	3	5/11
183.6	9	1/11	2.0%	138.7	3	5/11
195.3	71	0/11	1.8%	144.5	3	4/11
207.0	13	0/11	2.6%	150.4	5	2/11
218.8	21	0/11	4.2%	156.2	5	2/11
230.5	-1	9/11	8.8%	162.1	2	7/11
				168.0	17	1/11
				173.8	2	5/11
				179.7	11	0/11
				185.5	14	0/11
				191.4	17	0/11
				197.3	851	0/11
				203.1	-29	0/11
				209.0	47	0/11
				214.8	-137	0/11
				220.7	-55	0/11
				226.6	18	0/11
				232.4	9	6/11

Table 14: The raster test (`raster_sun_v0007`): the Sun's flux on GSM's scale, how many of the 11 fake Suns improve χ^2 as much or more, and the residual rms as a fraction of the power.

6.9 What this does and does not show

- **Box-air receives solar emission.** Solar radio bursts recorded by independent spectrographs appear in raw box-air data at the same seconds, with the type III drift and the low-frequency-weighted spectrum those stations saw (§6.5). That is a physical identification, not a coincidence in timing.
- **The raw crossings allow a quiet Sun of the expected size but do not detect it.** Across 3 uncalibrated terrain crossings on three days and three heights, the Sun term is 3–41 SFU at 121–219 MHz, against a concurrent prediction of 2–11 SFU (§6.7). It is a few-percent bump on raw power that drifts by as much for other reasons, and its kelvin stand-in is good only to a factor of 2.
- **The calibrated 07-17 excess is about twice the predicted Sun.** At 15:53 the calibrated spectrum jumps by 2.0 times the concurrent prediction (1.5 times scaled by a), and the local fit gives 0.9–1.2 times the other crossings' upper values at 199 MHz. RSTN shows no brightening that day. Anything that changes as the antenna and its box come into sunlight follows the same geometry, so it is not separated from the Sun. The excess rests on 7 one-minute bins at the end of the calibrated window, all directly after the transmitter switches off.
- **The timing does not separate the Sun from the lit ground.** Best timing shifts range from -30 to -2 min, with broad minima.
- **The beam's sweep during the raster does not isolate the Sun** with the present sky/ground/beam model: fake Suns fit as well (§6.8).
- $S_\odot$ inherits the unresolved amplitude a (0.9–1.7) and HFSS's gain at $\sim 50°$ from boresight.

7 Recovering the sky and the ground

The night fit (§5) holds one pointing at one height, so the ground temperature and the offset are degenerate there. This section tries three ways round that, each with a check that could fail.

7.1 Tying the frequencies together

At one pointing and height, the beam's ground fraction $f_{\mathrm{gnd}}(\nu)$ still changes with frequency. If the ground is one temperature at all frequencies (thermal emission) and the receiver offset is a smooth function of frequency,

the pedestal $f_{\rm gnd}(\nu)T_{\rm gnd} + T_{\rm off}(\nu)$ could separate them. `joint_fit.py` reduces each frequency of the night fit to a likelihood for $(T_{\rm gnd}, T_{\rm off})$, marginalizing the sky and the GSM amplitude. It then fits one $T_{\rm gnd}$ and a Legendre polynomial in $\log\nu$ for the offset over 105–235 MHz. If the data fixed $T_{\rm gnd}$, it would not move with the polynomial's order, nor between the halves of the band. It does: 245 to 902 K as the order goes from 1 to 6, and the two halves disagree by hundreds of kelvin (Table 15). The ground fraction's frequency dependence is too much like a smooth offset; smoothness alone does not fix $T_{\rm gnd}$.

offset order	$T_{\rm gnd}$ (K), all	lower half	upper half
1	245 ± 44	714 ± 158	427 ± 119
2	381 ± 79	749 ± 165	418 ± 119
3	339 ± 80	767 ± 166	888 ± 131
4	716 ± 98	773 ± 166	887 ± 131
5	902 ± 103	772 ± 166	886 ± 132
6	873 ± 103	772 ± 166	887 ± 132

Table 15: $T_{\rm gnd}$ from the night fit tied across frequency (`joint_v0007`), against the order of the offset's polynomial, over the whole band and its two halves.

7.2 The sky as calibrator on parked nights

Most nights have no kelvin scale, but the Galaxy drifts through the beam. In a stretch of fixed pointing, raw power at each frequency is fitted as gain × (GSM through the beam) + a constant + a linear drift (`sky_cal.py`). The gain is then in raw units per GSM kelvin. The constant over the gain is $T_{\rm rx} + f_{\rm gnd}T_{\rm gnd}$ in GSM kelvin, and stretches at different heights, with different $f_{\rm gnd}$, would separate the two. Each night is fitted at one height, on its rows at that height only (§4.1); the receiver regime is rx-A throughout. A stretch enters that comparison only if the sky fits it, with a median residual under 5% of the power. Figure 24 shows the raw data with the fit.

- **07-17, 87.5 m** (calibrated): the sky fits to 0.4% rms. The gain against `tcal`'s gives a =1.0–1.3 at 105–220 MHz, as the night fit found. This checks the pipeline, not where a comes from.
- **07-16, 30 m**: box-air's self-EMI is on all night; with its teeth masked the sky fits to 3.0%.
- **07-15**: the antenna was on the ground all night, not at 30 m, and there is no horizon profile at 2 m, so the night is not fitted. Earlier versions fitted it as 30 m from 05:12 UTC; that fit is withdrawn. It also explains why its raw power before about 04:45 was a few percent of its later level.

That gives 2 nights, whose ground fractions span only 0.04. A 10 K difference between the nights in anything else then shifts $T_{\rm gnd}$ by about 228 K, and the fit gives −2265 to 2670 K at 120–180 MHz. Parked nights at these two heights cannot separate the ground.

7.3 The ground from the rotating antenna

When the antenna moves, the beam's ground fraction swings between about 0.5 and 0.75 within seconds, while the receiver gain stays put. In four moving windows (`raster_sky.py`), raw power per integration is regressed on GSM through the beam, the ground fraction, a constant, the transmitter's leakage, the Sun and a drift. The gain multiplies every column, so the ground coefficient over the GSM coefficient is $T_{\rm gnd}$ in GSM kelvin, with no kelvin calibration. Errors are jackknifed over 2-minute blocks. Each window uses only its rows at the window's height (§4.1). The 07-18 night window keeps the 8290 of its 16260 rows after the re-raise; 4370 were on the ground and 3600 of unknown height. The raster loses its last 240 rows, of unknown height. Earlier versions also used a 30 m dusk window on 07-17 00:21–01:08, which was on the ground; it is dropped.

The excess in the raw data. Figure 25 shows it in the raster's raw power, integration by integration, as the beam's ground fraction swings between 0.60 and 0.75 over each elevation sweep. Take the raw power with no RFI flags at all, and subtract every fitted term except the ground: the sky, the constant, the transmitter, the Sun and the drift. What is left at 64 and 82 MHz rises and falls with the ground fraction by 18 and 20% of the power, peak to peak. A 300 K ground would move it by a few percent. The slope of that residual against the ground fraction is the ground's brightness: 1290 and 736 K in GSM kelvin with no flags, and 279 and 322 K




Figure 24: The parked nights at a confirmed height in raw data at 148 and 203 MHz, each divided by its median (points), with the fitted gain × GSM through the beam + constant + drift (lines) (`skycal_v0008`, HFSS beam). The 07-15 night, on the ground, is not fitted.

on the RFI-flagged samples (19 and 22% of them). So without flags the raw data carry a low-frequency excess, and on the flagged samples it is small. Neither value is clean: the unflagged power holds the interference the flags mark, and the flagged power a selection that follows the pointing. At 162 MHz the two differ too: −921 K unflagged and 225 K flagged (53% kept). At 113 MHz the unflagged raster is dominated by the transmitter's leakage, so only flagged values are quoted there. The flags keep fewer than a third of the raster's samples below 120 MHz, and the fraction kept depends on where the beam points. The flagged values above 100 MHz therefore carry a selection that the unflagged ones do not.

Figure 26 and Table 16 show the result in every window.

- **It repeats.** The three 91 m windows (two evenings, the transmitter and the Sun in different states) agree at 7 frequencies between 120 and 180 MHz (χ^2/dof up to 6.4).
- **It falls gently with frequency**, from 333 K at 125 MHz to 189–290 K at 150–175 MHz.
- **Below 85 MHz only the raster constrains it**, at 279±505 K at 64 MHz and 322±118 K at 82 MHz. The other two 91 m windows give errors of thousands of kelvin there (at 82 MHz, 4324±6533 and 1970±2288 K).
- **The 30 m window differs**: 2810±626 K at 64 MHz and 995±623 K at 82 MHz, and tens of kelvin above 130 MHz. It has the self-EMI on.

Thermal ground emission should not depend on frequency. At 125–180 MHz the values are those of soil near 290 K with an emissivity of 0.65–1, but that coincidence does not identify them as the ground's temperature. The nearest soil station puts the soil a metre or more below the site at 288.2–297.9 K, depending on depth, and shallower soil warmer (§5.2).

Not reflected sky. Ground reflects part of the sky, which is far brighter at low frequency. A specularly reflected sky term (each below-horizon direction mirrored to the sky above, flat ground) was added in the raster, with the ground's reflectivity R free. Soil would give $R \approx 0.1$–0.4. At 64–82 MHz the fit gives $R = -0.16$ to 0.18, and the ground's emission term −183–651 K: the flagged raster does not constrain a reflected sky.




Figure 25: The ground excess in the raster's raw power (`raster_sun_v0007`, raw correlator files for the unflagged values; channels midway between the transmitter's comb teeth). Top: four elevation sweeps of raw power with no RFI flags, less every fitted term except the ground (points), against the fitted ground term (blue) and what a 300 K ground would give (orange), in percent of the power. Bottom: the same residual, binned by the beam's ground fraction and converted to GSM kelvin, with no flags (filled) and with `flags@v3` (open); the slope is the ground's brightness.

Diffuse reflection is tested over phase C in §7.5.




Figure 26: (a) The ground term in each moving window, as the ratio of the ground and GSM coefficients (GSM kelvin; empirical beam unless marked; jackknife errors; points with errors above 300 K omitted). Shaded: 200–300 K. The axis is linear below 100 K and logarithmic above. (b) For the three 91 m windows, the same ground term less a thermal 250 K, divided by GSM's whole-sky mean (below 180 MHz). A sky leaking in with a fixed fraction would give a constant.

7.4 Tracing the excess

Earlier versions found the ground term reaching thousands of kelvin below 85 MHz in every window, and traced it: it was not specular reflection, not noise from the horizon, not concentrated in azimuth, and not a fixed fraction of the sky. On the present mount (with the axle tilt) and left-handed pointing, the flagged raster gives a few hundred kelvin there, so that excess is not established. The unflagged raster (1290 and 736 K at 64 and 82 MHz) and the 30 m window still carry one. The mount, the azimuth's handedness and the empirical beam all changed at once, and which of them removed the excess is not tested.

Where on the ground. `ground_excess.py` repeats the regression with the ground split into regions, each with its own brightness (Table 17). With flags, the deeper ground holds a few hundred kelvin at 64–82 MHz within its errors, and the band within 10° below the horizon scatters by more than its errors; without flags both

MHz	mean of three 91 m windows (GSM K)	χ^2/dof
125.0	333 ± 33	0.1/1
144.5	253 ± 23	1.6/1
150.4	290 ± 22	12.9/2
162.1	226 ± 8	1.3/2
168.0	272 ± 33	0.2/2
173.8	189 ± 9	1.3/2
179.7	233 ± 45	0.1/1

Table 16: The ground term, weighted mean of the three 91 m windows, where at least two have errors under 200 K, and the windows' agreement.

are higher. In the raster, eight azimuth sectors give 487–4622 K at 76 MHz (typical error 897 K). The other windows sample too few orientations to constrain eight sectors.

Against the sky. A sky seen where the model expects ground would give an excess proportional to the sky's brightness. Figure 26b divides the ground term less 250 K by GSM's whole-sky mean. In the raster the ratio is −0.00–0.04 at the frequencies below 90 MHz with errors under 300 K, and within ±0.2 at 150–175 MHz.

A gain that follows the orientation. If the system gain or the antenna's match changes as the antenna turns, by an amount that follows the ground fraction, the change has the shape of the ground column and these regressions cannot separate it. A gain change that multiplies only the sky (an extra column of ground fraction times GSM) leaves the ground term at 64–82 MHz unconstrained: −1789 to 12808 K, against 246–499 K without it. Testing an orientation-dependent match needs the antenna's reflection coefficient at different tilts. All 19 field antenna-S11 sweeps from 07-13 on were taken upright: those with pointing (19) are within 3.2° of el = 0.

So far. On the flagged raster the ground term is a few hundred kelvin from 64 to 190 MHz, falling gently with frequency. Whether the excess left in unflagged data and in the 30 m window is interference the flags remove, sky, or the beam below the horizon is open. The one-variable comparison of the old and new geometry would say what removed it from the flagged raster; the daily term over phase C (§7.5) is a separate test, with the sky drifting under fixed pointings.

window, region	64	70	76	82	113	125
raster, flags@v3: within 10° below horizon	-1557±719	-2749±1681	-1479±1152	1546±425	12±254	121±140
raster, flags@v3: deeper ground	99±553	-54±513	-48±292	541±110	392±38	315±38
raster, no flags: within 10° below horizon	–	4766±881	371±1418	379±861	–	632±244
raster, no flags: deeper ground	1238±980	2000±228	976±373	696±225	–	400±28
07-17/18 night, flags@v3: within 10° below horizon	–	–	–	–	–	16±1738
07-17/18 night, flags@v3: deeper ground	–	–	1123±1515	–	–	381±376
raster, 64 MHz, 8 azimuth sectors	9263±22052, 7654±30967, 2293±9070, 5136±16884, 8964±26934, 8936±19392, 16592±63978, 10373±31527					
raster, 76 MHz, 8 azimuth sectors	2216±664, 3875±1284, 487±312, 2844±790, 1664±713, 4500±1031, 4302±1003, 4622±1685					

Table 17: The ground split into regions (`ground_excess_v0007`, empirical beam; brightness in GSM kelvin, jackknife errors; "–": error above 2000 K). Top: the band within 10° below the terrain horizon and the deeper ground, at six frequencies (MHz). Bottom: the raster's eight azimuth sectors (0–45°, 45–90°, . . . , east of north).

7.5 The low-frequency daily term over phase C

The moving windows each last about an hour, so the sky hardly changes within one and cannot be told from a fixed ground term. Phase C adds what they lack: three days of sidereal time. `global_fit.py` models rx-A raw power in `raw_binned_v0006` over all of phase C (4236 bins in 36 gain windows that pass its residual gate), one frequency at a time. The model is a smooth gain times [GSM through the HFSS beam + a ground term + an offset]. The log gain is piecewise linear in time, with knots every 3 h and a 5% prior on each knot step. There are no sky corrections ($l = 0$). Alternate 30 min blocks are held out, and every model is scored by the held-out rms as a fraction of the power. The code is in data-analysis, run at commit 076e255. `validate_phase_c.py` in this directory runs each configuration on the 12 beam frequencies of the original analysis, 62.5–175.8 MHz; the

6 from 62.5 to 82.0 MHz are reported. The ground term takes one of these forms, each with one free parameter unless stated:

- a constant temperature T_{g};
- diffuse (Lambertian) reflection of the sky, $(1 - R)\,290\,\mathrm{K} + R\,\bar{S}(t)$. Here $\bar{S}$ is the cosine-weighted GSM mean over the open sky above the era's horizon, 1152–5189 K at these frequencies, so a ground term of thousands of kelvin is the size this form would give;
- the same, with $\bar{S}$ taken 6 h earlier, 6 h later or 12 h away in sidereal time, as controls;
- "beam": the beam's response below the horizon falling on the sky of the current pointing instead of on the ground;
- "pickup": $k\,\bar{S}$ without the ground fraction, for sky entering by a path that does not turn with the antenna;
- a housekeeping stream, averaged over each bin from the correlator metadata: three RF-switch thermistors, the load temperature and the system current;
- "free level": $T_0 + R\,\bar{S}$ with both free and a 290±300 K prior on T_0.

Heights. Phase C here is only its rows at 30, 87.5 and 91 m (§4.1). Earlier versions also fitted the 07-15 night and the 07-17 sunset park as 30 m, which were on the ground; most of the term was found with those rows in, and the results below differ from those versions accordingly.

Over all of phase C (Table 19), reflection with the real sky beats a constant ground by 0.14–0.27 points of held-out rms, with $R = 0.36$–0.75. The shifted skies gain up to 0.58 points, as much or more, so the real sky is not singled out. The beam form differs from a constant ground by at most 0.19 points, so the term does not follow the sky of the current pointing. Pickup does better than reflection (by 0.11–0.15 points, with 3 of its fits diverged), so the ground fraction is not what carries the term.

Against recorded temperatures and current. On the 100% of bins with all five streams, the best single stream does between −0.27 and 0.17 points worse than reflection at 62–78 MHz (a negative value: better); at 82 MHz the first thermistor reaches 2.84% against 2.87% for reflection. With all five streams in the fit at once, reflection with the real sky still lowers χ^2/dof by 1.8–3.0, against at most 1.3 for the shifted skies, and R stays 0.39–0.85 at 62–74 MHz. So χ^2 still prefers the real sky beside the streams, while the held-out residual does not separate it from the shifted skies.

With the level free the ground's constant part is no longer tied to R. The sky shifted by 6 h then does worse than the tied real sky at 62–70 MHz (by −0.36 to −0.20 points, negative meaning worse), and at 74–82 MHz the tied real sky beats every shifted free-level fit by 0.10–0.25 points. The free real sky itself (0 of 6 fits diverged) gives $R = 1.5$–3.2 with T_0 from −6 to −1 kK: far more time variation than reflection allows, offset by a negative level.

Disjoint subsets (Table 20). Splitting phase C at 07-16 18:10 UTC gives halves A and B. In half A the reflection form with the real sky (tied or with a free level) is the best model at 6 of 6 frequencies, and the 6 h-shifted sky does 0.2–0.5 points worse; χ^2/dof is 44–65. In half B (1424 bins, χ^2/dof 3–4) the real sky is also the best model at 6 of 6, at least 0.04 points ahead of every shifted sky, and reflection gains −0.07 to 1.21 points over a constant ground. Earlier versions found no term in half B; its misfit there was dominated by a look-down park that was the antenna tilted on the ground, now out. Half A split by day at 07-16 00:00 UTC gives 07-15 (311 bins, only the evening at 30 m), where every fit diverges, and 07-16 (2501 bins), where R is −0.09 to 0.36 (errors up to 0.08) and a shifted sky is the best model at 3 of 6 frequencies. So each half prefers the real sky, but 07-16 alone and phase C as a whole do not.

Window selection. `global_fit.py` drops gain windows whose median residual over the fitted frequencies is above 5% of the power, so which windows enter can depend on the frequencies fitted. Gated on 62–82 MHz alone, the same 4236 bins in 36 windows pass as with all 12 frequencies, so the gate does not change these results.

Faster gain. The fits above put a gain knot every 3 h, and §7.6 finds the gain changing within the hour. `validate_phase_c_fastgain.py` refits phase C with knots every 3 h, 1 h and 20 min, and injects that sub-hour gain into mocks (Table 18; data-analysis commit 076e255). Every fit anchors the GSM amplitude in tcal kelvin at $a_{\mathrm{tcal}} = 1.17$, this model's level at 62–82 MHz with 3-h knots, either softly (± 0.02 in log, which these data outweigh) or hard ($\pm 10^{-4}$).

- **With faster knots the term goes.** With 3-h knots, reflection with the real sky beats a constant ground by 0.15 to 0.40 points. With 1-h knots the difference is 0.04 to 0.12, and with 20-min knots −0.38 to 0.00. Freed this way, the gain level is $a_{\mathrm{tcal}} = 0.44$–0.92 against the soft anchor at 1-h knots. Held at 1.17 by the hard anchor, reflection is fitted with $R = 0.14$–0.52 at 1-h knots and does worse than a constant ground (−0.59 to −0.05 points).
- **The gain takes up the sky's daily shape.** Per gain segment, the log gain knots of the constant-ground fits correlate with $\log \bar{S}$ of the real sky at −0.47 to +0.81 (3-h knots), +0.85 to +0.93 (1-h) and +0.80 to +0.93 (20-min), and with the shifted skies at −0.64 to +0.15 (1-h) and −0.63 to +0.19 (20-min).
- **The sub-hour gain does not make the term.** The mocks take the 3-h fits as truth, times the frequency-flat fast gain of §7.6 (1.9% rms, its temperature-driven part included). Without a daily term, reflection gains −0.04 to 0.08 points with 3-h knots, and with the real data's own residuals multiplied in −0.07 to 0.15, against the real data's 0.15–0.40. A mock with a daily term gives 0.16–0.29 (and 0.01–0.27 with the residuals), like the real data.
- **The residuals would hide a real term.** A mock with a daily term ($R = 0.36$–0.73), the fast gain and the real residuals gains only 0.01 to 0.27 points with 3-h knots and −0.04 to 0.12 with 1-h knots, although R is recovered at 1-h knots (0.29–0.71). Without the residuals it gains 0.16–0.29 points with 3-h knots, 0.00–0.03 with 1-h and −0.35–0.28 with 20-min knots. With 20-min knots and the residuals, the comparison is erratic in both mocks: −0.36 to 0.09 points without a term and −0.19 to 0.13 with one. So with 1-h or 20-min knots the gain comparison cannot show a term of this size, and its loss with faster knots does not rule one out.

	3 h knots		1 h knots		20 min knots	
	gain	R	gain	R	gain	R
Real data, soft anchor	+0.15 to +0.40	0.36 to 0.73	+0.04 to +0.12	−0.21 to 0.19	−0.38 to +0.00	−0.22 to 0.48
Real data, hard anchor	–	–	−0.59 to −0.05	0.14 to 0.52	−3.97 to −0.34	0.12 to 0.66
Mock, no daily term, fast gain	−0.04 to +0.08	0.37 to 0.73	−0.04 to −0.01	0.21 to 0.69	−0.60 to +0.02	0.27 to 0.72
Mock, daily term, fast gain	+0.16 to +0.29	0.48 to 0.80	+0.00 to +0.03	0.31 to 0.71	−0.35 to +0.28	0.36 to 0.76
Mock, no daily term, fast gain and residuals	−0.07 to +0.15	0.37 to 0.69	−0.10 to +0.09	0.17 to 0.68	−0.36 to +0.09	−0.17 to 0.79
Mock, daily term, fast gain and residuals	+0.01 to +0.27	0.47 to 0.86	−0.04 to +0.12	0.29 to 0.71	−0.19 to +0.13	0.00 to 0.83
Gain knots against $\bar{S}$: real sky	−0.47 to +0.81		+0.85 to +0.93		+0.80 to +0.93	
shifted skies	−0.50 to +0.41		−0.64 to +0.15		−0.63 to +0.19	

Table 18: The daily term against faster gain (`validate_phase_c_fastgain.json`), at 62–82 MHz. "gain": held-out rms of a constant ground minus that of reflection with the real sky, in percentage points (positive: the daily term helps), over the converged fits; R: the fitted reflection coefficient. Soft and hard anchors hold a_{tcal} at 1.17 by ± 0.02 and $\pm 10^{-4}$ in log. The mocks are the 3-h fits times the fast gain of §7.6, with radiometer noise, and in the last two rows with the real fits' residuals multiplied in (rolled by 1500 bins). Bottom: the correlation of the constant fits' log gain knots with $\log \bar{S}$, per gain segment.

What this shows. On rows at confirmed heights, a slow additive term at 62–82 MHz improves the phase-C fit with 3-h gain knots over a constant ground, by as much as a mock with a daily term and more than mocks with only the fast gain. Within each half of phase C the real sky beats every shifted sky. Over the whole of phase C, and on 07-16 alone, skies shifted by 6 or 12 h in sidereal time do as well, so the term is not shown to be sky. With 3-h knots the fitted gain no longer follows the open-sky mean consistently (−0.47 to +0.81); with 1-h knots it does, and the term goes. Where the receiver's gain after the RF switch is measured (§7.6), it does not follow the sky. The 30 m era has one usable calibration interval, 07-16 01:41–02:24 UTC with the antenna moving, in half A; over it the gain after the switch changes by at most 1.4% at 110–205 MHz, against −2.7 to 1.4% for the 20-min fit's gain, and at 43 min it is far too short to show a daily term. Whether the term is additive, a gain change on 1–3 h scales, or the fit absorbing sky or beam mismatch, these data cannot tell. Over phase C sidereal time gains only 3.9 min a day on solar time, so separating a sky term from an instrumental daily cycle needs observations weeks apart; the ground rows could add a second height once a 2 m horizon exists.

MHz	constant	real	+6 h	−6 h	12 h	beam	pickup	R (real)
62.5	3.53	3.33	3.68	**3.31**	3.45	3.50	3.25†	0.75±0.03
66.4	3.54	3.28	3.34	**2.96**	3.01	3.73	2.88†	0.64±0.03
70.3	2.89	2.72	3.01	2.79	2.88	2.93	**2.59**	0.47±0.03
74.2	2.98	2.83	3.01	2.83	2.86	3.01	**2.72**	0.36±0.02
78.1	3.65	3.42	3.50	**3.26**	3.27	3.74	3.27	0.42±0.03
82.0	3.14	2.87	3.03	2.83	**2.83**	3.25	2.78†	0.68±0.04

Table 19: The phase-C fits on all bins (`validate_phase_c.json`): held-out rms in percent of the power for each ground form (one free parameter each; "real" and the shifted skies are the reflection form), and the reflection coefficient R of the real sky. Bold: the best fit in the row. †: the fit diverged (a gain knot below 1% of the median, or an offset above ten times the GSM column).

MHz	constant	real, tied	real, free	+6 h	−6 h	12 h	R (real, tied)
Half A							
62.5	4.74	4.43	**4.38**	4.96	4.60	4.73	0.78±0.05
66.4	4.75	4.36	**4.21**	4.79	4.73	4.85	0.65±0.04
70.3	3.72	3.50	**3.30**	3.82	3.71	3.75	0.47±0.04
74.2	4.01	3.85	**3.65**	4.08	4.00	3.96	0.35±0.04
78.1	5.15	4.92	**4.80**	5.16	5.15	5.08	0.38±0.04
82.0	4.52	4.28	**4.27**	4.57	4.52	4.50	0.65±0.05
Half B							
62.5	1.37	1.44	**1.35**	1.39	1.56	1.79	0.46±0.02
66.4	1.53	**1.41**	1.48	1.56	1.72	1.81	0.44±0.02
70.3	2.37	**1.66**	2.23	2.31	2.58	2.65	0.42±0.03
74.2	1.58	**1.49**	1.59	1.73	1.69	1.75	0.36±0.03
78.1	2.01	**1.66**	1.93	1.89	2.11	2.54	0.61±0.03
82.0	2.89	**1.68**	2.78	2.86	2.96	3.25	0.39±0.03
07-15							
62.5	nan†	nan†	–	nan†	nan†	nan†	-4605.38±117.25†
66.4	nan†	nan†	–	nan†	nan†	nan†	-6923.18±299.81†
70.3	nan†	nan†	–	nan†	nan†	nan†	-550.48±1392.71†
74.2	nan†	nan†	–	nan	nan†	nan†	2413.47±38.64†
78.1	nan†	nan†	–	nan	nan†	nan†	1.07±0.70†
82.0	nan†	nan†	–	nan†	nan†	nan†	231.47±2613.91†
07-16							
62.5	4.25	4.18	–	4.28	**4.17**	4.20	0.36±0.08
66.4	4.45	4.18	–	4.47	4.35	**4.07**	0.33±0.06
70.3	3.57	**3.45**	–	3.93	3.62	3.49	0.13±0.05
74.2	3.83	3.77	–	4.10	3.87	**3.68**	0.05±0.05
78.1	4.85	**4.59**	–	5.24	5.06	4.63	-0.09±0.06
82.0	4.23	**3.96**	–	4.33	4.18	4.33	0.11±0.06

Table 20: The same comparison on disjoint parts of phase C (`validate_phase_c.json`). Half A is phase C before 07-16 18:10 UTC, half B after it. 07-15 and 07-16 split half A at midnight UTC. "real, free" and, in the halves, the shifted skies have a free level with the 290±300 K prior; in the two days the shifted skies are tied, like the real sky. Bold: the best converged fit in each row; †: diverged.

7.6 The ground term with a fast gain

The phase-C fits above hold the gain smooth over 3 h. This subsection asks whether the data need a gain that changes faster, and what ground term follows at 110–205 MHz. `global_fit.py` (data-analysis commit 4db041a) fits rx-A in `raw_binned_v0006` over phase C at 21 frequencies from 51 to 234 MHz, one at a time: a smooth gain times [GSM through the HFSS beam + $T_{\rm gnd}$ times the ground fraction + an offset + the transmitter's leakage + the Sun], with sky corrections to $l = 2$. The gain prior allows 10% per knot step. Bins during the 07-16 23:50–00:02 UTC solar flare (§7.5) are excluded, which leaves 4236 bins in 36 windows. Alternate 5 min blocks are held out. `validate_ground_term.py` in this directory runs every fit quoted here.

Which gain. With knots every 20 min instead of 3 h, the held-out rms falls by 0.2–0.6 points, while $T_{\rm gnd}$ at 117–180 MHz changes by a factor 0.5–1.3 (at 203 MHz, 240 and 274 K). Earlier versions, on the previous mount and pointing, found it rising several-fold with the faster knots. Held-out blocks score how well each model predicts nearby bins, not whether its $T_{\rm gnd}$ is right, so the choice is tested by injection (Fig. 27). A mock is one fit's own prediction for every bin, with radiometer noise, times the 20-min fit's real residuals rolled in time, so they keep their size and timescale but not their alignment with pointing. Its $T_{\rm gnd}$ and gain are then known.

- A mock with the 3-h fit's slow gain is recovered by both knot spacings, the 20-min one to 103 K rms (at most 196 K; 514 K at 141 MHz, where the truth is 499 K).
- A mock with the 20-min fit's gain, refitted with 3-h knots, returns the low values the real data give with 3-h knots, within 74 K at every frequency (535 K at 141 MHz, against the 20-min fit's 579 K).

Both cases return the real 3-h values, and the two knot spacings give similar $T_{\rm gnd}$, so the injections no longer tell which gain the data behave like. The held-out rms prefers 20-min knots, and the multiplicative term they add is called the fast gain below; where the receiver's own gain is measured it is not that ("The receiver's own gain"). The 20-min fit is the one used below, with the caveat of "Where the fit collapses".




Figure 27: $T_{\rm gnd}$ at 110–205 MHz (rx-A, `raw_binned_v0006`, `validate_ground_term.json`; frequencies where the 20-min fit collapses omitted). Left: the real data with 20-min gain knots (errors from the injections) and 3-h knots, and a mock with the 20-min fit's gain refitted with 3-h knots, which returns the real 3-h values. Right: a mock with the 3-h fit's gain (its truth in grey), refitted with either knot spacing. Points are offset by 0.8 MHz for legibility.

The ground term and its error (Table 21). At 110–205 MHz $T_{\rm gnd}$ is 274–579 K. The error is the rms over 3 injections of recovered minus true $T_{\rm gnd}$ (two rolls of the residuals on the 20-min mock, one on the 3-h mock): 16–411 K, 0.4–5.0 times the fit's own error (43–154 K), which takes the residuals as white. The draws average 64 K. With three draws per frequency the error is a scale, not a precise σ. The residuals in the mocks do not follow the pointing. Real residuals that do, from terrain or beam errors, could bias $T_{\rm gnd}$ in a way these

injections do not test. This ground term and the rotating antenna's (Table 16) are both a few hundred kelvin over 125–180 MHz, but they come from different models, beams and data, and are not combined.

MHz	$T_{\rm gnd}$ (K)	injection (K)	fit (K)	$a_{\rm tcal}$	
50.8	4766	996	280	2.34	
58.6	2029	456	195	2.70	
66.4	424	403	420	1.39	
74.2	–	–	–	0.002	collapsed
82.0	771	349	333	1.55	unstable
117.2	342	315	147	0.62	
125.0	–	–	–	0.414	collapsed
132.8	577	65	79	0.88	
140.6	579	16	43	1.83	
148.4	158‡	71	–	0.71‡	shared-gain fit
156.2	405	411	82	0.90	
164.1	410	363	154	0.62	
171.9	–	–	–	0.001	collapsed
179.7	44	90	227	0.38	unstable
187.5	358	212	344	0.36	unstable
195.3	167‡	77	–	1.29‡	shared-gain fit
203.1	274	89	72	2.06	
210.9	–	–	–	1.572	collapsed
218.8	–	–	–	0.001	collapsed
226.6	250	48	35	65.09	unstable
234.4	–	–	–	21.369	collapsed

Table 21: $T_{\rm gnd}$ over phase C with 20-min gain knots (rx-A, `validate_ground_term.json`). "injection": the rms of recovered minus true $T_{\rm gnd}$ over the injections, the error to use; "fit": the fit's own error. $a_{\rm tcal}$ is the GSM amplitude, gain times `tcal`'s scale. "collapsed": $a_{\rm tcal}$ below 0.15 of the 3-h fit's at the same frequency (or either fit diverged), where no $T_{\rm gnd}$ is quoted. "unstable": collapses once the transmitter's bins are dropped. ‡: a different model, the high-band fast gain of the shared-gain fit ("A fast gain shared across frequencies", `validate_shared_gain.json`), which holds only at 141 MHz; quoted at 148, 195 MHz, where the per-frequency fit collapses. Its error is the injection rms over 3 mocks of that model.

Where the fit collapses. The fast gain takes up much of the sky's variation. At the frequencies of the table the 20-min fit's GSM amplitude is $a_{\rm tcal}$ =0.6–2.1, 0.27–1.48 of the 3-h fit's at the same frequency (1.4–2.5); the night fit on calibrated data gives 0.9–1.7. At 74, 125, 148, 172, 195, 211, 219, 234 MHz it falls below 0.15 of the 3-h value (0.11 at 148 MHz; at 6 of them one of the two fits diverges instead), and the ground, offset, transmitter and Sun terms take up the level. The amplitude falls by degrees, so the threshold is a choice, not a gap between two solutions. Holding $a_{\rm tcal}$ near 1.39 with a prior of ±0.2 in log brings it back above the threshold at 4 of these 8 frequencies (0.2–20.7), and changes χ^2/dof by at most 0.39% and the held-out rms by at most 0.01 points. The prior is soft: its rows are not scaled by χ^2/dof, so these data can outweigh it (§7.5 uses a hard one). Where it does pull $a_{\rm tcal}$ back, the fit is no worse, so the data do not fix the amplitude there. The sky is in those data. In the longest parked stretch (07-16 02:48–16:31 UTC, 30 m, 961 bins), raw power at these frequencies tracks the GSM column with correlations of 0.76–0.99, with no gain model at all; in a shorter 87.5 m stretch (07-17 16:29–18:50, 145 bins, where the sky changes less) 4 of the 8 exceed 0.5. In the long stretch the slope against the column is up to 2447 times the gain the collapsed fit puts there, where at the table's frequencies the ratio is 0.39–2.9. Then only the contrast between pointings ties the GSM's amplitude down, and the transmitter spoils it: the 4 windows with the transmitter on reach 9.0% residual, against a median of 0.6% in the others and 0.4% at 148–180 MHz. Dropping transmitter bins (3804 bins left) rescues 74, 125, 148, 195 MHz but turns 82, 180, 188, 227 MHz into collapsed fits. χ^2/dof is 3.9–18.4 where the fit collapses, against 4.0–11.7 at the frequencies of the table. These frequencies are left out of the ground term. At 51–66 MHz the fit holds with the fast gain, so the daily term of §7.5 could be tested with it there; this memo does not yet do so.

What the fast gain follows. Take the 20-min fit's log gain, average it over the frequencies of the table, and subtract a 3-h piecewise-linear fit: what is left varies by 1.9% rms over 110 knots, and the averages of two disjoint halves of those frequencies correlate at 0.96. `gain_drivers.py` projects every housekeeping stream in the correlator metadata onto the same knots, detrends it the same way and compares the two in two disjoint stretches: 07-15/16 at 30 m, and 07-17 at 30, 87.5 and 91 m (Fig. 29). The gain falls weakly as the RF switch warms, but the switch's temperature explains little of it. The robust (Theil–Sen) slopes are −0.9 and −0.7%/K; a bootstrap over blocks of 6 knots, which respects the correlation between neighbouring knots, puts the least-squares slope between −1.9 and 0.4%/K on 07-15/16 (95%, including zero) and between −3.1 and −0.1%/K on 07-17, and the temperature explains 0.04 and 0.04 of the variance. By fifths of the knots in temperature, the median fast gain runs from −0.3 to 1.2% in the four cooler fifths and is −1.1% in the warmest (−0.4 to 0.6 and −0.5% on 07-17). The switch's fast temperature varies by 0.4 and 0.4 K rms, so at these slopes it accounts for a few tenths of a percent of the 1.9% fast gain. No other stream (the system current, the IMU, the motors, the pot, calibration and VNA cycles, the flag occupancy) correlates with the same sign at better than $|r| = 0.12$ in both stretches. Around the end of calibration cycles, VNA sweeps and antenna moves, the 3-h fit's residual moves by at most 0.7% (1.9 times its standard error), against up to 0.1% around random times. 96% of the fast gain's variance has no recorded cause.

The receiver's own gain. The calibration visits of `tcal@v0004` measure the gain after the RF switch without any sky model: the load's power over its temperature, and the noise source's excess over the load, which also carries the source's own drift. `validate_cal_gain.py` compares them with the fits (data-analysis commit 076e255). They are dense only on 07-17, 04:11–16:22 UTC at 87.5 m, with the antenna mostly parked: 59 pairs 12 min apart. The 30 m era has too few to test, and nothing before the switch (antenna, balun, cable) is seen. Over that stretch (Fig. 28) the measured gain at 110–205 MHz varies by 0.8–1.3% rms by the load and 2.8–2.9% by the noise source, and follows the switch's thermistor (30.3–33.4 °C): the band's common mode changes by −1.1%/K by the load ($r = -0.92$) and −2.5%/K by the noise source ($r = -0.76$), slopes of the size the fast gain shows against the same thermistor but with far tighter correlation ("What the fast gain follows"). After a running 3-h mean is subtracted, its correlation with the GSM column is −0.19 to 0.09 at every frequency: no 1–3 h structure follows the sky. Its common mode then varies by 0.5% rms (noise source), against 1.6% for the 20-min fit's gain (correlation 0.60). Over the same hours the 20-min fit's gain at the table's frequencies moves by 3.3–25.3% rms and follows the GSM column, with correlations up to 0.88 (133 MHz) and above 0.5 at 2 of the 9 frequencies of the table. In this parked stretch the fast gain is therefore not the receiver's gain after the switch. It is a gain change before the switch, or the fit absorbing a mismatch in the sky or beam model at a fixed pointing. The 30 m era has one usable interval. The raw correlator files of 07-16 (`validate_cal_thirty.py`; 39 files, SHA-256 recorded) hold 3 load visits. A fixed rule drops a visit that lies more than 10% from the visits' median at more than 2 of the 21 frequencies. It drops the first, at 01:14 UTC with only 22 rows, which fails at 6. Those frequencies all lie within 0.125 MHz of an integer MHz, and every frequency where it passes at least 0.19 MHz from one, as contamination by a comb with 1.000 MHz spacing would give. Memo 001 dates the onset of box-air's 1.000 MHz comb on 07-16 to 01:16:57, after this visit, so its source is not identified. Between the other two, 01:41–02:24 UTC (43 min, the antenna moving through the 07-16 raster), the load gain after the switch changes by −0.9 to 1.4% at 110–205 MHz and by at most 0.6% at 51–82 MHz. The 20-min fit's gain changes over the same minutes by −2.7 to 1.4% at 110–205 MHz and −9.7–12.2% at 51–82 MHz, while the GSM column rises 8.1–8.6%. Over that one interval the two are of similar size at 110–205 MHz, but at 51–82 MHz the fitted gain moves far more than the gain after the switch. With the antenna moving, it may there absorb a mismatch that follows the pointing rather than the sky, and one interval says nothing about 1–3 h structure or about the nights where the daily term of §7.5 is found. Fixing the gain to either calibration series, with one free level (Table 22), cannot test the daily term of §7.5 on 07-17. That stretch has none even with free 3-h knots (reflection with the real sky against a constant ground: −0.01 to 0.10 points). With the gain fixed, the held-out residual rises from 0.37–1.02% to 0.63–2.19%, and a shifted sky beats the real one at 1 (load) and 6 (noise source) of the 6 frequencies: the calibration gain is not accurate to the 1% this test needs. (The fixed-gain fits tie two gain knots one span apart. Earlier versions put them 10^6 s apart, which left the second without training bins, so `global_fit.py`'s held-out rule scored only one bin per frequency; those held-out values are replaced here.)

A fast gain shared across frequencies. The 20-min fit's fast gain is nearly the same at every frequency ("What the fast gain follows"), so one fast term could serve many frequencies, and a frequency could then




Figure 28: The receiver's gain after the RF switch from the calibration visits (`tcal@v0004`; load power over load temperature, and noise source minus load), the 20-min fit's gain at the same times, and the GSM column through the beam, each as a log less its mean, on 07-17 at 87.5 m (`validate_cal_gain.json`). At both frequencies the fitted gain moves far more than the measured gain.

	3-h knots			fixed, noise source			fixed, load		
MHz	const	real	shifted	const	real	shifted	const	real	shifted
62.5	0.81	+0.10	+0.10	2.19	+0.23	+0.58	0.84	−0.02	−0.08
66.4	0.48	+0.02	+0.03	1.86	+0.22	+0.61	0.69	+0.13	+0.07
70.3	0.55	+0.00	−0.00	1.85	+0.20	+0.78	0.70	+0.09	+0.11
74.2	0.50	+0.03	−0.00	1.76	+0.29	+0.78	0.67	+0.16	+0.12
78.1	0.38	−0.01	−0.02	1.79	+0.28	+0.82	0.63	+0.14	+0.13
82.0	0.54	+0.01	−0.03	1.78	+0.34	+0.91	0.76	+0.20	+0.19

Table 22: The daily term on 07-17 04:11–16:22 UTC (87.5 m, 628 bins; the phase-C model of §7.5 with $l = 0$; `validate_cal_gain.json`): held-out rms of a constant ground (%), and the improvement on it of reflection with the real sky and with the best of the ±6 h and 12 h shifted skies (percentage points). Gain free with 3-h knots and a soft amplitude anchor, or fixed to the calibration series from the noise source or the load, with one free level.

no longer collapse by letting its own gain follow its sky. `validate_shared_gain.py` (data-analysis commit 4db041a) fits each frequency with a 3-h gain times one fast gain shared across the training frequencies: knots every 20 min, less what the 3-h gain can carry, with a gain and a tilt in frequency, or a quadratic in $\log \nu$. The shared gain and every training frequency's parameters are fitted jointly (Levenberg–Marquardt), until their steps are at most 9.4×10^{-7} in log gain (at most 139 iterations). An earlier version solved them in turn and did not converge, and its band comparison does not survive convergence. One selection rule is needed: a training frequency that collapses during the fit (a_{tcal} below 0.15 of its own 3-h fit's, or diverged) is dropped from the training set and the fit repeated, since otherwise it wanders along its own flat direction and drags the shared gain.

The rule leaves almost nothing to share. Trained on 117–203 MHz, the fits drop 117, 125, 133, 148, 156, 164, 172, 180, 188, 195, 203 MHz (gain and tilt) and 117, 125, 133, 148, 156, 164, 172, 180, 188, 195, 203 MHz (quadratic), keeping only 141 MHz. Trained on 51–203 MHz they drop 51, 74, 82, 117, 125, 133, 141, 148, 156, 164, 172, 180, 188, 195, 203 MHz (gain and tilt, keeping 59, 66 MHz) and 51, 66, 74, 82, 117, 125, 133, 148, 156, 164, 172, 180, 188, 195, 203 MHz (quadratic). Shared within each band, with its own gain and tilt below and above 100 MHz, the low band keeps 51, 59, 66, 74, 82 MHz and the high band only 141 MHz. Fitted jointly with a shared fast gain, every other high-band training frequency's GSM amplitude collapses, so the high band's "shared" gain is 141 MHz's own. The injection refits apply the same rule. Of the 12, none drops exactly what the real fit with the same training drops (0 and 0 of 6); they drop 203 MHz in 11, 188 or 195 MHz in 4 and a low-band frequency in 2. So which frequencies collapse depends on structure in the real data that the mocks, built from the model's own prediction, do not carry.

Applied to the rest of the high band, 141 MHz's fast gain holds a_{tcal} at 0.34–0.64 of each frequency's 3-h value, and in the within-band fit it removes the collapse at 74, 125, 148, 172, 195, 234 MHz (6 of the 8; a_{tcal} =0.7–2.2). There its held-out rms is −0.47 to 1.63 points from the per-frequency fit's, and at the frequencies of Table 21 −0.08 to 0.46 points. Trained on 59, 66 MHz alone, the gain shared across both bands does not fit the high band: its held-out rms at 110–205 MHz is 9.1–20.9%, against 1.2–1.8% for the per-frequency fit, and 6 of its fits there diverge. One fast term does not serve both bands. At the frequencies the within-band fit rescues, mocks of that fit (three rolls of the real residuals, the same drop rule) give its error. A value enters Table 21 only if no refit drops or collapses there and the injection rms is below 50% of the value. That admits 148, 195 MHz, with injection rms of 70.5–77.4 K, marked in the table as coming from this model. The two models differ: at 141 MHz the within-band fit gives 230 K where the table's per-frequency value is 579 K, so the marked rows continue the within-band fit's spectrum, not the table's.

7.7 Low-order sky corrections

The same regressions can add corrections to GSM: real spherical harmonics up to $l = 3$ in Galactic coordinates, whose fitted amplitudes over the GSM coefficient are corrections in GSM kelvin. Over an hour the sky turns only about 15° relative to the ground. So the same harmonics frozen to the ground (terrain, horizon or beam errors would look like that) are fitted too, and both are scored on held-out alternate 2-minute blocks (Fig. 30). The sky version beats the ground version at 17 of 25 frequencies in the raster, and in 2 of the 4 windows the corrections do not improve the held-out residual at most frequencies. Sky structure is not identified. With the harmonics in, T_{gnd} is no longer constrained either, since the lowest ones mimic the ground fraction. Telling sky from ground needs rotation at many sidereal times.




Figure 29: The fast log gain (20-min knots, mean over the frequencies of Table 21, less a 3-h piecewise-linear fit) against the RF switch's thermistor treated the same way, in two disjoint stretches of phase C (`validate_ground_term.json`). Grey: every knot; black: the median and interquartile range in each fifth of the temperature range; red: the Theil–Sen line, with the 95% block-bootstrap range of the least-squares slope above each panel. Neither stretch shows the gain following the temperature.




Figure 30: Held-out residual in each moving window, as a percentage of the power, without corrections to GSM (grey), with low-order sky-fixed corrections (blue) and with the same frozen to the ground (orange, open). The corrections do not reliably help: in 2 of the 4 windows (2 of the 3 at 91 m) the uncorrected model has the lowest held-out residual at most frequencies, and the sky-fixed corrections beat the ground-fixed ones at 17 of 25 frequencies in the raster. No sky structure is identified.

8 Open questions

- A horizon profile at 2 m (`horizon_profiles_v0005`) would bring back what is now left out because the antenna was on the ground: the 07-15 night, the ground half of the 07-18 night, and the geometry of the 07-17 sunset burst, including whether the terrain ended it (§4.1).
- What makes a vary from 0.9 to 1.7 across the band: the kelvin scale, the beam, GSM, or all three? Memo 004's noise-source temperature and memo 005's beam bear on it directly.
- The ground temperature needs calibrated data at more than one height. Extending `tcal` to the 30 m era would give the height lever of Table 1.
- box-gnd stands upright (like box-air at el = 0) with its box on the ground. Its position there is not yet solved. Before it can give a flux scale, a box-air × box-gnd cross-correlation, or a control for the Sun or for heating, its baseline, polarization and differential horizon need substantial work. It is not used here.
- Calibrating the 30 m-era crossings (07-16) would turn the raw-crossing fits of §6.7 into fluxes, and test whether the 07-17 excess is the Sun.
- What removed the low-frequency excess from the flagged raster: the mount's tilt, the azimuth's handedness or the empirical beam? Changing one at a time would say. Whether what remains in unflagged data and in the 30 m window is interference, sky, the beam below the horizon or a match that changes with tilt is open (§7.4); S11 or a calibration cycle at several tilts would test the last.
- Separating sky corrections from ground-fixed structure needs rotation at many sidereal times.
- Is the low-frequency daily term over phase C sky, instrumental, or a gain the 3-h knots miss? On rows at confirmed heights each half of phase C prefers the real sky, but phase C as a whole and 07-16 alone do not, and over phase C sidereal and solar time stay nearly in step (§7.5). Data weeks apart, or a recorded quantity that tracks it, would decide.
- What is the fast gain? Where the receiver's gain after the switch is measured it is not that, and 96% of it follows no recorded stream (§7.6). A gain change before the switch, or the fit absorbing sky or beam errors, is left; calibration cycles in the 30 m era, or a measurement of the antenna path's loss with temperature, would separate them. Until it is known, the choice between 3-h and 20-min knots changes $T_{\rm gnd}$ by up to a factor of two, and with 20-min knots the data do not fix the GSM amplitude.
- The model is still far from the noise. Candidates are the beam (HFSS against the empirical beam), the sky prior, a time-variable ground, and the 10% GSM prior absorbing structure it cannot predict.

Revision history

2026-11-06 The analyses behind §5.1 and §5.2 are named by what they are, and their frozen inputs carry neutral provenance; no number changes. §5.1 says which tilt the night fit used: the IMU's +0.19° from the readings before the firmware calibration's accelerometer bias was removed; with it removed the night's mean is +0.18° (memo 003). The raster beam on the night is now compared for v0018 alone: Fig. 8 shows the held-out χ^2 with the HFSS beam and with v0018 and their difference, and a new Fig. 9 shows the two fits' data, model and residual at three frequencies; v0017 is dropped from the comparison, and every v0018 number is unchanged. The Allan deviation (Fig. 10) is drawn here from the night-residual analysis's stored curves, now with the single channels' 16–84% range. "Native" becomes "HFSS" for the HFSS beam. Section 2 states that the sky columns' frame, from `eigsep_sim`'s `rot_gal2top_stack`, is 0.19–0.27° from the true local frame, with the fix pending upstream and its effect on the team's estimator.

2026-11-04 (draft) Added an independent night fit on `fit_v0025`'s inputs (§5.1): one sky scale θ and one additive level P per frequency, the HFSS beam integrated harmonically, and the axle tilt measured by box-air's IMU (+0.19 ± 0.6°, memo 003) instead of the beam fit's. At the measured tilt the held-out χ^2 falls from 139.8 to 27.2, and the generic-trend test no longer adopts a time trend. The fit cuts the 44 bins in which the IMU shows the box moving under wind; on the bins left, a scan of constant tilt is lowest at the IMU's. Means over frequency take their errors from jackknife replicates that delete the same time block at every frequency. A method chain records one path from `fit_v0025` to this fit. The night measures θ and the mean level far better than P, which is the level at zero sky; the strong θ–P correlation

is not a degeneracy of the data. On `tcal@v0004` the fit needs an antenna-side gain $\theta > 1$; with the bench's switch-path correction to the noise source $T - T_L$ would scale by $r(\nu)$, 0.85–0.88, a stated rescaling for a later release, and the gain would still exceed the loss model's. The residual's three unexplained features (a slow wave above 100 MHz, an early-night excess, extra low-frequency scatter) are stated as open; the multiplicative excess in the noise is a red part common to the band (its Allan deviation). With memo 005's raster beam (v0018 or v0017) in place of HFSS the night is predicted worse, also on the frequencies the raster measured, with the loss in the upper beam; HFSS stays the fit's beam. The wind over the night is quoted against the cut. Added the site's weather and soil temperature from public stations (§5.2): the outdoor air temperature over the night for $T_{\rm add}$, and a ground with no daily swing below 50 cm, at 297.9 K a metre down in July or 288.2 K at the deep annual mean; the site's thin soil over Cambrian limestone and shale, whose radio depth decides between the two. The ground-temperature prior, 293 ± 30 K, is centred between them; it is a prior on the ground's physical temperature, and its width does not cover the reflectivity's effect on the ground's brightness. The load term's dependence on the loss prior and its held-out cost are stated, and the candidate terms for the residual are tabulated with their controls.

2026-11-02 Moved to memo 005's beam `empirical_raster_v0017` (its mount, with the axle tilt), memo 003's left-handed `pointing@v5` and memo 004's `tcal@v0004`, with binning in raw power (60 s bins) and calibration as separate steps; every fit, product and validation rebuilt. The GSM amplitude is now $a \approx 1.1$, not 2. The flagged raster no longer shows thousands of kelvin of ground excess below 85 MHz. The calibrated 07-17 sunrise excess is about twice the predicted Sun and the size of the raw crossings, not several times larger. Each half of phase C now prefers the real sky for the daily term. With 20-min gain knots the ground term no longer rises several-fold; the fast gain instead takes up much of the sky's variation, a frequency collapses when its amplitude falls below 0.15 of its 3-h fit's, and a shared fast gain cannot be fitted jointly across the high band.

2026-10-23 Moved to geometry release `v0005` (memo 002): horizon profiles `horizon_profiles_v0005`, its transmitter, and pointing table v4, with every fit, product and validation rebuilt on them; the 91 m re-raise has its own profile. The empirical beam is still `empirical_raster_v0013`, fitted on release `v0004`.

2026-10-22 Heights from the mode table rebuilt on 2026-10-18 (memo 001), tested against box-air's lidar and IMU (§4.1), and pointing table v3. The 07-15 night and the 07-17 sunset were on the ground, not at 30 m, and there is no 2 m horizon profile, so every fit now uses only rows at 30, 87.5 or 91 m. Withdrawn: the 07-15 parked-night fit, the 07-17 dusk window, and the sunset burst's timing against the Sun going behind the ridge (the burst and its timing with the flare stand). Changed: the phase-C daily term no longer prefers the real sky over shifted skies; the low-band collapse of the fast-gain fit and the fast gain's relation to the RF switch's temperature both came with the ground rows and are gone; the impulsive events' rate is no longer the same with the Sun visible and behind the ridge, and one-variable controls (era, local hour, the Sun's height above the horizon) disagree, so their origin is open. The fixed-gain fits are now scored on their held-out bins, not on one.

2026-10-20 Added a fast gain shared across frequencies (§7.6), fitted jointly to convergence with collapsing training frequencies dropped by rule. It removed the collapse at most of the collapsing frequencies, and with a linear frequency dependence the high-band ground term changed little when the low band shared the gain. The ground-term table added its values where it rescued a frequency, marked, with injection errors.

2026-10-19 Moved every fit, product and validation to the extended terrain, DEM v0004 and `horizon_profiles_v0004`: `fit_v0013`–`v0016` and the `_v0004` products replace `fit_v0009`–`v0012` and the `_v0003` ones. No conclusion changes. The phase-C held-out comparisons move by a few hundredths of a point; the largest other moves are the joint-fit floor on $T_{\rm gnd}$ and the night fit's unconstrained $T_{\rm gnd}$ values, both of which the text already treats as unconstrained. Added the 30 m era's one usable calibration interval (§7.6): there too the gain after the RF switch stays within 1% while the fitted fast gain moves several per cent.

2026-10-18 Added the ground term with a fast gain (§7.6): over phase C the gain changes within the hour, injections show 3-h knots bias $T_{\rm gnd}$ low, and with 20-min knots it is quoted at 110–205 MHz with injection errors. The fit collapsed at several low- and high-band frequencies, and the fast gain followed the RF switch's temperature. Where measured (07-17, 87.5 m), the receiver's gain after the switch moves about 1% and does not follow the sky, so the fast gain is not that gain; §7.5 notes what this leaves open for

the daily term. Added the low-frequency daily term over phase C (§7.5): a slow additive term at 62–82 MHz that follows neither the pointing nor the recorded housekeeping. Diffuse reflection of the sky, and sky pickup, fit it over the whole phase. The sky origin and the reflection coefficient are withdrawn: the sidereal-time controls work only with the ground's level tied to R, and neither the preference nor R reproduces between 07-15 and 07-16. Faster gain knots (1 h, 20 min) take up the term into the gain, and a mock with a real term behaves the same way, so the term is stated as found with 3-h knots, with its additive nature open. Revised the 07-17 sunset at 30 m. The step in raw power there is mostly a solar burst during a C1.5 flare, ending as the disk goes behind the ridge before the elevation moves. Only the last 18% comes with the elevation step, which the earlier version blamed for all of it.

2026-10-16 Added the effect of extending the DEM 4 km south on the nominal horizon and on the predicted temperature (Section 2, Fig. 1), and the products that follow from it: DEM v0004 (6 km more to the south), `horizon_profiles_v0004` and its coverage check. No fit or product in this memo changed.

2026-10-07 Added the solar bursts, which identify solar emission in box-air; the raw context and raw-power fits of every terrain crossing; the local origin of the daytime impulsive events; the Sun's path through the beam and the raster test. The calibrated 07-17 excess is now read as larger than the Sun, not as the Sun. Added sky and ground recovery: the joint-frequency fit, the sky-calibrated parked nights, and the rotating-antenna regressions with their repeatable but frequency-dependent ground term. Traced the low-frequency excess: not specular reflection, horizon noise or localized sources, not a fixed fraction of the sky; the beam below the horizon or a tilt-dependent match remain, and need a field measurement.

2026-10-06 First version: the model, the degeneracy forecast, the night fit on `tcal@v0003` and `flags@v3`, and the first look for the Sun at the 07-17 terrain sunrise.

A Reproducing this memo

The products are regenerated, from `data-analysis/scripts/marjum-2026-07/ground_sky/` on the data-analysis branch `ground-sky-beam-v0017` (the products were built at b3cbff7; the degeneracy summary at 076e255), with `EIGSEP_CAMPAIGN_ROOT` set to the campaign directory, by the commands below. They read `pointing@v5` and take height eras from `curation/mode_table.jsonl` as rebuilt on 2026-10-18 (memo 001), by source file. Every product that depends on the antenna position reads the horizon profiles `curation/horizon_profiles_v0005` (geometry release `v0005`, memo 002, with its transmitter) and traces the Sun on the DEM mosaic `derived/dem/v0004`, and every product that rotates a beam uses the empirical beam and mount (with the axle tilt) of `empirical_raster_v0017`, fitted on release `v0004`. The fits calibrate the raw bins with `tcal@v0004`. The burst cutouts and the impulsive-event scan (`impulsive_v0004`, built on pointing v4 for its moving/parked state) are unchanged from the previous version.

- `python bin_raw.py v0006`;
- `python fit_ground_sky.py v0025 -binned raw_binned_v0006 -sun cut-flat`, `v0026 ... -sun cut-terrain` and `v0027 ... -sun model`;
- `python sun_events.py $EIGSEP_CAMPAIGN_ROOT/derived/ground_sky/fit_v0025 -out-dir $EIGSEP_CAMPAIGN_ROOT/derived/ground_sky/sun_events_v0007`;
- `python sun_bursts.py v0001`, `python impulsive_scan.py v0004`, `python sun_transitions.py v0007 -sun-events v0007 -fit v0025`, `python sun_profile.py v0007`, and `python raster_sun.py v0007 -beam hfss` then `-beam empirical`. The burst list they read is `curation/solar_bursts_v0001.json`.
- `python joint_fit.py v0007 -fit v0025`; `python sky_cal.py v0008 -beam hfss` and `-beam empirical`; `python raster_sky.py v0007` with `-beam hfss -label raster`, `-beam empirical -label raster -reflect`, and for the other moving windows `-beam empirical` with the `-label`, `-start`, `-stop`, `-era` (the 07-18 night: `91m-reraise`, which has its own profile) and `-row-step` recorded in each output (07-16 also `-self-emi`); and `python ground_excess.py v0007` with the `-split`, `-label` and window options recorded in each output; `python degeneracy_study.py`.

The low-frequency daily term (§7.5) comes from `python validate_phase_c.py -data-analysis <checkout> -work <dir>` in this directory, and `validate_phase_c_fastgain.py` uses the same checkout. Every validation

here takes a data-analysis checkout, checks that the commit it needs is an ancestor of it, and records the commit it ran at; the values below come from the commits named above, on the products listed there. In this version the validations ran on `ground-sky-beam-v0017` at 076e255, and the ground-term and shared-gain ones at 4db041a, which adds `-fast-collapse-frac` (the collapse rule relative to each frequency's 3-h fit); the heights and impulsive-event controls are unchanged. Commits from the branch `diffuse-ground` add the reflection, beam, pickup and housekeeping forms; the remaining options were added at c47da64 (`-ground-group`), fde5185 (`-ground-level-prior`), 6b10518 (`-t-range`), fc1dbb9 (`-night-only`, `-gain-segment-by-era`) and ae990d4 (`-t-exclude`). The 07-17 sunset burst comes from `python validate_sunset_burst.py -data-analysis <checkout>`, and the heights of §4.1 from `python validate_heights.py`, which reads every box-air raw file of 07-14 to 07-18 and records each one's SHA-256. Both record their inputs' checksums and the commit they ran at. The ground term with a fast gain (§7.6) comes from `python validate_ground_term.py -data-analysis <checkout> -work <dir>`. It runs `global_fit.py`, `inject_mock.py`, `hk_streams.py` and `gain_drivers.py` there, each pinned to at most five cores, and records the commit and the binned product's checksum. The receiver's own gain comes from `python validate_cal_gain.py -fit-checkout <dir> -cal-checkout <dir> -fit-work <dir> -cal-work <dir>`; it records both commits and the checksums of `raw_binned_v0006` and `tcal@v0004`. The 30 m-era load visits come from `python validate_cal_thirty.py -cal-checkout <dir> -fit-work <dir>`, run after `validate_ground_term.py` with the same work directory; it reads raw correlator files and records their SHA-256. The shared fast gain comes from `python validate_shared_gain.py -data-analysis <checkout> -fit-work <dir> -work <dir>`, run after `validate_ground_term.py` (whose per-frequency residuals its mocks use).

The independent night fit (§5.1) is not refitted here. Four of its figures and `nightfit_numbers.json` are outputs of the night fit of Christian Hellum Bye's EIGSEP D5 analysis (not shared), copied with its `results.json` (as `night_fit_results.json`) into `inputs/night_fit/`. The residual in kelvin (Fig. 4, bottom) is drawn from `night_fit_residual.npz`, which `python nightfit.py -extract <dir>` takes unchanged from the night fit's `fits.npz` and `data_v0025.npz` after checking their SHA-256. `nightfit.py`, called by `python make_products.py` (alone: `-only nightfit`), checks every input's SHA-256, copies the four figures, draws the fifth, and writes the method-chain table and the quoted numbers (Table 24). The night tilt's error, 0.6°, is memo 003's. The night-residual analysis's `results.json` and `PROVENANCE.json` (as `night_residual_results.json` and `night_residual_provenance.json`) and the data, model and residual of its two beam-swap fits (as `night_residual_beam_swap_native64_fit.npz` and `night_residual_beam_swap_v0018_fit.npz`) are copied into the same directory and checked the same way; `nightfit.py` draws its three figures from them (Figs. 8, 9 and 10) and quotes its numbers unchanged. The numbers Section 2 quotes on the team frame's error are copied from the first night fit of the same analysis into `sky_frame_effect.json` and checked the same way.

The site weather and ground temperature (§5.2) come the same way from the site-weather analysis: its two figures and `weather_numbers.json`, computed from the stations' published data, are copied into `inputs/weather/`; `weather.py`, called by `python make_products.py` (alone: `-only weather`), checks their SHA-256, copies the figures and writes the station table and the quoted numbers (Table 23).

Table 23: The inputs of the site weather and ground temperature (§5.2) and the commit they came from.

weather input (§5.2)	SHA-256 (first 16)
`memos/memo-008-marjum-2026-07-image-domain/inputs/weather/weather_air.pdf`	`271846815a2a11e2`
`memos/memo-008-marjum-2026-07-image-domain/inputs/weather/weather_numbers.json`	`0206cf64186703a1`
`memos/memo-008-marjum-2026-07-image-domain/inputs/weather/weather_soil.pdf`	`9f0ceb1a071a5b16`
site-weather analysis, source commit (not shared; run at `fef0e60`)	`343c27a`
eigsep (memo) commit, weather products	`38cd080`
generated	2026-11-06 13:53 UTC

The commits they ran at are below; each product's `manifest.json` records them along with its inputs' checksums. This memo's figures, tables and numbers come from `python make_products.py` in this directory, then `pdflatex memo-008.tex` twice. The script checks the SHA-256 of every product file it reads.

Table 24: The inputs of the independent night fit (§5.1) and the commits they came from.

night-fit input (§5.1)	SHA-256 (first 16)
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/imu_tilt_results.json	7b166e7934447134
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_fit_residual.npz	b135033311658c1f
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_fit_results.json	77f16cb3d9e422d0
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_residual_beam_swap_native64_fit.npz	98954f339df41cea
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_residual_beam_swap_v0018_fit.npz	b5a66eb60030444e
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_residual_provenance.json	2daf7ecda2e8e329
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/night_residual_results.json	4057c26d5a3826fd
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/nightfit_dmr.pdf	5debe49a66d725c1
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/nightfit_dmr_1d.pdf	2fbcb818a2dc139b
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/nightfit_heldout.pdf	7e53bbf905884667
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/nightfit_numbers.json	e175ed8cc34fa868
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/nightfit_params.pdf	846ac6e71a333968
memos/memo-008-marjum-2026-07-image-domain/inputs/night_fit/sky_frame_effect.json	89466ecbec4865da
night fit, source commit (not shared; run at 636ec64)	45563fa
night-residual analysis, source commit (not shared; run at e18a51c)	dad46d0
first night fit (the frame comparison), source commit (not shared)	81a5043
night tilt from the IMU tilt analysis (tau_box.night.wmean)	a27416d40ffc81c3
eigsep (memo) commit, night-fit products	6b4384b
generated	2026-11-07 00:06 UTC

input	SHA-256 (first 16)
derived/ground_sky/raw_binned_v0006/binned.npz	ebb0b559b141cc7a
derived/ground_sky/raw_binned_v0006/manifest.json	c28638551782ce85
derived/ground_sky/fit_v0025/fit.npz	272905d4b0a6884f
derived/ground_sky/fit_v0025/manifest.json	2c57254ffdc8b833
derived/ground_sky/fit_v0026/fit.npz	ae79f0e3e88d991b
derived/ground_sky/fit_v0026/manifest.json	4b385ed3ca84ec88
derived/ground_sky/fit_v0027/fit.npz	773dbad1e451c864
derived/ground_sky/fit_v0027/manifest.json	88d64ba42a5c358d
derived/ground_sky/sun_events_v0007/sun_events.json	50c5ab444e827f8c
derived/ground_sky/sun_events_v0007/sun_events.npz	e1add75a935b28da
derived/ground_sky/raw_binned_v0001/manifest.json	3ef90c83d3e3c0ad
derived/ground_sky/bursts_v0001/bursts.json	5e69db566e4dc2c0
derived/ground_sky/bursts_v0001/cutouts.npz	bca9c39857fd690e
derived/ground_sky/bursts_v0001/manifest.json	b9f52b6337f844e9
derived/ground_sky/impulsive_v0004/manifest.json	1aaa36233c75d496
derived/ground_sky/impulsive_v0004/events.jsonl	5eec1f86c51b1db0
derived/ground_sky/transitions_v0007/transitions.json	93c788bed577ee49
derived/ground_sky/transitions_v0007/transitions.npz	e9c37cd36f90e5d0
derived/ground_sky/transitions_v0007/manifest.json	a9e3b1fc94dc3033

derived/ground_sky/sun_profile_v0007/sun_profile.npz	37ff01764afa1760
derived/ground_sky/sun_profile_v0007/manifest.json	0cd82ec9912a2b09
derived/ground_sky/raster_sun_v0007/raster_hfss.json	290a66a8083f7dd9
derived/ground_sky/raster_sun_v0007/raster_empirical.json	15cd55fcc9798dac
derived/ground_sky/raster_sun_v0007/raster_empirical.npz	ba7524ba9b458041
derived/ground_sky/joint_v0007/joint_hfss.json	0991e8b459b8a1ca
derived/ground_sky/skycal_v0008/skycal_hfss.json	a0eac2c4c5961d2a
derived/ground_sky/skycal_v0008/skycal_empirical.json	cee148ff29870a7a
derived/ground_sky/skycal_v0008/skycal_hfss.npz	70d131b4c69cfec3
derived/ground_sky/raster_sky_v0007/raster_hfss.json	d6e2b6abc57da0ce
derived/ground_sky/ground_excess_v0007/raster_horizon_empirical.json	8df42472b5f08440
derived/ground_sky/ground_excess_v0007/raster_horizon_noflags_empirical.json	8fd0ca6708e99a98
derived/ground_sky/ground_excess_v0007/raster_azimuth_empirical.json	912c02744217e866
derived/ground_sky/ground_excess_v0007/w0718night91_horizon_empirical.json	71989ae12e0e45e6
derived/ground_sky/ground_excess_v0007/w0718night91_azimuth_empirical.json	230c4b540146bfa2
derived/ground_sky/ground_excess_v0007/w0717eve91_azimuth_empirical.json	c72f1005fbe6d274
derived/ground_sky/raster_sky_v0007/raster_empirical.json	09a2cc4c95df5a55
derived/ground_sky/raster_sky_v0007/w0717eve91_empirical.json	4af1f6c028d7355a
derived/ground_sky/raster_sky_v0007/w0718night91_empirical.json	9cc45b0192c1e053
derived/ground_sky/raster_sky_v0007/w0716night_empirical.json	242631d66a252aa4
marjum-2026-07/curation/horizon_profiles_v0005.npz	b497cebd1324291a
data-analysis/scripts/marjum-2026-07/ground_sky/summary.json	c229c3474fb6c1ef
memos/memo-008-marjum-2026-07-image-domain/inputs/solar_reference.json	e09895904559a8d0
validate_phase_c.json (this memo)	d3d7914d6d152c81
validate_sunset_burst.json (this memo)	945bbe8274eb9291
validate_sunset_burst.npz (this memo)	c198e636a7ec7140
validate_ground_term.json (this memo)	2e3c818acf0b635c
validate_cal_gain.json (this memo)	2ae94ba52b98df40
validate_cal_thirty.json (this memo)	7581a9837aa7b332
validate_shared_gain.json (this memo)	1be16dab0d4d5fed
validate_phase_c_fastgain.json (this memo)	d0b4e0e00b62a2f7
validate_heights.json (this memo)	d3abe5647327ae9a
validate_heights.npz (this memo)	287f7d7899b770ba
validate_impulsive.json (this memo)	df289264d24e6655
products made with data-analysis	347cecc, 72b9943, b3cbff7
products made with eigsep_base	840c10b
products made with eigsep_data	9ac9b93, e1da3c5, e941c51
products made with eigsep_sim	d050c83
phase-C fits: data-analysis ground-sky-beam-v0017	076e255
fast-gain fits: data-analysis ground-sky-beam-v0017	4db041a
shared fast gain: data-analysis ground-sky-beam-v0017	4db041a
30 m receiver gain: data-analysis ground-sky-beam-v0017	076e255
receiver-gain fits: data-analysis ground-sky-beam-v0017	076e255
eigsep (memo) commit	bc309e0
data-analysis commit (degeneracy study file)	4db041a
generated	2026-11-05 08:42 UTC