

Design and Analysis of Experiments

04 - Statistical Intervals

Version 2.12.2018

Felipe Campelo

<http://orcslab.cpdee.ufmg.br/>

Graduate Program in Electrical Engineering

Belo Horizonte
April 2018

*"I attribute my success to this: I never
gave or took an excuse."*

*"I think one's feelings waste themselves in words;
they ought all to be distilled into actions,
and into actions which bring results."*

Florence Nightingale
1820-1910

English statistician and founder of modern nursing



Statistical Intervals

Introduction

Statistical intervals are important in quantifying the uncertainty associated to a given estimate;

As an example, let's recap the coaxial cables example: *a coaxial cable manufacturing operation produces cables with a target resistance of 50Ω and a standard deviation of 2Ω . Assume that the resistance values can be well modeled by a normal distribution.*

Let us now suppose that a sample mean of $n = 25$ observations of resistance yields $\bar{x} = 48$. Given the sampling variability, it is very likely that this value is not exactly the true value of μ , but we are so far unable quantify how much uncertainty there is in this estimate.

Statistical Intervals

Definition

Statistical intervals define regions that are likely to contain the true value of an estimated parameter.

More formally, it is generally possible to quantify the level of uncertainty associated with the estimation, thereby allowing the derivation of sound conclusions at predefined levels of certainty.

Three of the most common types of interval are:

- Confidence intervals;
- Tolerance intervals;
- Prediction intervals;

Statistical Intervals

Definition

Statistical intervals define regions that are likely to contain the true value of an estimated parameter.

More formally, it is generally possible to quantify the level of uncertainty associated with the estimation, thereby allowing the derivation of sound conclusions at predefined levels of certainty.

Three of the most common types of interval are:

- **Confidence intervals;**
- Tolerance intervals;
- Prediction intervals;

Confidence Intervals

Definition


Confidence intervals quantify the degree of uncertainty associated with the estimation of population parameters such as the mean or the variance.

Can be defined as “*the interval that contains the true value of a given population parameter with a confidence level of $100(1 - \alpha)\%$* ”;

Another useful definition is to think about confidence intervals in terms of confidence *in the method*: “The method used to derive the interval has a hit rate of 95%” - i.e., the interval generated has a 95% chance of ‘capturing’ the true population parameter.”

Confidence Intervals

Example: 100 $CI_{.95}$ for a sample of 25 observations



For an interactive demonstration of the factors involved in the definition of a confidence interval, download the files from <https://git.io/vxXGj> and run on RStudio.

Confidence Intervals

CI on the Mean of a Normal Variable

The two-sided $CI_{(1-\alpha)}$ for the mean of a normal population with known variance σ^2 is given by:

$$\bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{1-\alpha/2} \frac{\sigma}{\sqrt{n}}$$

where $(1 - \alpha)$ is the confidence level and z_x is the x -quantile of the standard normal distribution.

For the more usual case with an unknown variance,

$$\bar{x} + t_{\alpha/2}^{(n-1)} \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t_{1-\alpha/2}^{(n-1)} \frac{s}{\sqrt{n}}$$

where $t_x^{(n-1)}$ is the x -quantile of the t distribution with $n - 1$ degrees of freedom.

Confidence Intervals

CI on the Variance and Standard Deviation of a Normal Variable

A two-sided confidence interval on the variance of a normal variable can be easily calculated:

$$\frac{(n-1)s^2}{\chi^2_{1-\alpha/2}(n-1)} \leq \sigma^2 \leq \frac{(n-1)s^2}{\chi^2_{\alpha/2}(n-1)}$$

where $\chi^2_x(n-1)$ represents the x-quantile of the χ^2 distribution with $n-1$ degrees of freedom. For the standard deviation one simply needs to take the squared root of the confidence limits.

Statistical Intervals

Wrapping up

Statistical intervals quantify the uncertainty associated with different aspects of estimation;

Reporting intervals is always better than point estimates, as it provides the necessary information to quantify the location and uncertainty of your estimated values;

The correct interpretation is a little tricky (although not very difficult)^[3], but it is essential in order to derive the correct conclusions based on the statistical interval of interest.

[3] See the table at the end of <https://git.io/v5ZFh>

Bibliography

Required reading

- 1 J.G. Ramírez, *Statistical Intervals: Confidence, Prediction, Enclosure*:
<https://git.io/v5ZFh>
- 2 D.C. Montgomery and G.C. Runger, *Applied Statistics and Probability for Engineers*, Chapter 8. 3rd Ed., Wiley 2005.
- 3 J. Orloff and J. Bloom, *Bootstrap confidence intervals*: <https://goo.gl/XrT1ao>

Recommended reading

- 1 Simply Statistics (blog) - <http://simplystatistics.org>
- 2 R. Dawkins, *Climbing Mount Improbable*, W.W.Norton&Co.,1997.

About this material

Conditions of use and referencing

This work is licensed under the Creative Commons CC BY-NC-SA 4.0 license (Attribution Non-Commercial Share Alike International License version 4.0).

<http://creativecommons.org/licenses/by-nc-sa/4.0/>

Please reference this work as:

Felipe Campelo (2018), *Lecture Notes on Design and Analysis of Experiments*.

Online: <https://github.com/fcampelo/Design-and-Analysis-of-Experiments>
Version 2.12. Creative Commons BY-NC-SA 4.0.

```
@Misc{Campelo2018,  
  title={Lecture Notes on Design and Analysis of Experiments},  
  author={Felipe Campelo},  
  howPublished={\url{https://github.com/fcampelo/Design-and-Analysis-of-Experiments}},  
  year={2018},  
  note={Version 2.12. Creative Commons BY-NC-SA 4.0.},  
}
```

