

# Population models of *Minecraft* sheep

Evan Freeland

January 17, 2024

## 1 Introduction

*Minecraft* is an open-world sandbox video game<sup>1</sup>. The game takes place in a procedurally generated world made of cubic voxels, or *blocks*. Among other activities, players can engage in agriculture, including the raising of sheep.

Players can gather blocks of wool by raising a flock of sheep in a pasture (Figure 1). When a sheep is sheared, they must eat grass to regrow their wool. When a grass block is eaten, it converts into dirt, and can regrow if grass spreads from an adjacent grass block.

One may be tempted to pack the pasture as densely as possible to maximize space efficiency. However, if a pasture is packed too densely with sheep, they will deplete the grass, and wool production will be low. On the other hand, if the pasture is not densely packed, wool production will still be low because there are not many sheep. To understand where the happy medium lies, we will develop mathematical models for the populations of sheep and grass in the pasture.

### 1.1 Deterministic vs. Stochastic

Models of biological populations can be broadly divided into *deterministic* and *stochastic*. Deterministic models are simpler to analyze, but stochastic models are generally more informative since the biological populations, both in the real world and in *Minecraft*, are stochastic processes. In the real world, populations are influenced by random ecological processes, and in *Minecraft*, the growth of grass and the appetite of sheep are governed by random number generation.

We will begin with a deterministic approach to modelling a pasture, and then move to a stochastic model.

---

<sup>1</sup>This investigation was conducted in *Minecraft* Java edition version 1.19.4.

Figure 1: A pasture in a meadow, with woolly and sheared sheep, and grass and dirt blocks.



## 2 Deterministic model

### 2.1 Grass population

The *Minecraft* physics engine operates in discrete time increments of 0.05 seconds, called *ticks*. Since the increments are small compared to the expected time between sheep eating and grass block growing, we will employ a continuous-time model for this discrete-time process.

Every tick, a sheep has a chance to attempt to eat the block it's standing on. The attempt will only succeed if the sheep is standing on grass. If the sheep and grass in the pasture are uniformly distributed, the expected number of sheep standing on grass is the product of the number of sheep and the fraction of grass blocks in the pasture. The rate at which grass is eaten is therefore given by:

$$\mu = \frac{NRN_g}{A} \quad (1)$$

Where  $\mu$  is the rate at which grass is eaten,  $A$  is the size of the pasture,  $N_g$  is the number of grass blocks in the pasture,  $N$  is the number of sheep, and  $R$  is the frequency with which an individual sheep attempts to eat.

Every tick, a grass block has a chance to attempt to spread to an adjacent dirt block. If  $N_g$  is small, then grass growth will be limited by the lack of places the grass can spread *from*. If  $N_g$  is large, grass growth will be limited by a lack of places for the grass to spread *to*. We need some equation such that the grass growth rate,  $\lambda$ , is low for extreme values of  $N_g$  and higher for moderate values. Let's assume  $\lambda$  has a parabolic dependence on  $N_g$ :

$$\lambda = kN_g - jN_g^2$$

The growth rate must be zero for  $N_g = 0$  and  $N_g = A$ , which requires  $j = \frac{k}{A}$ :

$$\lambda = kN_g - \frac{k}{A}N_g^2 \quad (2)$$

This leaves  $k$  as a constant related to the rate at which grass spreads.  
The overall rate of change of  $N_g$  is the difference of  $\lambda$  and  $\mu$ :

$$\dot{N}_g = \lambda - \mu = \left(k - \frac{NR}{A}\right)N_g - \frac{k}{A}N_g^2 \quad (3)$$

This model, in which  $\dot{N}_g$  has quadratic dependence on  $N_g$ , is known as the *logistic* model. This model is one of the oldest models for population growth, originally proposed in 1838 [4]. It has since been joined by many more general models [5] which are more useful for real-world ecology, but since we are modelling a video game, we will employ it for its simplicity.

The exact solution [5] of equation 3 is:

$$N_g(t) = \frac{P}{1 + me^{-at}} \quad (4)$$

Where

$$a = \left(k - \frac{NR}{A}\right) \quad P = A - \frac{NR}{k} \quad m = \frac{P}{N_g(0)} - 1$$

Of note here is that the population asymptotically approaches  $P$  as  $t \rightarrow \infty$ . In ecology,  $P$  is known as the carrying capacity, the maximum population that a habitat can sustain. We will refer to  $P$  as the equilibrium grass population, to avoid confusion with  $A$ , which is the number of blocks in the pasture and therefore the maximum number of grass blocks it can contain.

We should also note that for  $N \geq \frac{Ak}{R}$ , the equilibrium grass population is 0. According to the logistic model, putting this many sheep in a pasture will guarantee that they eat all the grass.

## 2.2 Experimental determination of rate constants

Before proceeding further, it's worth mentioning the actual values of  $k$  and  $R$ . These were measured as  $0.00888 \text{ s}^{-1}$  and  $0.00772 \text{ s}^{-1}$ , respectively. More details about the experiments used to obtain these values can be found on the author's Github<sup>2</sup>.

## 2.3 Woolly sheep population

For simplicity, let's imagine the farmer has superhuman speed, and is able to shear each sheep the instant it grows its wool. In this case, the rate at which

---

<sup>2</sup>github.com/

wool is gathered is equal to  $\mu^3$ . Denoting the rate of wool gathering with  $F$ , and substituting  $N_g = P$  into equation 1 (assuming the pasture has reached equilibrium):

$$F_{fast} = \mu = NR - \frac{N^2 R^2}{Ak} \quad (5)$$

This is a simple parabolic relationship with a maximum value of  $F_{fast} = \frac{Ak}{4}$  at  $N = \frac{Ak}{2R}$ .

*Minecraft* players typically don't want to put all of their time into shearing their sheep as quickly as they can. For a more realistic model, we'll need to consider how the population of woolly sheep evolves. A sheep will only regrow wool upon eating if it doesn't already have any, so we should expect that the rate at which sheep are regrowing wool is the rate at which they are eating,  $\mu$ , times the fraction of sheep that are wool-less. Denoting  $N_w(t)$  as the number of wooly sheep in the pasture:

$$\dot{N}_w = \frac{dN_w}{dt} = \frac{N - N_w}{N} \mu = (N - N_w) \frac{RN_g}{A} \quad (6)$$

Let's assume the farmer shears all the sheep at time  $t = 0$ , leaving  $N_w = 0$  and then goes off to do other things. At time  $T$ , they come back to shear the sheep again. The average rate at which the farmer gathers wool is:

$$F_{slow} = \frac{N_w(T)}{T}$$

How many sheep regrew their wool in time  $T$ ? Equation 6 is a simple linear ODE, and we can solve it for the initial condition of  $N_w(0) = 0$ , and assuming  $N_g = P$  (and is therefore time-independent):

$$N_w(T) = N - Ne^{-TR + \frac{TR^2 N}{Ak}}$$

Therefore:

$$F_{slow} = \frac{N - Ne^{-TR + \frac{TR^2 N}{Ak}}}{T} \quad (7)$$

To find the optimal sheep population, we can differentiate equation 7 w.r.t.  $N$  and set  $F_{slow} = 0$ :

$$\frac{dF_{slow}}{dN} = \frac{1 - \left( \frac{TR^2 N}{Ak} + 1 \right) e^{-TR + \frac{TR^2 N}{Ak}}}{T} = 0$$


Solving for  $N$  yields:

$$N = \frac{Ak}{TR^2} (W(e^{TR+1}) - 1) \quad (8)$$

---

<sup>3</sup>Shearing a sheep actually yields between 1-3 blocks of wool, but this is dictated purely by chance, so we'll assume one sheep = one wool.

Figure 2: Contours of  $F_{slow}$  as a function of  $N$  at constant  $T$ . The dashed line traces the optimal  $N$  and  $F_{slow}$  as  $T$  increases.



Where  $W$  is the Lambert  $W$  function, which satisfies  $W(xe^x) = x$ . Plugging this back into equation 7, we obtain the value of  $F_{slow}$  at optimal  $N$ :

$$F_{slow} = \frac{Ak(W(e^{TR+1}) - 1)(1 - e^{-TR+W(e^{TR+1})-1})}{R^2T^2} \quad (9)$$

Figure 2 shows a plot of equations 7-9. As  $T$  increases, the optimal sheep population also increases, and the optimal rate of wool gathering decreases. Clearly there is a tradeoff between the frequency with which the farmer shears the flock and the rate at which they gather wool.

We should expect that for small  $T$ ,  $F_{slow} \approx F_{fast}$ . This is already qualitatively evident from the parabolic shape of the  $T = 1$  curve in Figure 2, which matches the quadratic relationship between  $N$  and  $F_{fast}$  in equation 5. We can show this mathematically by applying the following approximation, for  $x \ll 1$ :

$$e^x \approx x + 1$$

In equation 7, the exponent is small for small values of  $T$ , therefore:

$$F_{slow} \approx \frac{N - N(-TR + \frac{TR^2N}{Ak} + 1)}{T} = NR - \frac{N^2R^2}{Ak} = F_{fast}$$

It appears in Figure 2 that for large  $T$  and small  $N$ ,  $F_{slow}$  is approximately linear w.r.t.  $N$ . Under these conditions, all of the sheep have regrown their wool by the time the farmer comes back, and  $N_w \approx N$ . The exponent  $-TR + \frac{TR^2N}{Ak}$  in equation 7 is large and negative, and:

$$F_{slow} \approx \frac{N - 0}{T} = \frac{N}{T}$$

Lines of  $\frac{N}{T}$  are plotted over top of the curves in Figure 2.

### 3 Stochastic model

#### 3.1 Grass population

In the deterministic model, we considered the grass population as a single number,  $N_g$ , which evolved with time. In the stochastic approach, we will consider the evolution of the *probability distribution* of  $N_g$  with time. The method used here is described in [1], and applied here for the case of the logistic model.

Consider a pasture at time  $t = t_0$ , with  $N_g(t_0) = x$ . A short time ago, at time  $t = t_0 - \Delta t$ , the pasture had  $N_g(t_0 - \Delta t)$ . The pasture may have arrived at  $N_g(t_0) = x$  in one of four ways:

1.  $N_g(t_0 - \Delta t) = x$ , and neither eating nor growth take place.
2.  $N_g(t_0 - \Delta t) = x - 1$ , and one grass block regrows.
3.  $N_g(t_0 - \Delta t) = x + 1$ , and one sheep eats a grass block.
4. Two or more growth and eating events occur.

The probability of each pathway is the probability that the pasture was in the required state  $N_g(t_0 - \Delta t)$  times the probability of the event (eating or growth) required to bring the pasture to  $N_g(t_0) = x$ .

$\lambda$  and  $\mu$  are the expected number of grass blocks that regrow in unit time, and the expected number of sheep that eat in unit time, respectively. For a sufficiently short time interval  $\Delta t$ ,  $\lambda\Delta t$  and  $\mu\Delta t$  are the probability that one grass block regrows or one sheep eats in time  $\Delta t$ , respectively. This gives the probability of nothing happening in time  $\Delta t$  as  $(1 - \lambda\Delta t - \mu\Delta t)$ . The probability of two or more events is some product of  $\lambda\Delta t$  and  $\mu\Delta t$ , and is therefore dependent on  $\Delta t$  with order 2 or higher.

We'll introduce the following notation for the state of the pasture at time  $t$ :

$$p_x(t) = \Pr(N_g(t) = x)$$

The probability of each path is therefore:

$$\begin{aligned} \Pr(1) &= p_x(t_0 - \Delta t)(1 - \lambda\Delta t - \mu\Delta t) \\ \Pr(2) &= p_{x-1}(t_0 - \Delta t)\lambda\Delta t \\ \Pr(3) &= p_{x+1}(t_0 - \Delta t)\mu\Delta t \\ \Pr(4) &\approx O(\Delta t^2) \end{aligned}$$

The probability of the pasture arriving at  $N_g(t_0) = x$  is the sum of the probability of each pathway:

$$\begin{aligned} p_x(t_0) &= p_x(t_0 - \Delta t)(1 - \lambda\Delta t - \mu\Delta t) \\ &\quad + p_{x-1}(t_0 - \Delta t)\lambda\Delta t \\ &\quad + p_{x+1}(t_0 - \Delta t)\mu\Delta t \\ &\quad + O(\Delta t^2) \end{aligned}$$

subtracting  $p_x(t_0 - \Delta t)$ , dividing by  $\Delta t$ , and taking  $\lim_{\Delta t \rightarrow 0}$  converts this into a differential equation:

$$\dot{p}_x(t) = -p_x(t)(\lambda(x) + \mu(x)) + p_{x-1}(t)\lambda(x-1) + p_{x+1}(t)\mu(x+1) \quad (10)$$

Note that  $\mu(x)$  simply represents the value of  $\mu$  with  $N_g = x$ , and likewise for  $\lambda$ . Note also that the  $O(\Delta t^2)$  term has vanished upon taking the limit  $\Delta t \rightarrow 0$ .

We will have one copy of equation 10 for every possible value of  $x$ , i.e. we will have a system of  $A + 1$  equations. Systems of equations such as these, which describe the evolution of a probability distribution with time, are known as *Kolmogorov equations* [1]. In general, Kolmogorov equations can be used to describe a *Markov process*, i.e. a process by which a system transitions between multiple possible states. In our case the possible states are the possible values of  $N_g$ .

We'll wrap up the system into vector form:  $\dot{\mathbf{p}}(t) = \mathbf{p}(t)\mathbf{R}$ , where  $\mathbf{p}(t) = [p_0(t), p_1(t), p_2(t) \dots p_A(t)]$ , and  $\mathbf{R}$  is a matrix of coefficients, such that the coefficient  $r_{i,j}$  relates  $p_i$  with  $\dot{p}_j$ :

$$\dot{p}_j = \sum_i r_{ij} p_i$$

$$r_{i,j} = \begin{cases} \lambda(j-1) & \text{for } i = j-1 \\ \mu(j+1) & \text{for } i = j+1 \\ -(\lambda(i) + \mu(i)) & \text{for } i = j \\ 0 & \text{otherwise} \end{cases}$$

Then:

$$\dot{\mathbf{p}}(t) = \mathbf{p}(t)\mathbf{R}$$

This system can now be solved numerically. We'll solve it for a range of  $N$  and three different pasture sizes. *Minecraft* pastures typically begin fully covered with grass, so we will assume  $\mathbf{p}(0) = [0, 0, 0 \dots 1]$ .

Let's consider the equilibrium expected value of  $N_g$ . Figure 3 compares the expected value of  $N_g$  after 10000 seconds with  $P$  from the deterministic logistic model. They line up very closely for small sheep densities, but for large sheep densities, the expected  $N_g$  is significantly less than  $P$ , especially for  $A = 100$ .

Let's look in a little more detail at the behavior of the pasture. Figure 4 shows the evolution of the distribution of  $N_g$  for  $N = 40$  and  $N = 70$  in a 100-block pasture. The pasture with  $N = 40$  approaches a stable distribution, evidenced by the indistinguishable distributions at  $t = 1600$  s and  $t = 9500$  s. In contrast, the pasture with  $N = 70$  continues evolving toward smaller and smaller grass populations, with  $p_0$  increasing dramatically as time proceeds.

The reason for this isn't hard to see by considering the equation for  $\dot{p}_0$ . Beginning with equation 10 and considering that  $p_{-1} = 0$  and  $\lambda(0) = \mu(0) = 0$ :

$$\dot{p}_0(t) = p_1(t)\mu(1)$$

Figure 3: Expected value of  $N_g$  at  $t = 10000$ , for  $A = 100$  and  $A = 500$ .  $P$  from the deterministic model is shown as a dotted line.





Figure 4: Evolution of  $N_g$  distribution for a pasture of size  $A = 100$ , for (a)  $N = 40$  and (b)  $N = 70$ .





Since  $p_1$  and  $\mu(1)$  are always positive,  $p_0$  is monotonically increasing, bounded above by  $p_0 = 1$ . Therefore the true equilibrium state is  $\mathbf{p} = [1, 0, 0, 0, \dots]$  i.e. extinction. However, even if this is the equilibrium state, the pasture may remain at a different *metastable* (or *quasi-equilibrium*) distribution if the probability of extinction remains negligible across reasonable time scales. This is what is seen in Figure 4a. The distribution for  $N = 40$  "stabilizes" with a value of  $p_1$  so low that  $\dot{p}_0$  remains negligible.

Rapid grass extinction at high sheep densities poses a problem, since the deterministic model predicts that the optimal  $N$  increases as  $T$  increases (Figure 2). Fortunately, an exact solution can be obtained for the expected time to extinction ETE. Let  $\mathbf{M}$  be a matrix of mean residence times, such that the element  $m_{ij}$  of  $\mathbf{M}$  is the expected amount of time the pasture spends with  $N_g = j$  before reaching extinction, given that it began with a population  $i$ . It can be shown [2] that  $\mathbf{M}$  is obtained by removing the first row and column from  $\mathbf{R}$  and taking the negative of the inverse of this modified  $\mathbf{R}$  matrix.

The sum of the  $i$ th row of  $\mathbf{M}$  gives the total time spent with nonzero population before becoming extinct, given the pasture started with  $N_g = i$ . The typical *Minecraft* pasture begins fully populated with grass, i.e.  $N_g(0) = A$ . Therefore the value we are interested in is:

$$\text{ETE} = \sum_j m_{Aj}$$

We can apply this solution to obtain the ETE for any value of  $N$  and  $A$ . Figure 5 shows the ETE for pastures of a few select sizes, as a function of  $N$ . For populations close to the deterministic sheep population limit  $\frac{N}{A} = \frac{k}{R} = 0.869$ , the ETE is on the order of a few hours for all three pasture sizes. As the sheep population density decreases, the ETE increases dramatically - For a 100 block pasture with a sheep density of 0.5, the ETE is over 2 years. ETE also increases dramatically with larger pastures - A 100-block pasture with sheep density 0.7 has ETE of 3 hours, while a 500-block pasture with the same density has ETE of 9 months. The variation of ETE with pasture size explains the closer match of the  $N = 500$  pasture with the deterministic model in Figure 3.


We have seen that pastures with long ETE, such as the one depicted in Figure 4a, approach a metastable distribution as they evolve. This distribution can actually be calculated analytically. The pasture will cease evolving when all  $\dot{p}_x = 0$ . This implies that the likelihood of the pasture having  $N_g = x$  and losing one grass block equals the probability of  $N_g = x - 1$  and growing one grass block [3]. In symbols:

$$p_x \mu(x) = p_{x-1} \lambda(x-1) \quad (11)$$

We will assume  $N_g = 0$  is an unreachable state and exclude it from consideration. Writing equation 11 for  $2 \leq x \leq A$  and combining with the normalization condition, we obtain a system of  $A$  equations in  $A$  variables, which can be solved to obtain the metastable  $N_g$ .

Figure 6 shows the metastable distribution for a few select conditions (it is assumed that  $p_0 = 0$ ). The metastable distribution predicted by equation

Figure 5: Expected time to extinction for pastures of  $A = 100, 200$ , and  $500$  as a function of sheep population density.



11 closely matches the numerical solution of the Kolmogorov equations for  $t = 10000$  in all cases except  $A = 100, N = 65$ . In that case, the ETE is not sufficiently long to allow a metastable distribution to exist. Equation 11 also produces unrealistic results in this case - the distribution shows an increase in  $p_x$  for small  $x$ ; such a distribution would not actually be metastable.

### 3.2 Woolly sheep population

A model for both the sheep and grass mechanics will have two variables evolving at the same time,  $N_g$  and  $N_w$ . Therefore, we will need to consider a bivariate probability distribution.

We'll use the following notation for concision:


$$N_{gw}(t) = [N_g(t), N_w(t)]$$

$$q_{xy}(t) = \Pr(N_{gw}(t) = [x, y])$$

We'll follow the same derivation as we did in the previous section. Consider a pasture at time  $t_0$ , with  $N_c(t_0) = [x, y]$ . A short time ago, at time  $t = t_0 - \Delta t$ , the pasture had  $N_c(t_0 - \Delta t)$ . The pasture may have arrived at the state at  $t_0$  in one of several ways:

- $N_{gw}(t_0 - \Delta t) = [x, y]$ , and neither eating nor growth take place.
- $N_{gw}(t_0 - \Delta t) = [x - 1, y]$ , and one grass block regrows.
- $N_{gw}(t_0 - \Delta t) = [x + 1, y]$ , and one woolly sheep eats grass.
- $N_{gw}(t_0 - \Delta t) = [x + 1, y - 1]$ , and one woolless sheep eats grass.

Figure 6: Metastable population distributions for pastures of size (a)  $A = 100$  and (b)  $A = 500$ , for three different sheep population densities. Numerical results are shown as dotted lines.



- Two or more growth and eating events occur.


As with the grass model, the probability of each path is the probability of the pasture existing at the required initial state, times the probability of the required change occurring. For the two cases involving one block of grass being eaten, the probability of the event will include a term pertaining to the number of sheep who have their wool. By analogy with equation 6, the rate at which woolless sheep eat grass is  $\frac{N-N_{gw}}{N}\mu$ , and the rate at which woolly sheep eat grass is  $\frac{N_{gw}}{N}\mu$ . The probability of the pasture arriving at  $N_{gw} = [x, y]$  is the sum of the probabilities of each path:

$$\begin{aligned}
 q_{x,y}(t) = & q_{x,y}(t - \Delta t)(1 - \mu(x)\Delta t - \lambda(x)\Delta t) \\
 & + q_{x-1,y}(t - \Delta t)\lambda(x-1)\Delta t \\
 & + q_{x+1,y}(t - \Delta t)\mu(x+1)\frac{y}{N}\Delta t \\
 & + q_{x+1,y-1}(t - \Delta t)\mu(x+1)\frac{N-(y-1)}{N}\Delta t \\
 & + O(\Delta t^2)
 \end{aligned}$$

Subtracting  $q_{x,y}(t - \Delta t)$ , dividing by  $\Delta t$ , and taking the limit  $\Delta t \rightarrow 0$ :

$$\begin{aligned}
 \dot{q}_{x,y}(t) = & q_{x,y}(t)(-\mu(x) - \lambda(x)) \\
 & + q_{x-1,y}(t)\lambda(x-1) \\
 & + q_{x+1,y}(t)\mu(x+1)\frac{y}{N} \\
 & + q_{x+1,y-1}(t)\mu(x+1)\frac{N-(y-1)}{N}
 \end{aligned} \tag{12}$$

Figure 7:  $F_{slow}$  calculated from the stochastic model for sheep and grass, for a pasture with  $A = 100$ . The initial state was obtained from the numerical solution of  $p_x$  at  $t = 10000$ .



We will have a system of  $(A + 1)(N + 1)$  equations. We can express this in matrix form as  $\dot{\mathbf{q}}(t) = \mathbf{q}(t)\mathbf{R}$ , where  $\mathbf{q}(t)$  is a matrix of elements  $q_{ij} = q_{xy}(t)$ , and  $\mathbf{R}$  is a rank-4 tensor, where element  $r_{ijkl}$  relates  $q_{ij}$  with  $\dot{q}_{kl}$ :

$$\dot{q}_{kl} = \sum_{i,j} q_{ij} r_{ijkl}$$

$$r_{ijkl} = \begin{cases} -\mu(i) - \lambda(i) & \text{for } (i, j) = (k, l) \\ \lambda(k-1) & \text{for } (i, j) = (k-1, l) \\ \mu(k+1) \frac{l}{N} & \text{for } (i, j) = (k+1, l) \\ \mu(k+1) \frac{N-(l-1)}{N} & \text{for } (i, j) = (k+1, l-1) \\ 0 & \text{otherwise} \end{cases}$$

As before, this system can be solved numerically. Let's assume the pasture has been equilibrating for 10000 seconds, and then the farmer shears the sheep, leaving  $N_w = 0$ .  $T$  seconds later, they come back again. We can set the initial values of  $q_{x,0}$  using the values of  $p_x(t = 10000)$  from the grass population model. 7 shows  $F_{slow}$  as calculated from the results of this model. Similarly to Figure 3, the stochastic model predicts significantly lower  $F_{slow}$  than the deterministic model.

The root cause of this deviation is the same as for the grass growth model. The equation for  $\dot{q}_{0,y}$  is:

$$\begin{aligned} \dot{q}_{0,y}(t) &= q_{1,y}(t) \mu(1) \frac{y}{N} \\ &= q_{1,y-1}(t) \mu(1) \frac{N - (y-1)}{N} \end{aligned}$$

Since all terms here are positive,  $q_{0,y}$  are monotonically increasing, and given enough time, the pasture will tend to evolve into one of these states and never leave. It's important to note that equation 12 can be summed across all  $y$  to obtain equation 10:

$$\begin{aligned}\sum_y \dot{q}_{x,y}(t) &= \left[ \sum_y q_{x,y}(t) \right] (-\mu(x) - \lambda(x)) \\ &+ \left[ \sum_y q_{x-1,y}(t) \right] \lambda(x-1) \\ &+ \left[ \sum_y q_{x+1,y}(t) \frac{y}{N} \right] \mu(x+1) \\ &+ \left[ \sum_y q_{x+1,y-1}(t) \frac{N-(y-1)}{N} \right] \mu(x+1)\end{aligned}$$

The last two sums result in:

$$\begin{aligned}\sum_y q_{x+1,y}(t) \frac{y}{N} &= q_{x+1,1} \frac{1}{N} + q_{x+1,2} \frac{2}{N} + \cdots + q_{x+1,N} \frac{N}{N} \\ \sum_y q_{x+1,y-1}(t) \frac{N-(y-1)}{N} &= q_{x+1,0} \frac{N}{N} + q_{x+1,1} \frac{N-1}{N} + \cdots + q_{x+1,N-1} \frac{1}{N}\end{aligned}$$

Each term combines with another in the other sum to give  $q_{x+1,y} \frac{N}{N}$ . Therefore we can combine the two sums into one  $\sum_y q_{x+1,y}$ :

$$\begin{aligned}\sum_y \dot{q}_{x,y}(t) &= \left[ \sum_y q_{x,y}(t) \right] (-\mu(x) - \lambda(x)) \\ &+ \left[ \sum_y q_{x-1,y}(t) \right] \lambda(x-1) \\ &+ \left[ \sum_y q_{x+1,y}(t) \right] \mu(x+1)\end{aligned}$$

Since  $\sum_y q_{x,y} = p_x$ , this is identical to equation 10. The fact that summing across  $y$ , i.e. removing consideration of wooly sheep, reduces this model to the model for grass population, means the ETE for a pasture with this model is the same as for a pasture .

The biggest drawback of this model is its computational intensiveness. If we want to apply this analysis to the same conditions as the model for grass population, we will have at most  $(500+1)(425+1) = 213426$  simultaneous equations, and an  $\mathbf{R}$  tensor with  $(500+1)^2(425+1)^2 = 45550657476$  elements.

Such a matrix would occupy 364 GB of memory if stored as 64-bit floats, making computation infeasible for all but top-shelf consumer hardware.

## 4 What next?

The logistic model, and the Kolmogorov equations, have proven useful for predicting the behavior of a pasture in *Minecraft*. However, there are still a few unanswered questions.

First, a pasture in *Minecraft* is often constructed by fencing off an area of grass from a larger meadow (see Figure 1). Grass can spread under fences, effectively introducing an immigration term to the logistic model that is nonzero even when the grass in the pasture is extinct. Such a model is studied in [1] chapter 7, but the deterministic model becomes rather unwieldy, so the effect of immigration was ignored here. However, introducing immigration would have an important effect on the behavior of the pasture for small  $N_g$ , since  $N_g = 0$  is no longer an absorbing state.

Second, time in *Minecraft* is discretized into increments of 0.05 seconds, while we have been modelling the pasture with continuous-time processes. How would a discrete-time model differ from a continuous-time model?

## References

- [1] Matis, James, and Thomas Kiffe, eds. *Stochastic Population Models: A Compartmental Perspective*. 1st ed. Lecture Notes in Statistics. New York: Springer, n.d.
- [2] Matis, J. H., T. E. Wehrly, and C. M. Metzler. “On Some Stochastic Formulations and Related Statistical Moments of Pharmacokinetic Models.” *Journal of Pharmacokinetics and Biopharmaceutics* 11, no. 1 (1983): 77-92. <https://doi.org/10.1007/BF01061769>.
- [3] Bartlett, M. S., J. C. Gower, and P. H. Leslie. “A Comparison of Theoretical and Empirical Results for Some Stochastic Population Models.” *Biometrika* 47, no. 1-2 (June 1, 1960): 1-11. <https://doi.org/10.1093/biomet/47.1-2.1>.
- [4] Verhulst, P. “Notice sur la loi que la population suit dans son accroissement.” *Correspondance Mathématique et Physique* 10 (1838): 113-21.
- [5] Banks, Robert B. *Growth and Diffusion Phenomena*. Texts in Applied Mathematics. Springer Berlin, Heidelberg, 1994.