

SULIT



UTM
UNIVERSITI TEKNOLOGI MALAYSIA

Faculty of
Computing

**UNIVERSITI TEKNOLOGI MALAYSIA
FINAL EXAMINATION SEMESTER II, 2015 /2016**

SUBJECT CODE : SCSI2143
SUBJECT NAME : PROBABILITY & STATISTICAL DATA ANALYSIS
SECTION : 01/02/03/04/05/06/07/08/09/10/11/12/13/14/15/16/17
TIME :
DATE/DAY :
VENUES :

INSTRUCTIONS :

(Please Write Your Lecture Name And Section In Your Answer Booklet)

Name	
I/C No.	
Year / Course	
Section	
Lecturer Name	

This questions paper consists of **ten (10)** printed pages excluding this page.

Question 1**10 marks**

- a) Sample of two different models of cars were selected, and the actual speed for each car was determined when the speedometer registered 50 mph. The resulting 95% confidence intervals for mean actual speed were (50.3, 52.7) and (49.8, 50.6) respectively. Assuming that the two sample standard deviations are equal, which confidence interval is based on the larger sample size? Explain your reasoning.

(3 marks)

- b) Three different statistics are being considered for estimating a population characteristic. The sampling distributions of the three statistics are shown in Figure 1.




Figure 1

Which statistic would you recommend? Explain your choice.

(3 marks)

- c) A manufacturer conducted a survey among 500 randomly selected target market households in the test market for its new disposable diapers. The objective of the survey was to determine the market share for its new brand. If the sample point estimate for market share is 15%, develop a 90% confidence interval.

(4 marks)

Question 2**20 marks**

- a) A marketing study found that the mean spending in 15 categories of consumer items for 150 respondents in the 18 to 34 age group was RM75.50 with a standard deviation of RM53.20. For 250 respondents in the 35+ age group, the mean and the standard deviation were RM65.20 and RM46.10, respectively. Test the hypothesis that there is no difference in the mean spending between these two populations. Use a significance level of 0.05.

(Note: Assume that the population variances are not equal). (10 marks)

- b) In a study of memory recall, ten students from a large statistics and data analysis class were selected at random and given 15 minutes to memorize a list of 20 nonsense words. Each was asked to list as many of the words as he or she could remember both 1 hour and 24 hours later. The data are as shown in Table 1. Is there evidence to suggest the mean number of words recalled after 1 hour exceeds the mean recall after 24 hours by more than 3? Use a significance level of 0.05. (10 marks)

Table 1

Student	1	2	3	4	5	6	7	8	9	10
1 hour later (words)	14	12	18	7	11	9	16	15	13	17
24 hour later (words)	10	4	14	5	8	5	11	12	9	10

Question 3**20 marks**

- a) The following table lists the frequency distribution of cars sold at an auto dealership during the past 4 quarters of 2015. Using the 10% significance level, will you reject the claim that the number of cars sold at this dealership is the same for each quarter?

Table 2

Quarter	1 st Quarter	2 nd Quarter	3 rd Quarter	4 th Quarter
Car Sold	59	40	51	90

- i) State the null and alternative hypotheses. (2 marks)
- ii) Find the critical value. (1 mark)
- iii) Calculate the test statistic. (4 marks)

iv) What does the result suggest about the claim that the number of cars sold at this dealership is the same for each quarter? (3 marks)

b) Clinical test of the allergy drugs Seldane yielded results summarized in the Table 3.

i) At the 0.05 significance level, test the claim that the occurrence of headaches is independent of the group (Seldane, placebo, control). (9 marks)

ii) Based on (i), should Seldane users be concerned about getting headaches? (1 mark)

Table 3

	Seldane Users	Placebo Users	Control
Headache	49	49	24
No Headache	732	616	602

Question 4

30 marks

a) A company is to replace its fleet of cars. Eight possible models are considered and the transport manager is asked to rank them, from 1 to 8. The current market price of the car is also recorded. Table 4 shows the ranking given by the transport manager and the current market price of the car.

Table 4

Model	Transport manager's ranking	Price (RM)
S	5	49000
T	1	70000
U	7	56000
V	2	73000
W	8	52000
X	6	57300
Y	4	68300
Z	3	65000

i) Calculate the coefficient correlation, r , using spearman rank correlation. (10 marks)

ii) Does the value of r suggest any relationship between the rank given by the transport manager and the market price value? (2 marks)

- b) A biologist assumes that there is a linear relationship between the amount of fertilizer supplied to mango plants and subsequent yield of mango obtained. Eight mango plants, of the same variety were selected at random and treated, weekly, with a solution in which x gram of fertilizer was dissolved in a fixed quantity of water. The yield y kilogram of mango was recorded as in Table 5.

Table 5

Plant	A	B	C	D	E	F	G	H	Total
x	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	22
y	3.9	4.4	5.8	6.6	7.0	7.1	7.3	7.7	49.8

- Calculate the equation of the least squares regression line of y on x (12 marks)
- Estimate the yield of the plant treated weekly with 2.8 grams of fertilizer (4 marks)
- If a plant is treated weekly with 15 gram fertilizer, it that appropriate to use the equation to predict the yield of the plant. (2 marks)

QUESTION 5

20 marks

A randomly selected research papers submitted by students during five semesters were analysed for plagiarism. Suppose that the number of papers sampled in each of the five semesters and the means and standard deviations for percentage of plagiarized words are as given in Table 6. Do these data provide evidence to support the claim that mean percentage of plagiarized words is not the same for all five semesters? Test the appropriate hypotheses using $\alpha = 0.05$. Assume that the conditions necessary for the ANOVA F test are reasonable.

Table 6

Semester	n	Mean	Standard deviation
1	32	6.31	3.75
2	32	3.31	3.06
3	32	1.79	3.25
4	32	1.83	3.13
5	32	1.50	2.37

(20 marks)

List of Formula

$$a) \hat{p} \pm (z_{\text{critical value}}) \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

$$b) z_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{\sqrt{\frac{\sigma^2}{n_1} + \frac{\sigma^2}{n_2}}}$$

$$c) T_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}; S_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

$$d) T_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}; v = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{(s_1^2/n_1)^2}{n_1 - 1} + \frac{(s_2^2/n_2)^2}{n_2 - 1}}$$

$$e) \chi_0^2 = \sum_i^n \frac{(o_{ij} - e_{ij})^2}{e_{ij}}; \text{ degree of freedom } (r-1)(c-1)$$

$$f) \chi_0^2 = \sum_i^n \frac{(o_i - e_i)^2}{e_i}; \text{ degree of freedom } = k-1$$

$$g) t = \frac{D - \mu_D}{S_D / \sqrt{n}}; D = \frac{\sum D}{n}; S_D = \sqrt{\frac{\sum D^2 - \frac{(\sum D)^2}{n}}{n-1}}$$

$$h) r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)} \quad \text{where } d_i = y_i - x_i$$

$$i) b_1 = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sum x^2 - \frac{(\sum x)^2}{n}} \quad b_0 = y - b_1 x$$

$$j) F = \frac{\text{variance between samples } \frac{s_x^2}{n}}{\text{variance within sample } s_p^2}; \text{ numerator} = k-1; \text{ denominator} = k(n-1)$$

F Values for $\alpha = 0.05$

d_2	d_1								
	1	2	3	4	5	6	7	8	9
1	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	240.5
2	18.51	19.00	19.16	19.25	19.3	19.33	19.35	19.37	19.38
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04
120	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96
inf	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88




TABLE A-2 Standard Normal (z) Distribution

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	* .4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	↑ .4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	* .4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	↑ .4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990
3.10 and higher	.4999									

NOTE: For values of z above 3.09, use 0.4999 for the area.

*Use these common values that result from interpolation:

z score	Area
1.645	0.4500
2.575	0.4950

From Frederick C. Mosteller and Robert E. K. Rourke, *Sturdy Statistics*, 1973, Addison-Wesley Publishing Co., Reading, MA. Reprinted with permission of Frederick Mosteller.




TABLE A-3 t Distribution

Degrees of Freedom	α					
	.005 (one tail)	.01 (one tail)	.025 (one tail)	.05 (one tail)	.10 (one tail)	.25 (one tail)
	.01 (two tails)	.02 (two tails)	.05 (two tails)	.10 (two tails)	.20 (two tails)	.50 (two tails)
1	63.657	31.821	12.706	6.314	3.078	1.000
2	9.925	6.965	4.303	2.920	1.886	.816
3	5.841	4.541	3.182	2.353	1.638	.765
4	4.604	3.747	2.776	2.132	1.533	.741
5	4.032	3.365	2.571	2.015	1.476	.727
6	3.707	3.143	2.447	1.943	1.440	.718
7	3.500	2.998	2.365	1.895	1.415	.711
8	3.355	2.896	2.306	1.860	1.397	.706
9	3.250	2.821	2.262	1.833	1.383	.703
10	3.169	2.764	2.228	1.812	1.372	.700
11	3.106	2.718	2.201	1.796	1.363	.697
12	3.054	2.681	2.179	1.782	1.356	.696
13	3.012	2.650	2.160	1.771	1.350	.694
14	2.977	2.625	2.145	1.761	1.345	.692
15	2.947	2.602	2.132	1.753	1.341	.691
16	2.921	2.584	2.120	1.746	1.337	.690
17	2.898	2.567	2.110	1.740	1.333	.689
18	2.878	2.552	2.101	1.734	1.330	.688
19	2.861	2.540	2.093	1.729	1.328	.688
20	2.845	2.528	2.086	1.725	1.325	.687
21	2.831	2.518	2.080	1.721	1.323	.686
22	2.819	2.508	2.074	1.717	1.321	.686
23	2.807	2.500	2.069	1.714	1.320	.685
24	2.797	2.492	2.064	1.711	1.318	.685
25	2.787	2.485	2.060	1.708	1.316	.684
26	2.779	2.479	2.056	1.706	1.315	.684
27	2.771	2.473	2.052	1.703	1.314	.684
28	2.763	2.467	2.048	1.701	1.313	.683
29	2.756	2.462	2.045	1.699	1.311	.683
Large (z)	2.575	2.326	1.960	1.645	1.282	.675

TABLE A-4		Chi-Square (χ^2) Distribution								
Degrees of Freedom	Area to the Right of the Critical Value									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	—	—	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.071	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.299
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.042	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.194	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.257	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.954	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169