


SULIT



UTM
UNIVERSITI TEKNOLOGI MALAYSIA

Faculty of
Computing

UNIVERSITI TEKNOLOGI MALAYSIA
FINAL EXAMINATION SEMESTER I, 2017 / 2018

SUBJECT CODE : **SCSI 2143**

SUBJECT NAME : **PROBABILITY & STATISTICAL DATA ANALYSIS**

SECTION : **ALL**

TIME : **9.00 AM – 12.00 NOON**

DATE/DAY : **16 / 01 / 2018 (TUESDAY)**

VENUES : **BK 2**

INSTRUCTIONS :

Answer **ALL** questions in the **ANSWER BOOKLET**.

(Please Write Your Lecture Name And Section In Your Answer Booklet)

Name	
I/C No.	
Year / Course	
Section	
Lecturer Name	

This questions paper consists of **TEN (10)** printed pages including this page.

QUESTION 1**[10 Marks]**

A study of 40 bowlers showed that their mean score was 186. The standard deviation of the population is 6.

- a) Find the 95% confidence interval of the mean score for all bowlers. (4 marks)
- b) Find the 95% confidence interval of the mean score if a sample of 100 bowlers is used instead of a sample of 40. (4 marks)
- c) Which interval is smaller? Justify your answer. (2 marks)

QUESTION 2**[10 Marks]**

In May, the fill weight of 6 kg boxes of laundry soap had a mean of 6.13kg with a standard deviation of 0.095. The goal was to decrease the standard deviation. The company decided to adjust the filling machines and then test $H_0: \sigma = 0.095$ against $H_1: \sigma < 0.095$. In June a random sample of size $n = 20$ yielded $\bar{x} = 6.10$ and $s = 0.065$ with $\alpha = 0.05$ (significance level), was the company successful?

QUESTION 3**[20 Marks]**

- a) A consumer-research organization routinely selects several car models each year and evaluates their fuel efficiency. In this year's study of two similar subcompact models from two different automakers, the average fuel mileage for 12 cars of brand A was 27.2 miles per gallon (mpg), with a standard deviation of 3.8 mpg. The nine brand B cars that were tested averaged 32.1 mpg, with a standard deviation of 4.3 mpg. At $\alpha = 0.01$, should it conclude that brand B cars have higher average fuel mileage than brand A cars? Assume the populations to be approximately normally distributed with equal variance. (10 marks)
- b) A composition teacher wishes to see whether a new grammar program will reduce the number of grammatical errors her students make when writing a two-page essay. The data are shown in Table 1. At $\alpha = 0.025$, can it be concluded that the number of errors have been reduced? (10 marks)

Table 1

Student	1	2	3	4	5	6
Errors before	12	9	0	5	4	3
Errors after	9	6	1	3	2	3

Question 4

[20 Marks]

Table 2 shows the data on the age at which smoking started for a sample of 1031 men who smoked low-tar cigarettes

Table 2

Age	Frequency
<16	237
16-17	258
18-20	320
≤ 21	216

- a) What does the result suggest about the claim that the number of men started smoking at all age categories are equal. Test the claim at $\alpha = 0.05$ (10 marks)
- b) At $\alpha = 0.05$ test another claimed that the percentage of men started smoking at age below 16 is 25%, between 16-17 is 20%, between 18-20 is 30% and after age 21 is 25%. (10 marks)

Question 5**[20 Marks]**

The time (x) in years that an employee spent at a company and the employee's hourly pay (y), for 5 employees are listed in the Table 3 below.

Table 3

x	y
5	25
3	20
4	21
10	35
15	38

- a) Calculate the correlation coefficient r (4 marks)
- b) Interpret the result found in 5(a) (2 marks)
- c) Find the regression slope, b_1 (3 marks)
- d) Find the regression intercept, b_0 (3 marks)
- e) Find the regression equation, $\hat{y}_i$ (5 marks)
- f) Use the equation $\hat{y}_i$ to predict the hourly pay rate of an employee who has worked for 20 years. (3 marks)

Question 6**[20 Marks]**

Four brands of flashlight batteries are to be compared by testing each brand in five flashlights. Twenty flashlights are randomly selected and divided into four groups of five flashlights each. Then each group of flashlights uses a different brand of battery. The lifetimes of the batteries, to the nearest hour, are as shown in Table 4.

Table 4

Brand	n	Mean	Standard deviation
A	5	33.6	6.43
B	5	30.8	3.56
C	5	29.8	4.71
D	5	28.6	6.8

Preliminary data analyses indicate that the independent samples come from normal populations with equal standard deviations. At the 5% significance level, does there appear to be a difference in mean lifetime among the four brands of batteries? (20 marks)

List of Formula

$$a) \quad \bar{x} \pm (z \text{ critical value}) \frac{\sigma}{\sqrt{n}}$$

$$b) \quad \chi^2 = \frac{(n-1)s^2}{\sigma^2}$$

$$c) \quad z_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{\sqrt{\frac{\sigma^2}{n_1} + \frac{\sigma^2}{n_2}}}$$

$$d) \quad T_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}; S_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

$$e) \quad T_o = \frac{\bar{x}_1 - \bar{x}_2 - \Delta_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}; v = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{(s_1^2/n_1)^2}{n_1 - 1} + \frac{(s_2^2/n_2)^2}{n_2 - 1}}$$

$$f) \quad \chi_0^2 = \sum_i^n \frac{(o_{ij} - e_{ij})^2}{e_{ij}}; \text{ degree of freedom } (r-1)(c-1)$$

$$g) \quad \chi_0^2 = \sum_i^n \frac{(o_i - e_i)^2}{e_i}; \text{ degree of freedom } = k-1$$

$$h) \quad t = \frac{\bar{D} - \mu_D}{s_D / \sqrt{n}}; \quad \bar{D} = \frac{\sum D}{n}; \quad s_D = \sqrt{\frac{\sum D^2 - \frac{(\sum D)^2}{n}}{n-1}}$$

$$i) \quad r = \frac{\sum xy - (\sum x \sum y)/n}{\sqrt{(\sum x^2 - (\sum x)^2/n)(\sum y^2 - (\sum y)^2/n)}}; r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)} \quad \text{where } d_i = y_i - x_i$$

$$j) \quad b_1 = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sum x^2 - \frac{(\sum x)^2}{n}} \quad b_0 = \bar{y} - b_1 \bar{x}$$

$$k) \quad F = \frac{\text{variance between sample}}{\text{variance within sample}} = \frac{ns^2_x}{s_p^2}; \text{ numerator } = k-1; \text{ denominator } = k(n-1)$$

F Values for $\alpha = 0.05$

d_2	d_1								
	1	2	3	4	5	6	7	8	9
1	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	240.5
2	18.51	19.00	19.16	19.25	19.3	19.33	19.35	19.37	19.38
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04
120	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96
inf	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88




TABLE A-2 Standard Normal (z) Distribution

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990
3.10 and higher	.4999									

NOTE: For values of z above 3.09, use 0.4999 for the area.

*Use these common values that result from interpolation:

z score	Area
1.645	0.4500
2.575	0.4950

From Frederick C. Mosteller and Robert E. K. Rourke, *Sturdy Statistics*, 1973, Addison-Wesley Publishing Co., Reading, MA. Reprinted with permission of Frederick Mosteller.




TABLE A-3 t Distribution

Degrees of Freedom	α					
	.005	.01	.025	.05	.10	.25
	(one tail) .01 (two tails)	(one tail) .02 (two tails)	(one tail) .05 (two tails)	(one tail) .10 (two tails)	(one tail) .20 (two tails)	(one tail) .50 (two tails)
1	63.657	31.821	12.706	6.314	3.078	1.000
2	9.925	6.965	4.303	2.920	1.886	.816
3	5.841	4.541	3.182	2.353	1.638	.765
4	4.604	3.747	2.776	2.132	1.533	.741
5	4.032	3.365	2.571	2.015	1.476	.727
6	3.707	3.143	2.447	1.943	1.440	.718
7	3.500	2.998	2.365	1.895	1.415	.711
8	3.355	2.896	2.306	1.860	1.397	.706
9	3.250	2.821	2.262	1.833	1.383	.703
10	3.169	2.764	2.228	1.812	1.372	.700
11	3.106	2.718	2.201	1.796	1.363	.697
12	3.054	2.681	2.179	1.782	1.356	.696
13	3.012	2.650	2.160	1.771	1.350	.694
14	2.977	2.625	2.145	1.761	1.345	.692
15	2.947	2.602	2.132	1.753	1.341	.691
16	2.921	2.584	2.120	1.746	1.337	.690
17	2.898	2.567	2.110	1.740	1.333	.689
18	2.878	2.552	2.101	1.734	1.330	.688
19	2.861	2.540	2.093	1.729	1.328	.688
20	2.845	2.528	2.086	1.725	1.325	.687
21	2.831	2.518	2.080	1.721	1.323	.686
22	2.819	2.508	2.074	1.717	1.321	.686
23	2.807	2.500	2.069	1.714	1.320	.685
24	2.797	2.492	2.064	1.711	1.318	.685
25	2.787	2.485	2.060	1.708	1.316	.684
26	2.779	2.479	2.056	1.706	1.315	.684
27	2.771	2.473	2.052	1.703	1.314	.684
28	2.763	2.467	2.048	1.701	1.313	.683
29	2.756	2.462	2.045	1.699	1.311	.683
Large (z)	2.575	2.326	1.960	1.645	1.282	.675

Formulas and Tables
for *Essentials of Statistics*, by Mario F. Triola
©2002 by Addison-Wesley.

TABLE A-4 Chi-Square (χ^2) Distribution										
Degrees of Freedom	Area to the Right of the Critical Value									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	—	—	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.071	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.299
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.042	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.194	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.257	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.954	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169

From Donald B. Owen, *Handbook of Statistical Tables*, ©1962 Addison-Wesley Publishing Co., Reading, MA. Reprinted with permission of the publisher.