


Inference of pairwise relatedness

Ida Moltke, Naples, May 2017



Outline

1. Introduction
 - Goal and motivation
 - Definition of key concepts
 - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
 - Overview and intuition
 - Model
 - ML inference
3. Problems and more advanced methods
 - Problems with the standard methods
 - NGSrelate (for NGS data)
 - Methods for ancient DNA
 - RelateAdmix (for admixed individuals)
4. Exercises

Outline

1. Introduction

- Goal and motivation
- Definition of key concepts
- Overview of current methods for inference

2. A simple maximum likelihood solution for called genotypes

- Overview and intuition
- Model
- ML inference


3. Problems and more advanced methods

- Problems with the standard methods
- NGSrelate (for NGS data)
- Methods for ancient DNA
- RelateAdmix (for admixed individuals)

4. Exercises

Goal

- ▶ We want to infer how two individuals are related based on DNA
- ▶ For example:



Motivation

- ▶ Many methods assume individuals are unrelated.
- ▶ Violations can lead to wrong conclusions!

Motivation

- ▶ Many methods assume individuals are unrelated.
- ▶ Violations can lead to wrong conclusions!
- ▶ Can be of interest in it self, e.g.:
 - ▶ forensics (paternity testing, DNA evidence)



Motivation

- ▶ Many methods assume individuals are unrelated.
- ▶ Violations can lead to wrong conclusions!
- ▶ Can be of interest in it self, e.g.:
 - ▶ forensics (paternity testing, DNA evidence)




- ▶ can reveal cultural practices in the past



Key concept: Identity-By-Descent

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color.

- Note with this definition:
 - Individuals can be identical in a locus without being IBD

Key concept: Identity-By-Descent

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color.

- Note with this definition:
 - Individuals can be identical in a locus without being IBD
 - IBD sharing cannot be observed, **so needs to be inferred**

Key concept: Identity-By-Descent

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color.

- Note with this definition:
 - Individuals can be identical in a locus without being IBD
 - IBD sharing cannot be observed, **so needs to be inferred**
 - Any non-inbred pair will in a locus **share 0, 1 or 2 alleles IBD**.

Key concept: Identity-By-Descent

- One definition: DNA sequence **identity due to recent common ancestry**



The individuals are IBD in regions where they have the same color.

- Note with this definition:
 - Individuals can be identical in a locus without being IBD
 - IBD sharing cannot be observed, **so needs to be inferred**
 - Any non-inbred pair will in a locus **share 0, 1 or 2 alleles IBD**.
 - Importantly: **the closer related, the more IBD sharing**

IBD and relatedness coefficients (R)

- $\mathbf{R} = (\mathbf{k}_0, \mathbf{k}_1, \mathbf{k}_2)$: fractions of genome with 0, 1 and 2 alleles IBD

IBD and relatedness coefficients (R)

- $R = (k_0, k_1, k_2)$: fractions of genome with 0, 1 and 2 alleles IBD
- We expect different relationships to have different values of R , e.g.:

Relationship	k_0	k_1	k_2
Monozygotic twins	0	0	1
Parent-offspring	0	1	0
Siblings	0.25	0.5	0.25
Halfsiblings/Uncle-nephew/grandparent-child	0.5	0.5	0
First cousins	0.75	0.25	0
Second cousins	0.9375	0.0625	0
Unrelated	1	0	0

- Hence we can (often) use R to infer how two individuals are related

IBD and relatedness coefficients (R)


- ▶ $R = (k_0, k_1, k_2)$: fractions of genome with 0, 1 and 2 alleles IBD
- ▶ We expect different relationships to have different values of R , e.g.:

Relationship	k_0	k_1	k_2
Monozygotic twins	0	0	1
Parent-offspring	0	1	0
Siblings	0.25	0.5	0.25
Halfsiblings/Uncle-nephew/grandparent-child	0.5	0.5	0
First cousins	0.75	0.25	0
Second cousins	0.9375	0.0625	0
Unrelated	1	0	0

- ▶ Hence we can (often) use R to infer how two individuals are related
- ▶ There are other measures like kinship
- ▶ But they are all functions of R so we will focus on that


How?

- This can e.g. be done by plotting k_1 against k_2 :



How?

- This can e.g. be done by plotting k_1 against k_2 :



- However, we need to estimate R (because it can't be observed)!

How do we infer R?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most used program is probably PLINK

Importantly, all of them are based on several assumptions

- ▶ that the individuals are from a **diploid** species

How do we infer R?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most used program is probably PLINK

Importantly, all of them are based on several assumptions

- ▶ that the individuals are from a **diploid** species
- ▶ that the individuals are **not inbred**

How do we infer R ?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most used program is probably PLINK

Importantly, all of them are based on several assumptions

- ▶ that the individuals are from a **diploid** species
- ▶ that the individuals are **not inbred**
- ▶ that we have **called genotypes** for the two individuals in L loci

How do we infer R?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most used program is probably PLINK

Importantly, all of them are based on several assumptions

- ▶ that the individuals are from a **diploid** species
- ▶ that the individuals are **not inbred**
- ▶ that we have **called genotypes** for the two individuals in L loci
- ▶ that the individuals are from the same **homogenous population**

How do we infer R?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most used program is probably PLINK

Importantly, all of them are based on several assumptions

- ▶ that the individuals are from a **diploid** species
- ▶ that the individuals are **not inbred**
- ▶ that we have **called genotypes** for the two individuals in L loci
- ▶ that the individuals are from the same **homogenous population**
- ▶ that the population **allele frequencies** are known
(or that you provide enough samples to estimate them)

Outline

1. Introduction
 - Goal and motivation
 - Definition of key concepts
 - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
 - Overview and intuition
 - Model
 - ML inference
3. Problems and more advanced methods
 - Problems with the standard methods
 - NGSrelate (for NGS data)
 - Methods for ancient DNA
 - RelateAdmix (for admixed individuals)
4. Exercises

A simple ML approach (overview)

1. Define a model that allows us to calculate $P(data|R)$ (likelihood)
2. Estimate R by finding the R value the maximises $P(data|R)$ (MLE)

Intuition behind (what information is used)

Although we cannot observe IBD, only genotypes, we note that

- ▶ **If alleles are not identical they cannot be IBD**
- ▶ If alleles are identical it could be due to either IBD or chance
- ▶ If an allele is frequent, identity will occur often simply by chance
- ▶ If an allele is rare, identity will occur rarely by chance
- ▶ **So the rarer the allele, the more identity makes us believe the individuals are IBD**

Model

- ▶ We want a model that allows us to calculate $P(data|R)$
- ▶ Notation: for a pair of non-inbred individuals genotyped in L loci we let
 - ▶ G_j be their genotypes in locus j , e.g. $G_j = (AA, aa)$
 - ▶ Z_j indicate how many alleles they share IBD in locus j (unobserved)

Model

- ▶ We want a model that allows us to calculate $P(data|R)$
- ▶ Notation: for a pair of non-inbred individuals genotyped in L loci we let
 - ▶ G_j be their genotypes in locus j , e.g. $G_j = (AA, aa)$
 - ▶ Z_j indicate how many alleles they share IBD in locus j (unobserved)
- ▶ For a single locus, j , first we can write:

$$\begin{aligned} P(G_j|R) &= P(Z_j = 0|R)P(G_j|Z_j = 0) + \\ &\quad P(Z_j = 1|R)P(G_j|Z_j = 1) + \\ &\quad P(Z_j = 2|R)P(G_j|Z_j = 2) \quad = \sum_{i=0}^2 P(Z_j = i|R)P(G_j|Z_j = i) \end{aligned}$$

Model

- ▶ We want a model that allows us to calculate $P(data|R)$
- ▶ Notation: for a pair of non-inbred individuals genotyped in L loci we let
 - ▶ G_j be their genotypes in locus j , e.g. $G_j = (AA, aa)$
 - ▶ Z_j indicate how many alleles they share IBD in locus j (unobserved)
- ▶ For a single locus, j , first we can write:

$$\begin{aligned} P(G_j|R) &= P(Z_j = 0|R)P(G_j|Z_j = 0) + \\ &\quad P(Z_j = 1|R)P(G_j|Z_j = 1) + \\ &\quad P(Z_j = 2|R)P(G_j|Z_j = 2) \quad = \sum_{i=0}^2 P(Z_j = i|R)P(G_j|Z_j = i) \end{aligned}$$

- ▶ Note that $P(Z_j = i|R)$ is simply k_i for all $i \in \{0, 1, 2\}$, so we get

$$P(G_j|R) = \sum_{i=0}^2 k_i P(G_j|Z_j = i)$$

Model ($P(G_j|Z_j = i)$)

- Assuming Hardy-Weinberg Equilibrium we can derive $P(G_j|Z_j = i)$:

G_j	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	f_A^4	f_A^3	f_A^2	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

Model ($P(G_j|Z_j = i)$)

- Assuming Hardy-Weinberg Equilibrium we can derive $P(G_j|Z_j = i)$:

G_j	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	f_A^4	f_A^3	f_A^2	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

$$1. P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$$

Model ($P(G_j|Z_j = i)$)

- Assuming Hardy-Weinberg Equilibrium we can derive $P(G_j|Z_j = i)$:

G_j	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	f_A^4	f_A^3	f_A^2	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$

Model ($P(G_j|Z_j = i)$)

- Assuming Hardy-Weinberg Equilibrium we can derive $P(G_j|Z_j = i)$:

G_j	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	f_A^4	f_A^3	f_A^2	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$
- $P(AA, Aa |Z_j = 2) = P(AA)P(Aa|AA, Z_j = 2) = P(AA) \times 0 = 0$

Model ($P(G_j|Z_j = i)$)

- Assuming Hardy-Weinberg Equilibrium we can derive $P(G_j|Z_j = i)$:

G_j	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	f_A^4	f_A^3	f_A^2	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$
- $P(AA, Aa |Z_j = 2) = P(AA)P(Aa|AA, Z_j = 2) = P(AA) \times 0 = 0$

- Captures connection between IBD, genotypes and allele frequencies:

- 3) captures that alleles have to be identical to be IBD
- 1) and 2) captures that the rarer an allele is, the more will observing identity make us believe the individuals are IBD

ML inference based on the model

- **Assuming loci are independent** we get the full likelihood:


$$P(G_1, G_2, \dots, G_L | R) = \prod_{j=1}^L P(G_j | R)$$

ML inference based on the model

- ▶ **Assuming loci are independent** we get the full likelihood:

$$P(G_1, G_2, \dots, G_L | R) = \prod_{j=1}^L P(G_j | R)$$

- ▶ This function is optimized for R and we get MLE of R
- ▶ Most often done using an EM algorithm
- ▶ NB does not always give you the MLE!




Outline

1. Introduction
 - Goal and motivation
 - Definition of key concepts
 - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
 - Overview and intuition
 - Model
 - ML inference
3. Problems and more advanced methods
 - Problems with the standard methods
 - NGSrelate (for NGS data)
 - Methods for ancient DNA
 - RelateAdmix (for admixed individuals)
4. Exercises

Problems

Most current methods


- ▶ work on called genotypes and **perform poorly on low depth NGS data**
- ▶ E.g. PLINK on simulated data from 20 1st cousins (k0 is overestimated):



Problems (cont.)

For ancient DNA there is a range of additional issues:

- ▶ very low coverage
- ▶ increased error rates (especially transitions)




- ▶ not enough samples available for proper allele frequency estimation

Problems (cont.)



Finally, most current methods have problems with admixed individuals.

E.g. PLINK:



Handling NGS data (NGSrelate)



- A solution is to use genotype likelihoods (GLs) instead of called genotypes



- Quantifies the uncertainty in what the genotype is

Handling NGS data (NGSrelate)

- ▶ A solution is to use genotype likelihoods (GLs) instead of called genotypes




- ▶ Quantifies the uncertainty in what the genotype is
- ▶ We can write a new likelihood for a locus j which takes this uncertainty into account using these:

$$P(D_j|R) = \sum_{G_j \in \{\text{all possible genotype pairs}\}} P(D_{j,i1}|G_{j,i1})P(D_{j,i2}|G_{j,i2})P(G_j|R)$$

Handling NGS data (NGSrelate)

- ▶ A solution is to use genotype likelihoods (GLs) instead of called genotypes




- ▶ Quantifies the uncertainty in what the genotype is
- ▶ We can write a new likelihood for a locus j which takes this uncertainty into account using these:

$$P(D_j|R) = \sum_{G_j \in \{\text{all possible genotype pairs}\}} P(D_{j,i1}|G_{j,i1})P(D_{j,i2}|G_{j,i2})P(G_j|R)$$

- ▶ NB when genotypes are known the GLs are 0 for all but the true genotype, so the likelihood becomes the same as before!

Handling NGS data (NGSrelate)

► NGSrelate vs PLINK:




Methods for ancient DNA

- ▶ In some cases methods like NGSrelate can be used (if frequency info is available and one removes transitions)
- ▶ In other cases you need special methods (e.g. due to lack of allele frequencies)
- ▶ Still highly active research field, this year at least 2 new methods came out (Kuhn et al. BioRXiv and Theunert et al. Genetics)

Handling admixture

- ▶ the likelihood can also be adjusted to take admixture into account
- ▶ this is done in the software relateAdmix
- ▶ comparison of relateAdmix and PLINK on admixed individuals:



Outline

1. Introduction
 - Goal and motivation
 - Definition of key concepts
 - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
 - Overview and intuition
 - Model
 - ML inference
3. Problems and more advanced methods
 - Problems with the standard methods
 - NGSrelate (for NGS data)
 - Methods for ancient DNA
 - RelateAdmix (for admixed individuals)
4. Exercises

Exercise

Go to <http://popgen.dk/ida/EMBONaples2017/web/> and solve exercise 1 & 2

(run the exercises on the server logged in with `ssh -Y`)