


# Inference of pairwise relatedness

Ida Moltke, Naples, April 2018



# Outline

## 1. Introduction

- Goal and motivation
- Definition of key concepts
- Overview of current methods for inference

## 2. A simple maximum likelihood solution for called genotypes

- Overview and intuition
- Model
- ML inference


## 3. Problem and more advanced methods

- Problems with standard methods
- NGSrelate (for NGS data)
- RelateAdmix (for admixed individuals)

## 4. Exercise

# Goal

- We want to infer how two individuals are related based on DNA
- For example:



# Motivation

- ▶ Many methods assume individuals are unrelated
- ▶ Violations can lead to wrong conclusions!

# Motivation

- ▶ Many methods assume individuals are unrelated
- ▶ Violations can lead to wrong conclusions!
- ▶ Can be of interest in itself, e.g.:
  - ▶ forensics (paternity testing, DNA evidence)




- ▶ can help reveal past or current cultural/social practices



# Key concept: Identity by Descent (IBD)

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color

- Note with this definition:
  - Individuals can be identical in a locus without being IBD

# Key concept: Identity by Descent (IBD)

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color

- Note with this definition:
  - Individuals can be identical in a locus without being IBD
  - Hence IBD sharing cannot be observed, **so needs to be inferred**

# Key concept: Identity by Descent (IBD)

- One definition: DNA sequence **identity due to recent common ancestry**




The individuals are IBD in regions where they have the same color

- Note with this definition:
  - Individuals can be identical in a locus without being IBD
  - Hence IBD sharing cannot be observed, **so needs to be inferred**
  - Any non-inbred pair will in a locus **share 0, 1 or 2 alleles IBD**



# Key concept: Identity by Descent (IBD)

- One definition: DNA sequence **identity due to recent common ancestry**



The individuals are IBD in regions where they have the same color

- Note with this definition:
  - Individuals can be identical in a locus without being IBD
  - Hence IBD sharing cannot be observed, **so needs to be inferred**
  - Any non-inbred pair will in a locus **share 0, 1 or 2 alleles IBD**
  - And **the closer related, the more IBD sharing**

# IBD and relatedness coefficients (R)

- $\mathbf{R} = (\mathbf{k}_0, \mathbf{k}_1, \mathbf{k}_2)$ : fractions of genome with 0, 1 and 2 alleles IBD

# IBD and relatedness coefficients (R)

- $\mathbf{R} = (k_0, k_1, k_2)$ : fractions of genome with 0, 1 and 2 alleles IBD
- We expect different relationships to have different values of  $R$ , e.g.:

Relationship	$k_0$	$k_1$	$k_2$
Monozygotic twins	0	0	1
Parent-offspring	0	1	0
Siblings	0.25	0.5	0.25
Halfsiblings/Uncle-nephew/grandparent-child	0.5	0.5	0
First cousins	0.75	0.25	0
Second cousins	0.9375	0.0625	0
Unrelated	1	0	0

- Hence we can (often) use  $R$  to infer how two individuals are related

# IBD and relatedness coefficients ( $R$ )


- ▶  $R = (k_0, k_1, k_2)$ : fractions of genome with 0, 1 and 2 alleles IBD
- ▶ We expect different relationships to have different values of  $R$ , e.g.:

Relationship	$k_0$	$k_1$	$k_2$
Monozygotic twins	0	0	1
Parent-offspring	0	1	0
Siblings	0.25	0.5	0.25
Halfsiblings/Uncle-nephew/grandparent-child	0.5	0.5	0
First cousins	0.75	0.25	0
Second cousins	0.9375	0.0625	0
Unrelated	1	0	0

- ▶ Hence we can (often) use  $R$  to infer how two individuals are related
- ▶ There are other measures like kinship
- ▶ But they are all functions of  $R$  so we will focus on  $R$


# How?

- This can e.g. be done by plotting  $k_1$  against  $k_2$ :



# How?

- This can e.g. be done by plotting  $k_1$  against  $k_2$ :



- However, we need to estimate  $R$  (because it can't be observed)

# How do we infer R?

There are a number of estimators:

- ▶ several method of moments estimators (e.g. PLINK)
- ▶ some maximum likelihood (ML) estimators (e.g. Thompson, 1975)
- ▶ most well known program is probably PLINK

Importantly, almost all of them are based on several assumptions, namely:

- ▶ that the individuals are from a **diploid species**
- ▶ that the individuals are **not inbred**
- ▶ that we have **called genotypes** for the two individuals in  $L$  loci
- ▶ that the individuals are from the **same homogenous population**
- ▶ that the population **allele frequencies are known**  
(or that you have provide enough samples to estimate them)

# Outline

1. Introduction
  - Goal and motivation
  - Definition of key concepts
  - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
  - Overview and intuition
  - Model
  - ML inference
3. Problem and more advanced methods
  - Problems with standard methods
  - NGSrelate (for NGS data)
  - RelateAdmix (for admixed individuals)
4. Exercise



## A simple ML approach (overview)

1. Define a model that allows us to calculate  $P(data|R)$  (likelihood)
2. Estimate  $R$  by finding the  $R$  value that maximises  $P(data|R)$  (MLE)

## Intuition behind (what information is used)

Although we cannot observe IBD, only genotypes, we note that

- ▶ **If alleles are not identical they cannot be IBD**
- ▶ If alleles are identical it could be due to either IBD or chance
- ▶ If an allele is frequent, identity will occur often simply by chance
- ▶ If an allele is rare, identity will only occur rarely by chance
- ▶ **So the rarer the allele, the more will identity makes us believe the individuals are IBD**

# Model

- ▶ We want a model that allows us to calculate  $P(\text{data}|\mathbf{R})$  (data=genotypes)
- ▶ Notation: for a pair of non-inbred individuals genotyped in  $L$  loci we let
  - ▶  $G_j$  be their genotypes in locus  $j$ , e.g.  $G_j = (AA, aa)$  (observed)
  - ▶  $Z_j$  indicate how many alleles they share IBD in locus  $j$  (unobserved)

# Model

- ▶ We want a model that allows us to calculate  $P(\text{data}|\mathbf{R})$  (data=genotypes)
- ▶ Notation: for a pair of non-inbred individuals genotyped in  $L$  loci we let
  - ▶  $G_j$  be their genotypes in locus  $j$ , e.g.  $G_j = (AA, aa)$  (observed)
  - ▶  $Z_j$  indicate how many alleles they share IBD in locus  $j$  (unobserved)
- ▶ For a single locus,  $j$ , we can write:

$$\begin{aligned} P(G_j|\mathbf{R}) &= P(Z_j = 0|\mathbf{R})P(G_j|Z_j = 0) + \\ &\quad P(Z_j = 1|\mathbf{R})P(G_j|Z_j = 1) + \\ &\quad P(Z_j = 2|\mathbf{R})P(G_j|Z_j = 2) = \sum_{z=0}^2 P(Z_j = z|\mathbf{R})P(G_j|Z_j = z) \end{aligned}$$

# Model

- ▶ We want a model that allows us to calculate  $P(\text{data}|\mathbf{R})$  (data=genotypes)
- ▶ Notation: for a pair of non-inbred individuals genotyped in  $L$  loci we let
  - ▶  $G_j$  be their genotypes in locus  $j$ , e.g.  $G_j = (AA, aa)$  (observed)
  - ▶  $Z_j$  indicate how many alleles they share IBD in locus  $j$  (unobserved)
- ▶ For a single locus,  $j$ , we can write:

$$\begin{aligned} P(G_j|\mathbf{R}) &= P(Z_j = 0|\mathbf{R})P(G_j|Z_j = 0) + \\ &\quad P(Z_j = 1|\mathbf{R})P(G_j|Z_j = 1) + \\ &\quad P(Z_j = 2|\mathbf{R})P(G_j|Z_j = 2) = \sum_{z=0}^2 P(Z_j = z|\mathbf{R})P(G_j|Z_j = z) \end{aligned}$$

- ▶ Note that  $P(Z_j = z|\mathbf{R})$  is simply  $k_z$  for all  $z \in \{0, 1, 2\}$ , so we get

$$P(G_j|\mathbf{R}) = \sum_{z=0}^2 k_z P(G_j|Z_j = z)$$

## Model ( $P(G_j|Z_j = z)$ )

- Assuming HWE we can derive:

$G_j$	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	$f_A^4$	$f_A^3$	$f_A^2$	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

# Model ( $P(G_j|Z_j = z)$ )

- Assuming HWE we can derive:

$G_j$	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	$f_A^4$	$f_A^3$	$f_A^2$	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

1.  $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$

# Model ( $P(G_j|Z_j = z)$ )

- Assuming HWE we can derive:

$G_j$	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	$f_A^4$	$f_A^3$	$f_A^2$	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$



# Model ( $P(G_j|Z_j = z)$ )

- Assuming HWE we can derive:

$G_j$	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	$f_A^4$	$f_A^3$	$f_A^2$	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$
- $P(AA, Aa |Z_j = 2) = P(AA)P(Aa|AA, Z_j = 2) = P(AA) \times 0 = 0$

## Model ( $P(G_j|Z_j = z)$ )

- Assuming HWE we can derive:

$G_j$	$Z_j=0$	$Z_j=1$	$Z_j=2$	
AA,AA	$f_A^4$	$f_A^3$	$f_A^2$	$\forall A$
AA,aa	$2f_A^2 f_a^2$	0	0	$A \neq a$
AA,Aa	$4f_A^3 f_a$	$2f_A^2 f_a$	0	$A \neq a$
Aa,Aa	$2f_A^2 f_a^2$	$f_A^2 f_a + f_A f_a^2$	$2f_A f_a$	$A \neq a$

- E.g.

- $P(AA, AA|Z_j = 0) = P(AA)P(AA|AA, Z_j = 0) = P(AA)P(AA) = f_A^2 \times f_A^2$
- $P(AA, AA|Z_j = 2) = P(AA)P(AA|AA, Z_j = 2) = P(AA) \times 1 = f_A^2 \times 1$
- $P(AA, Aa |Z_j = 2) = P(AA)P(Aa|AA, Z_j = 2) = P(AA) \times 0 = 0$

- Captures connection between IBD, genotypes and allele frequencies:
  - 3) captures that alleles have to be identical to be IBD
  - 1) and 2) captures that the rarer an allele is, the more will observing identity make us believe the individuals are IBD

# ML inference based on the model

- **Assuming loci are independent** we get the full likelihood:


$$P(G_1, G_2, \dots, G_M | R) = \prod_{j=1}^L P(G_j | R)$$

# ML inference based on the model

- **Assuming loci are independent** we get the full likelihood:

$$P(G_1, G_2, \dots, G_M | R) = \prod_{j=1}^L P(G_j | R)$$

- This function is optimized for  $R$  and we get MLE of  $R$
- Most often done using an EM algorithm
- NB does not always give you the MLE!




# Outline

1. Introduction
  - Goal and motivation
  - Definition of key concepts
  - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
  - Overview and intuition
  - Model
  - ML inference
3. Problem and more advanced methods
  - Problems with standard methods
  - NGSrelate (for NGS data)
  - RelateAdmix (for admixed individuals)
4. Exercise

# Problems

## Most current methods


- ▶ work on called genotypes and **perform poorly on low depth NGS data**
- ▶ E.g. PLINK on simulated data from 20 1st cousins ( $k_0$  is overestimated):



## Problems (cont.)

For ancient DNA there is a range of additional issues:


- ▶ very low coverage
- ▶ high error rates (especially transitions):



- ▶ not enough samples available for proper allele frequency estimation

## Problems (cont.)



Also most current methods have problems with admixed individuals.  
E.g. PLINK:





# Handling NGS data (NGSrelate)



- ▶ A solution is to use genotype likelihoods (GLs) instead of called genotypes



- ▶ Quantifies the uncertainty in what the true genotype is

# Handling NGS data (NGSrelate)

- A solution is to use genotype likelihoods (GLs) instead of called genotypes




- Quantifies the uncertainty in what the true genotype is
- We can write a new likelihood for a locus  $j$  which takes this uncertainty into account using these:

$$P(D_j|R) = \sum_{G_j \in \{\text{all possible genotype pairs}\}} P(D_{j,i1}|G_{j,i1})P(D_{j,i2}|G_{j,i2})P(G_j|R)$$

# Handling NGS data (NGSrelate)

- ▶ A solution is to use genotype likelihoods (GLs) instead of called genotypes




- ▶ Quantifies the uncertainty in what the true genotype is
- ▶ We can write a new likelihood for a locus  $j$  which takes this uncertainty into account using these:

$$P(D_j|R) = \sum_{G_j \in \{\text{all possible genotype pairs}\}} P(D_{j,i1}|G_{j,i1})P(D_{j,i2}|G_{j,i2})P(G_j|R)$$

- ▶ NB when genotypes are known the GLs are 0 for all but the true genotype, so the likelihood becomes the same as before!

# Handling NGS data (NGSrelate)

## ► NGSrelate vs. PLINK




# Methods for ancient DNA

- ▶ In some cases methods like NGSrelate can be used (if frequency info is available and one removes transitions)
- ▶ In other cases you need special methods (e.g. due to lack of allele frequencies)
- ▶ Still highly active research field, recently 3 new methods came out (Theunert et al. Genetics, Kuhn et al. BioRXiv, Waples et al. BioRXiv)

# Handling admixture

- ▶ the likelihood can also be adjusted to take admixture into account
- ▶ this is done in the software relateAdmix
- ▶ comparison of relateAdmix and PLINK on admixed individuals:



# Outline

1. Introduction
  - Goal and motivation
  - Definition of key concepts
  - Overview of current methods for inference
2. A simple maximum likelihood solution for called genotypes
  - Overview and intuition
  - Model
  - ML inference
3. Problem and more advanced methods
  - Problems with standard methods
  - NGSrelate (for NGS data)
  - RelateAdmix (for admixed individuals)
4. Exercise

# Exercise with PLINK and NGSrelate

Go to <http://popgen.dk/ida/Naples2018/web/> and solve exercises 1 & 2  
(run the exercises on the server logged in with ssh -Y)