

Linear Regression

Section 3.1

Dr. Cathy Poliak, cpoliak@uh.edu

University of Houston

Beginning Example

The goal is to predict the *stock_index_price* (the dependent variable) of a fictitious economy based on two independent/input variables:

- *Interest_Rate*
- *Unemployment_Rate*

The data is in the *stock_price.csv* data set in BlackBoard. This is from {<https://datatofish.com/multiple-linear-regression-in-r/>}

Questions We Want To Answer

1. Is there a relationship between *stock index price* and *interest rate*?
2. How strong is the relationship between *stock index price* and *interest rate*?
3. Is the relationship linear?
4. How accurately can we predict the *stock index price*?
5. Do both *interest rate* and *unemployment rate* contribute to the *stock index price*?
6. What is the statistical learning problem?

General Approach

- *Stock index price* is the **response** or **output**. We refer to the response usually as Y .
- *Interest rate* is an **input** or **predictor**, we will name it X_1 .
- Also, *Unemployment rate* is an **input**, we will name it X_2 .
- Let $X = (X_1, X_2, \dots, X_p)$ be p different predictors (independent) variables.
- For this example we will have an input vector as

$$\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}$$

- We assume there is some sort of relationship between X and Y , which can be written in the general form thus our model is

$$Y = f(X) + \epsilon$$

- Where ϵ captures the measurement errors and other discrepancies.
- Statistical learning refers to a set of approaches for estimating f .

Estimators

A statistic $\hat{\theta}$ used to estimate an unknown population parameter θ is called an **estimator**.

- Properties of an estimator $\hat{\theta}$
 - ▶ Accuracy - measured by **bias**

$$\text{Bias}(\hat{\theta}) = E(\hat{\theta}) - \theta$$

- ▶ Precision - measured by its variance, $\text{Var}(\hat{\theta})$. The estimated standard deviation of an estimator θ is referred to as its **standard error (SE)**.
 - ▶ The **mean squared error (MSE)** combines both measures.

$$\text{MSE}(\hat{\theta}) = E(\hat{\theta} - \theta)^2 = \text{Var}(\hat{\theta}) + [\text{Bias}(\hat{\theta})]^2$$

- In MATH 3339 we studied estimators for μ and p . In this class we will we will want estimators for $f(X)$.

Example, Estimate of μ

Suppose we take a random sample of 4 from a Normal distribution with $\mu = 10$ and $\sigma = 2$.

- Let $\bar{x} = \frac{1}{4} \sum_{i=1}^4 x_i$ be an estimator of μ . What is the expected value, bias, variance, and MSE of \bar{x} .

$$E(\bar{x}) = \mu = 10, \text{Bias}(\bar{x}) = E(\bar{x}) - \mu = 10 - 10 = 0$$

$$\text{Var}(\bar{x}) = \frac{\sigma^2}{n} = \frac{4}{4} = 1$$

$$\text{SD}(\bar{x}) = \frac{\sigma}{\sqrt{n}}$$

$$\text{MSE}(\bar{x}) = \text{Var}(\bar{x}) + \text{Bias}(\bar{x})^2 = 1$$

$$\text{MSE}(\bar{x}) = \sigma^2/n$$

- Let 8 be an estimator of μ . What is the expected value, bias, variance, and MSE of 8?

$$E(8) = 8 \quad \text{bias}(8) = 8 - 10 = -2$$

$$\text{Var}(8) = 0$$

$$\text{MSE}(8) = 0 + (-2)^2 = 4$$

$$\text{MSE}(8) = (8 - \mu)^2$$

$$\hat{p}_1 = \frac{x}{n} = \frac{\text{\# of successes in "n" trials}}{\text{\# of trials}}$$


$$X \sim \text{Bin}(n, p) \quad E(X) = np \quad \text{var}(X) = np(1-p)$$

$$E(\hat{p}_1) = E\left(\frac{X}{n}\right) = \frac{1}{n} E(X)$$

$$\text{var}(\hat{p}_1) = \text{var}\left(\frac{X}{n}\right) = \frac{1}{n^2} \text{var}(X)$$

$$p = \text{seg}(0, 1, \text{by} = 0.01)$$

$$\text{Plot}(p, \text{mse1})$$



Bill Howe, UW
src: domingo 2012

Simple Linear Regression Model

- The data are n observations on an explanatory variable x and a response variable y ,

$$(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$$

- The statistical model for simple linear regression states that the observed response y_i when the explanatory variable takes the value x_i is

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i.$$

- $\mu_y = \beta_0 + \beta_1 x_i$ is the mean response for y when $x = x_i$ a specific value of x .
- ϵ_i are the error terms for predicting y_i for each value of x_i .
- Notice in our general form that $f(X) = \beta_0 + \beta_1 X$.

Parameters of the Simple Regression Model

- The intercept: β_0 .
- The slope: β_1 .
- The goal is to obtain coefficient estimates $\hat{\beta}_0$ and $\hat{\beta}_1$ such that for each observed y_i , $y_i \approx \hat{\beta}_0 + \hat{\beta}_1 x_i$, for $i = 1, 2, \dots, n$.
- The most common approach is by the minimizing the least squares criterion.

Principle of Least Squares

- Let $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$ be the prediction for Y based on the i th value of X .
- Then $e_i = y_i - \hat{y}_i$ be the i th residual, the difference between the i th observed response value and the i th predicted value by our linear equation.
- The **residual sum of squares** (RSS) is defined by

$$\text{RSS} = e_1^2 + e_2^2 + \cdots + e_n^2$$

- The point estimates of β_0 and β_1 , denoted by $\hat{\beta}_0$ and $\hat{\beta}_1$ and called the **least squares estimates**, are those values that minimize the RSS.

$$\text{minimize } \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n [y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i)]^2$$

The Least - Squares Estimates

- The method of **least squares** selects estimators $\hat{\beta}_0$ and $\hat{\beta}_1$ that minimizes the **residual sum of squares** (RSS).
- Where the estimate of the slope coefficient β_1 is:

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

$$\hat{\beta}_1 = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2} = \frac{(n-1) \text{cov}(x, y)}{(n-1) \text{var}(x)}$$

$$= \frac{\text{cov}(x, y) s_x s_y}{s_x^2} = \text{cor}(x, y) \frac{s_y}{s_x}$$

- The estimate for the intercept β_0 is:

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- Where $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$ and $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$.

minimize

$$Q = \sum_{i=1}^n \{y_i - (\tilde{\beta}_0 + \tilde{\beta}_1 x_i)\}^2$$

$$\frac{\partial Q}{\partial \tilde{\beta}_0} = -2 \sum_{i=1}^n \{y_i - (\tilde{\beta}_0 + \tilde{\beta}_1 x_i)\}$$

$$0 = \sum_{i=1}^n \{y_i - \tilde{\beta}_0 - \tilde{\beta}_1 x_i\}$$

$$0 = \sum_{i=1}^n y_i - n \tilde{\beta}_0 - \tilde{\beta}_1 \sum_{i=1}^n x_i$$

$$\tilde{\beta}_0 = \frac{1}{n} \sum_{i=1}^n y_i - \tilde{\beta}_1 \left[\frac{1}{n} \sum_{i=1}^n x_i \right]$$

$$\boxed{\tilde{\beta}_0 = \bar{y} - \tilde{\beta}_1 \bar{x}}$$

$$\frac{\partial Q}{\partial \hat{\beta}_1} = -2 \sum_{i=1}^n x_i (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)$$

$$0 = \sum_{i=1}^n x_i y_i - \hat{\beta}_0 \sum_{i=1}^n x_i - \hat{\beta}_1 \sum_{i=1}^n x_i^2$$

$$0 = \sum_{i=1}^n x_i y_i - \left[\frac{\sum_{i=1}^n y_i - \hat{\beta}_1 \sum_{i=1}^n x_i}{n} \right] \sum_{i=1}^n x_i - \hat{\beta}_1 \sum_{i=1}^n x_i^2$$


$$\hat{\beta}_1 \left[n \sum_{i=1}^n x_i^2 - \left(\sum_{i=1}^n x_i \right)^2 \right] = n \sum_{i=1}^n x_i y_i - \sum_{i=1}^n y_i \sum_{i=1}^n x_i$$

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n x_i y_i - n \bar{y} \bar{x}}{\sum_{i=1}^n x_i^2 - n \bar{x}^2} = \frac{\text{cov}(x, y)}{\text{var}(x)}$$

Stock Prices Example

- Use the *stock_price.csv* data.
- We want to predict *stock index price* based on *interest rate*.
 1. Determine if it is a linear relationship. How can we tell?
 2. Get an estimate of the model.
 3. Is this a good fit for the data?

Do We Have A Linear Relationship?



The Estimate of the Model

```
stock.lm <- lm(Stock_Index_Price ~ Interest_Rate, data = stock_price)
summary(stock.lm)
```

Call:

```
lm(formula = Stock_Index_Price ~ Interest_Rate, data = stock_price)
```

Residuals:

| Min | 1Q | Median | 3Q | Max |
|----------|---------|--------|--------|---------|
| -183.892 | -30.181 | 4.455 | 56.608 | 101.057 |

Coefficients:

| | Estimate | Std. Error | t value | Pr(> t) |
|---------------|--------------------------|------------|---------|--------------|
| (Intercept) | -99.46 = $\hat{\beta}_0$ | 95.21 | -1.045 | 0.308 |
| Interest_Rate | 564.20 = $\hat{\beta}_1$ | 45.32 | 12.450 | 1.95e-11 *** |
| --- | | | | |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 75.96 on 22 degrees of freedom

Multiple R-squared: 0.8757, Adjusted R-squared: 0.8701

F-statistic: 155 on 1 and 22 DF, p-value: 1.954e-11

$$\hat{y}_i = -99.46 + 564.2 x_i$$

$$RSE = S = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-2}} = \sqrt{\frac{\sum_{i=1}^n \text{residual}_i^2}{n-2}}$$

$$= \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2}{n-2}}$$

Confidence Intervals for β_1

$$n=24$$

If we want to know a range of possible values for the slope we can use a confidence interval. The confidence interval for β_1 is

$$\hat{\beta}_1 \pm t_{\alpha/2, n-2} \times SE(\hat{\beta}_1)$$

where

$$SE(\hat{\beta}_1) = \sqrt{\frac{s^2}{\sum_{i=1}^n (x_i - \bar{x})^2}}$$

and $s^2 = \hat{Var}(\epsilon)$.

Given the following excerpt from the R output, determine a 95% confidence interval for the slope.

| | Estimate | Std. Error | t value | Pr(> t) |
|---------------|-----------|------------|----------|--------------|
| (Intercept) | -99.46432 | 95.21031 | -1.04468 | 3.075145e-01 |
| Interest_Rate | 564.20389 | 45.31738 | 12.45006 | 1.953975e-11 |

$$qt(1.95/2, 22) = 2.074$$

$$CI: \hat{\beta}_1 \pm t_{\alpha/2, n-2} SE(\hat{\beta}_1) = 564.204 \pm 2.074 (45.3174) = [470.21, 658.19]$$

R Function for Confidence Intervals

```
confint(stock.lm,"Interest_Rate")
```

```
                2.5 %    97.5 %  
Interest_Rate 470.2214 658.1864
```

t Test for Significance of β_1

$$y = \beta_0 + \beta_1 x + \varepsilon$$

- Hypothesis

$$H_0 : \beta_1 = 0 \text{ versus } H_a : \beta_1 \neq 0$$

Or we can think about it in this way

H_0 : There is no relationship between X and Y

versus

H_0 : There is a relationship between X and Y

- Test statistic

$$t = \frac{\hat{\beta}_1 - 0}{SE(\hat{\beta}_1)}$$

$$\text{standard error} = SE(\hat{\beta}_1) = \frac{s}{\sqrt{\sum (x_i - \bar{x})^2}}$$

With degrees of freedom $df = n - 2$.

- P -value: based on a t distribution with $n - 2$ degrees of freedom.
- Decision: Reject H_0 if $p\text{-value} \leq \alpha$.
- Conclusion: If H_0 is rejected we conclude that the explanatory variable x can be used to predict the response variable y .

$$s^2 = \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-2}$$

$$s = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-2}}$$

$$s_x^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1}$$

Example of Hypothesis Test

Given the following excerpt from the R output, Test $H_0 : \beta_1 = 0$ against $H_a : \beta_1 \neq 0$.

| | Estimate | Std. Error | t value | Pr(> t) |
|---------------|-----------|------------|----------|--------------|
| (Intercept) | -99.46432 | 95.21031 | -1.04468 | 3.075145e-01 |
| Interest_Rate | 564.20389 | 45.31738 | 12.45006 | 1.953975e-11 |

$$\text{Test statistic: } T = \frac{\hat{\beta}_1 - 0}{SE(\hat{\beta}_1)} = \frac{564.20389 - 0}{45.31738} = 12.45$$

$$\text{p-value} = P(T_{22} \leq -12.45 \text{ or } T \geq 12.45)$$

$$= 2 * pt(-12.45, 22)$$

$$= 1.95397e^{-11} \approx 0$$

Decision: Reject H_0 , since p-value ≤ 0.05

Conclusion: There is strong evidence between the relationship of interest rate and stock price

Is this good at predicting the response?

- Once we have said that this model can help predict the **output** we want to quantify at how well the model fits the data.
- Two quantities that we use is the **residual standard error** (RSE) and the **coefficient of determination** (R^2).
- These quantities are in the summary output of the `lm()` function.

Residual Standard Error

- The RSE is an estimate of the standard deviation of the ϵ .
- We can think about it as the average amount that the response will deviate from the true regression line.

$$\text{RSE} = \sqrt{\frac{1}{n-2} \sum_{i=1}^n e_i^2} = \sqrt{\frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

- The lower the RSE the better our model fits the data.

R^2 Statistic

R^2 is the percent (fraction) of variability in the response variable (Y) that is explained by the least-squares regression with the explanatory variable.

- This is a measure of how successful the regression equation was in predicting the response variable.
- The closer R^2 is to one (100%) the better our equation is at predicting the response variable.
- In the R output it is the **Multiple R-squared** value.

Calculating R^2

1. The **residual sum of squares**, denoted by RSS is

$$RSS = \sum (y_i - \hat{y}_i)^2$$

2. The **regression sum of squares**, denoted SSR is the amount of total variation that *is* explained by the model

$$SSR = \sum (\hat{y}_i - \bar{y})^2$$

3. A quantitative measure of the total amount of variation in observed values is given by the **total sum of squares**, denoted by TSS .

$$TSS = \sum (y_i - \bar{y})^2$$

Note: $TSS = SSR + RSS$

4. The **coefficient of determination**, R^2 is given by

$$R^2 = \frac{SSR}{TSS} = \frac{TSS - RSS}{TSS} = 1 - \frac{RSS}{TSS}$$

Information from the Summary in R

Call:

```
lm(formula = Stock_Index_Price ~ Interest_Rate, data = stock_price)
```

Residuals:

| Min | 1Q | Median | 3Q | Max |
|----------|---------|--------|--------|---------|
| -183.892 | -30.181 | 4.455 | 56.608 | 101.057 |

Coefficients:

| | Estimate | Std. Error | t value | Pr(> t) |
|---------------|----------|------------|---------|--------------|
| (Intercept) | -99.46 | 95.21 | -1.045 | 0.308 |
| Interest_Rate | 564.20 | 45.32 | 12.450 | 1.95e-11 *** |

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 75.96 on 22 degrees of freedom

Multiple R-squared: 0.8757 Adjusted R-squared: 0.8701

F-statistic: 155 on 1 and 22 DF, p-value: 1.954e-11

About 87.51% of the variation in stock index price can be accounted for by this model.

RSE and R^2

- The RSE is considered a measure of the *lack of fit* of the model to the data. Recall this is the estimate of the standard deviation of the residuals $y_i - \hat{y}_i$.
 - ▶ If \hat{y}_i is very far from y_i , then the RSE may be quite large.
 - ▶ This measurement depends on the units of the original values.
- The R^2 takes the form of a proportion of variance in y that is explained.
 - ▶ R^2 thus always takes on a value between 0 and 1.
 - ▶ If R^2 is close to 1 indicates that a large proportion of the variability in the response has been explained by the regression.
 - ▶ *Note:* For a simple linear regression $R^2 = \text{Cor}(X, Y)^2$.

Assumptions about the Model


The linear regression model has assumptions that we need to prove is true. We use the acronym **LINE** to remember these assumptions.

- **L**inear relationship: can we determine a linear relationship between the response and other variables?
- **I**ndependent observations: are the observations a result of a simple random sample?
- **N**ormal distribution: for any fixed value of X , Y is normally distributed.
- **E**qual variance: the variance of the residual is the same for any value of X .
- Be careful of extreme values.

Plots to Check Assumptions


Linear

Residuals vs Fitted




Normal

Q-Q Residuals



Equal Variance

Scale-Location



Outliers

Residuals vs Leverage

