

## 18.369 Problem Set 4

Due Wed, 20 March 2024.

### Problem 1: Projection operators

(You can use the Great Orthogonality Theorem [GOT] and/or the various orthogonality results for the rows/columns of the character table.)

- The representation-theory handout gives a formula for the projection operator from a function onto its component that transforms as a particular representation. Using the GOT, prove the correctness of this formula: in particular, for any function  $\psi$ , show that  $\hat{P}_i^{(\alpha)}\psi$  transforms as the  $i$ -th partner function representation  $\alpha$ . (You can use the fact, from class, that we showed that any function can be decomposed into a sum of partner functions of irreducible representations. Consider what  $\hat{P}_i^{(\alpha)}$  does to one of these partner functions.)
- Prove that the sum of the projection operators  $\hat{P}^{(\alpha)}$  over all the irreducible representations  $\alpha$  gives  $\sum_{\alpha} \hat{P}^{(\alpha)} = \sum_{\alpha,i} \hat{P}_i^{(\alpha)} = 1$  (the identity operator), by using the column-orthogonality property of the character table.
- Prove that the projection operators  $\hat{P}_i^{(\alpha)}$  are Hermitian for any unitary representation  $D^{(\alpha)}$ , assuming that  $\hat{g}$  is unitary for all  $g \in G$  (unitarity of  $\hat{g}$  is trivial to prove for symmetry groups where  $g$  is a rotation and/or translation).
- Using the previous parts, prove that  $\langle \phi_i^{(\alpha)}, \psi_j^{(\beta)} \rangle = 0$  if  $\phi_i^{(\alpha)}$  and  $\psi_j^{(\beta)}$  are partner functions of different irreducible representations  $\alpha \neq \beta$ , or if they correspond to different components  $i \neq j$  of the same (unitary) representation  $\alpha = \beta$ . (Hint: insert 1 into the inner product.)

### Problem 2: A square metal box

In class, we considered a two-dimensional ( $xy$ ) problem of light in an  $L \times L$  square of air ( $\varepsilon = 1$ ) surrounded by perfectly conducting walls (in which

$\mathbf{E} = 0$ ). We solved the case of  $\mathbf{H} = H_z(x, y)\hat{\mathbf{z}}$  and saw solutions corresponding to five different representations of the symmetry group ( $C_{4v}$ ).

- Solve for the eigenfunctions of the other polarization:  $\mathbf{E} = E_z(x, y)\hat{\mathbf{z}}$  (you will need the  $\mathbf{E}$  eigenproblem from problem set 1), with the boundary condition that  $E_z = 0$  at the metal walls. (These are separable, i.e.  $X(x)Y(y)$  for some functions  $X$  and  $Y$ .)
- Sketch (or plot on a computer) and classify these solutions according to the representations  $\Gamma_{1\dots 5}$  of  $C_{4v}$  enumerated in class. (Like in class, you will get some reducible accidental degeneracies.) Look at enough solutions to find *all five* irreps, and to illustrate the general pattern (you should find that the irreps appear in repeating patterns).

### Problem 3: Group Velocity and Material Dispersion

In class, we showed (following the book) that the group velocity  $d\omega/dk = \langle H_k, \frac{\partial \hat{H}_k}{\partial k} H_k \rangle / \langle H_k, H_k \rangle$  was equal to Poynting flux divided by energy density (both averaged over the unit cell).

Now, go through the same derivation, but in this case assume that we have a lossless *dispersive* material with a real  $\varepsilon(\mathbf{x}, \omega)$ , still assuming  $\mu = 1$  for simplicity. In this case, when you take the  $k$  derivative (assuming  $\mathbf{k} = k\hat{x}$ , i.e.  $k = k_x$ ), apply the chain rule to obtain a term with  $\frac{\partial \varepsilon}{\partial \omega} \frac{d\omega}{dk}$  on the right-hand side. Solve for  $d\omega/dk$  and show that it is Poynting flux divided by energy density, but the energy density is now the “Brillouin” energy density of a lossless dispersive medium, which we gave in the notes for Lecture 6:

$$\frac{1}{4} \left[ \frac{\partial(\omega\varepsilon)}{\partial \omega} |\mathbf{E}|^2 + |\mathbf{H}|^2 \right]$$

(for  $\mu = 1$ , where we have an additional  $1/2$  factor from the time-average).