

Quantum Amplitude Estimation – Applications to Derivative Pricing

Victorio Úbeda Sosa, Wilhelm Ågren
Svenska Handelsbanken

Quantum Autumn School
December 2-4, 2024

Classical Monte Carlo

Classical Monte Carlo

We have seen how classical Monte Carlo can be used to compute the expected payoff $\mu = \mathbb{E}[X]$ of a simple option $X = f(S_T)$:

Classical Monte Carlo

We have seen how classical Monte Carlo can be used to compute the expected payoff $\mu = \mathbb{E}[X]$ of a simple option $X = f(S_T)$:

- ▶ Sample $x_1, \dots, x_N \stackrel{i.i.d}{\sim} S_T$,

Classical Monte Carlo

We have seen how classical Monte Carlo can be used to compute the expected payoff $\mu = \mathbb{E}[X]$ of a simple option $X = f(S_T)$:

- ▶ Sample $x_1, \dots, x_N \stackrel{i.i.d}{\sim} S_T$,
- ▶ Set $\tilde{\mu}_N = \frac{1}{N} \sum_{i=1}^N f(x_i)$.

Classical Monte Carlo

We have seen how classical Monte Carlo can be used to compute the expected payoff $\mu = \mathbb{E}[X]$ of a simple option $X = f(S_T)$:

- ▶ Sample $x_1, \dots, x_N \stackrel{i.i.d}{\sim} S_T$,
- ▶ Set $\tilde{\mu}_N = \frac{1}{N} \sum_{i=1}^N f(x_i)$.

By the law of large numbers,

$$\tilde{\mu}_N \xrightarrow{\text{a.s.}} \mu, \quad \text{as } N \rightarrow \infty.$$

Classical Monte Carlo

We have seen how classical Monte Carlo can be used to compute the expected payoff $\mu = \mathbb{E}[X]$ of a simple option $X = f(S_T)$:

- ▶ Sample $x_1, \dots, x_N \stackrel{i.i.d}{\sim} S_T$,
- ▶ Set $\tilde{\mu}_N = \frac{1}{N} \sum_{i=1}^N f(x_i)$.

By the law of large numbers,

$$\tilde{\mu}_N \xrightarrow{\text{a.s.}} \mu, \quad \text{as } N \rightarrow \infty.$$

Moreover, by the central limit theorem,

$$\|\tilde{\mu}_N - \mu\|_2 = \sqrt{\mathbb{V}(\tilde{\mu}_N)} = \frac{\mathbb{V}(f(S_T))}{\sqrt{N}} \sim \mathcal{O}\left(\frac{1}{\sqrt{N}}\right).$$

Can a quantum computer do better?

Encoding the problem

Björn showed us how to construct a unitary operator \mathcal{U} encoding the desired quantity μ as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle .$$

Encoding the problem

Björn showed us how to construct a unitary operator \mathcal{U} encoding the desired quantity μ as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle .$$

I.e., $a \in [0, 1]$ is the probability of measuring $|1\rangle$ in the $(n+1)$ -th qubit.

Encoding the problem

Björn showed us how to construct a unitary operator \mathcal{U} encoding the desired quantity μ as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle.$$

I.e., $a \in [0, 1]$ is the probability of measuring $|1\rangle$ in the $(n+1)$ -th qubit.

Can we estimate a ?

Amplitude estimation

- ▶ Quantum Amplitude Estimation (QAE) does just that.
- ▶ But first:

Amplitude estimation

- ▶ Quantum Amplitude Estimation (QAE) does just that.
- ▶ But first: Phase estimation.

Phase estimation

Given:

- ▶ Unitary operator \mathcal{U} ,

Phase estimation

Given:

- ▶ Unitary operator \mathcal{U} ,
- ▶ Known eigenvector $|\psi\rangle$,

Phase estimation

Given:

- ▶ Unitary operator \mathcal{U} ,
- ▶ Known eigenvector $|\psi\rangle$, ...
- ▶ ... with associated **unknown** eigenvalue $\lambda = e^{i2\pi q/2^n}$,
 $q \in [0, 2^n)$.

Phase estimation

Given:

- ▶ Unitary operator \mathcal{U} ,
- ▶ Known eigenvector $|\psi\rangle$, ...
- ▶ ... with associated **unknown** eigenvalue $\lambda = e^{i2\pi q/2^n}$,
 $q \in [0, 2^n)$.

Task: Estimate λ .

Phase estimation

Given:

- ▶ Unitary operator \mathcal{U} ,
- ▶ Known eigenvector $|\psi\rangle$, ...
- ▶ ... with associated **unknown** eigenvalue $\lambda = e^{i2\pi q/2^n}$, $q \in [0, 2^n)$.

Task: Estimate λ .




Figure: Phase estimation circuit.

Theorem (Quantum phase estimation)

With probability $1 - \delta$, the estimate \tilde{q} provided by QPE satisfies

$$|\tilde{q} - q| \leq \frac{2}{N} \left(2 + \frac{1}{2\delta} \right) \sim \mathcal{O} \left(\frac{1}{N} \right),$$

where $N = 2^n$.

Phase estimation

“Proof idea” (integer case).

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$.

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$|0\rangle^{\otimes n} |\psi\rangle$$

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$|0\rangle^{\otimes n} |\psi\rangle \xrightarrow{H^{\otimes n}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle |\psi\rangle$$

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$\begin{aligned} |0\rangle^{\otimes n} |\psi\rangle &\xrightarrow{H^{\otimes n}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle |\psi\rangle \xrightarrow{\text{Controlled } \mathcal{U}\text{'s}} \\ &\frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle e^{2\pi i j q / 2^n} |\psi\rangle \end{aligned}$$

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$\begin{aligned} |0\rangle^{\otimes n} |\psi\rangle &\xrightarrow{H^{\otimes n}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle |\psi\rangle \mapsto \\ \text{Controlled } \mathcal{U}\text{'s} &\xrightarrow{\quad} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle e^{2\pi i j q / 2^n} |\psi\rangle = \\ \text{Phase \underline{kickback}} &\left(\frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle \right) |\psi\rangle \end{aligned}$$

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$\begin{aligned} |0\rangle^{\otimes n} |\psi\rangle &\xrightarrow{H^{\otimes n}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle |\psi\rangle \mapsto \\ \text{Controlled } \mathcal{U}\text{'s} &\xrightarrow{\quad} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle e^{2\pi i j q / 2^n} |\psi\rangle = \\ \text{Phase kickback} &\left(\frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle \right) |\psi\rangle = \\ &= \text{QFT } |q\rangle |\psi\rangle \end{aligned}$$

Phase estimation

“Proof idea” (integer case).

Assume $q \in \{0, 1, \dots, 2^n - 1\}$. Then,

$$\begin{aligned} |0\rangle^{\otimes n} |\psi\rangle &\xrightarrow{H^{\otimes n}} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle |\psi\rangle \mapsto \\ \text{Controlled } \mathcal{U}\text{'s} &\xrightarrow{\quad} \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle e^{2\pi i j q / 2^n} |\psi\rangle = \\ \text{Phase kickback} &\left(\frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle \right) |\psi\rangle = \\ &= \text{QFT } |q\rangle |\psi\rangle \xrightarrow{\text{QFT}^\dagger} |q\rangle |\psi\rangle. \end{aligned}$$



Phase estimation

Quantum Fourier Transform acts as

$$\text{QFT } |q\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle .$$

Phase estimation

Quantum Fourier Transform acts as

$$\text{QFT } |q\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle .$$

Therefore,

$$|0\rangle^{\otimes n} |\psi\rangle$$

Phase estimation

Quantum Fourier Transform acts as

$$\text{QFT} |q\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle.$$

Therefore,

$$|0\rangle^{\otimes n} |\psi\rangle \xrightarrow{\text{Phase estimation}} |q\rangle |\psi\rangle$$



The eigenvalue can be recovered as $\lambda = e^{2\pi i j q / 2^n}$.

Phase estimation

Quantum Fourier Transform acts as

$$\text{QFT} |q\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle.$$

Therefore,

$$|0\rangle^{\otimes n} |\psi\rangle \xrightarrow{\text{Phase estimation}} |q\rangle |\psi\rangle$$



The eigenvalue can be recovered as $\lambda = e^{2\pi i j q / 2^n}$.

Note. For $q \in \{0, 1, \dots, 2^n - 1\}$, phase estimation introduces no error.

Phase estimation

Quantum Fourier Transform acts as

$$\text{QFT} |q\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} e^{2\pi i j q / 2^n} |j\rangle.$$

Therefore,

$$|0\rangle^{\otimes n} |\psi\rangle \xrightarrow{\text{Phase estimation}} |q\rangle |\psi\rangle$$



The eigenvalue can be recovered as $\lambda = e^{2\pi i j q / 2^n}$.

Note. For $q \in \{0, 1, \dots, 2^n - 1\}$, phase estimation introduces no error.

Note'. For the general case $q \in [0, 2^n)$, see [[Kit95](#)], [[MW23](#)].

Amplitude estimation

Amplitude estimation

Given: unitary operator \mathcal{U} acting on $n + 1$ qubits as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle.$$

Amplitude estimation

Given: unitary operator \mathcal{U} acting on $n + 1$ qubits as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle.$$

Task: estimate a , i.e., the probability of measuring 1 in the $(n + 1)$ -th qubit.

Amplitude estimation

Given: unitary operator \mathcal{U} acting on $n + 1$ qubits as

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle.$$

Task: estimate a , i.e., the probability of measuring 1 in the $(n + 1)$ -th qubit.

Amplitude estimation

Amplitude estimation uses a combination of phase estimation and amplitude amplification [Bra+02] in order to get an estimate of a .

Amplitude estimation

Amplitude estimation uses a combination of phase estimation and amplitude amplification [Bra+02] in order to get an estimate of a .




Figure: Amplitude estimation circuit.

Where \mathcal{Q} is the Grover operator

$$\mathcal{Q} = \mathcal{U} \mathcal{S}_0 \mathcal{U}^\dagger \mathcal{S}_{\psi_0},$$

and

$$\mathcal{S}_{\psi_0} = \mathbb{I} - 2|\psi_0\rangle_n \langle \psi_0|_n \otimes |0\rangle \langle 0|, \quad \mathcal{S}_0 = \mathbb{I} - 2|0\rangle_{n+1} \langle 0|_{n+1}.$$

Amplitude estimation




Figure: Amplitude estimation circuit.

The measured number $\tilde{q} \in \{0, 1, \dots, 2^m - 1\}$ is then mapped to the estimate

$$\tilde{a} = \sin^2 \left(\tilde{\theta}_a \right),$$

where

$$\tilde{\theta}_a = \frac{\pi \tilde{q}}{M}, \quad M = 2^m.$$

Amplitude estimation

Theorem (Quantum amplitude estimation)

With probability $8/\pi^2 \approx 0.81$, the estimate \tilde{a} provided by QAE satisfies

$$|\tilde{a} - a| \leq \frac{2\pi\sqrt{a(1-a)}}{M} + \frac{\pi^2}{M^2} \sim \mathcal{O}\left(\frac{1}{M}\right).$$

Amplitude estimation

Theorem (Quantum amplitude estimation)

With probability $8/\pi^2 \approx 0.81$, the estimate \tilde{a} provided by QAE satisfies

$$|\tilde{a} - a| \leq \frac{2\pi\sqrt{a(1-a)}}{M} + \frac{\pi^2}{M^2} \sim \mathcal{O}\left(\frac{1}{M}\right).$$

Quadratic speedup over classical Monte Carlo!

- ▶ Monte Carlo: $\mathcal{O}(1/\sqrt{N})$,
- ▶ Amplitude estimation: $\mathcal{O}(1/M)$.

Amplitude estimation

Theorem (Quantum amplitude estimation)

With probability $8/\pi^2 \approx 0.81$, the estimate \tilde{a} provided by QAE satisfies

$$|\tilde{a} - a| \leq \frac{2\pi\sqrt{a(1-a)}}{M} + \frac{\pi^2}{M^2} \sim \mathcal{O}\left(\frac{1}{M}\right).$$

Quadratic speedup over classical Monte Carlo!

- ▶ Monte Carlo: $\mathcal{O}(1/\sqrt{N})$,
- ▶ Amplitude estimation: $\mathcal{O}(1/M)$.

In practice: run QAE k times and take the median value. This boost success probability to $1 - (1 - 8/\pi^2)^k$

Amplitude estimation

“Proof idea”

Amplitude estimation

“Proof idea”

Use the QPE theorem and the fact that the Grover operator acts as

$$\mathcal{Q}^k \mathcal{U} |0\rangle^{\otimes(n+1)} = \cos((2k+1)\theta_a) |\psi_0\rangle_n |0\rangle + \sin((2k+1)\theta_a) |\psi_1\rangle_n |1\rangle,$$

which implies

$$\mathbb{P}(|1\rangle) = \sin^2((2k+1)\theta_a),$$

where $\theta_a = \arcsin(\sqrt{a})$.

A toy example

A toy example

Let \mathcal{U} describe a Bernoulli random variable with success probability p :

$$\mathcal{U}|0\rangle = \sqrt{1-p}|0\rangle + \sqrt{p}|1\rangle.$$

A toy example

Let \mathcal{U} describe a Bernoulli random variable with success probability p :

$$\mathcal{U}|0\rangle = \sqrt{1-p}|0\rangle + \sqrt{p}|1\rangle.$$

We can model \mathcal{U} with a rotation around the Y -axis:

$$\mathcal{U} = R_Y(\theta_p), \quad \theta_p = 2 \arcsin(\sqrt{p}).$$

A toy example

Let \mathcal{U} describe a Bernoulli random variable with success probability p :

$$\mathcal{U}|0\rangle = \sqrt{1-p}|0\rangle + \sqrt{p}|1\rangle.$$

We can model \mathcal{U} with a rotation around the Y -axis:

$$\mathcal{U} = R_Y(\theta_p), \quad \theta_p = 2 \arcsin(\sqrt{p}).$$

The Grover operator for this case is particularly simple:

$$\mathcal{Q} = R_Y(2\theta_p),$$

A toy example

Let \mathcal{U} describe a Bernoulli random variable with success probability p :

$$\mathcal{U}|0\rangle = \sqrt{1-p}|0\rangle + \sqrt{p}|1\rangle.$$

We can model \mathcal{U} with a rotation around the Y -axis:

$$\mathcal{U} = R_Y(\theta_p), \quad \theta_p = 2 \arcsin(\sqrt{p}).$$

The Grover operator for this case is particularly simple:

$$\mathcal{Q} = R_Y(2\theta_p),$$

whose powers are very easy to calculate:

$$\mathcal{Q}^k = R_Y(2k\theta_p).$$

A toy example




Figure: QAE circuit for binomial random variable, implemented in Qiskit.





Figure: QAE estimates

QAE for derivative pricing

We saw how to encode the expected payoff μ of a european call option into an operator \mathcal{U}

$$\mathcal{U} |0\rangle_{n+1} = \sqrt{1-a} |\psi_0\rangle_n |0\rangle + \sqrt{a} |\psi_1\rangle_n |1\rangle,$$

after an appropriate re-scaling to the interval $[0, 1]$.



Amplitude estimation can be used to approximate a .

QAE for option pricing




Figure: QAE circuit for option pricing. Implemented in Qiskit.

Notebook time!

Is QAE NISQ-ready?

Is QAE NISQ-ready?




Figure: Depth vs ancillas.

Is QAE NISQ-ready?




Figure: Depth vs ancillas.

IBM Heron can run ≈ 1800 gates within the coherence time of its qubits [IBM].

QAE alternatives

- ▶ Iterative quantum amplitude estimation [[Gri+21](#)].
- ▶ Faster amplitude estimation [[Nak20](#)].
- ▶ Maximum likelihood amplitude estimation [[Suz+20](#)].

QAE alternatives

- ▶ Iterative quantum amplitude estimation [[Gri+21](#)].
- ▶ Faster amplitude estimation [[Nak20](#)].
- ▶ Maximum likelihood amplitude estimation [[Suz+20](#)].




Figure: Convergence rates of different QAE algorithms (Figure 3 in [[Gri+21](#)]).

- ▶ Quadratic speedup over classical Monte Carlo,

Conclusions

- ▶ Quadratic speedup over classical Monte Carlo,
- ▶ Useful in areas of mathematical finance where Monte Carlo is common practice, such as derivative pricing,

Conclusions

- ▶ Quadratic speedup over classical Monte Carlo,
- ▶ Useful in areas of mathematical finance where Monte Carlo is common practice, such as derivative pricing,
- ▶ Not NISQ-ready, but active area of research.

- ▶ [Sta+20]: Portfolios of options, basket options, path-dependent options.
- ▶ [WK24]: Asian and barrier options under Heston model.
- ▶ [ZLW19]: Option Pricing with qGANs.
- ▶ [DL21]: Quantum Support Vector Regression for Disability Insurance.

References I

- [Bra+02] Gilles Brassard et al. *Quantum amplitude amplification and estimation*. 2002. DOI: [10.1090/conm/305/05215](https://doi.org/10.1090/conm/305/05215) (cit. on pp. 38, 39).
- [DL21] Boualem Djehiche and Björn Löfdahl. “Quantum Support Vector Regression for Disability Insurance”. In: *Risks* 9.12 (2021). ISSN: 2227-9091. DOI: [10.3390/risks9120216](https://doi.org/10.3390/risks9120216) (cit. on p. 64).
- [Gri+21] Dmitry Grinko et al. “Iterative quantum amplitude estimation”. In: *npj Quantum Information* 7.1 (Mar. 2021). ISSN: 2056-6387. DOI: [10.1038/s41534-021-00379-1](https://doi.org/10.1038/s41534-021-00379-1) (cit. on pp. 59, 60).
- [IBM] IBM. *IBM Quantum Summit 2023*. URL: <https://www.ibm.com/quantum/summit-2023> (cit. on pp. 56, 57, 58).

- [Kit95] Alexei Y. Kitaev. “Quantum measurements and the Abelian Stabilizer Problem”. In: *Electron. Colloquium Comput. Complex.* TR96 (1995) (cit. on pp. 29, 30, 31, 32, 33).
- [MW23] Nikhil S. Mande and Ronald de Wolf. “Tight Bounds for Quantum Phase Estimation and Related Problems”. In: (2023). arXiv: [2305.04908 \[quant-ph\]](https://arxiv.org/abs/2305.04908) (cit. on pp. 29, 30, 31, 32, 33).
- [Nak20] Kouhei Nakaji. “Faster amplitude estimation”. In: *Quantum Information and Computation* 20.13,14 (Nov. 2020), pp. 1109–1123. ISSN: 1533-7146. DOI: [10.26421/qic20.13-14-2](https://doi.org/10.26421/qic20.13-14-2) (cit. on pp. 59, 60).

- [Sta+20] Nikitas Stamatopoulos et al. “Option Pricing using Quantum Computers”. In: *Quantum* 4 (July 2020), p. 291. ISSN: 2521-327X. DOI: [10.22331/q-2020-07-06-291](https://doi.org/10.22331/q-2020-07-06-291) (cit. on p. 64).
- [Suz+20] Yohichi Suzuki et al. “Amplitude estimation without phase estimation”. In: *Quantum Information Processing* 19.2 (Jan. 2020). ISSN: 1573-1332. DOI: [10.1007/s11128-019-2565-2](https://doi.org/10.1007/s11128-019-2565-2) (cit. on pp. 59, 60).
- [WK24] Guoming Wang and Angus Kan. “Option pricing under stochastic volatility on a quantum computer”. In: *Quantum* 8 (Oct. 2024), p. 1504. ISSN: 2521-327X. DOI: [10.22331/q-2024-10-23-1504](https://doi.org/10.22331/q-2024-10-23-1504) (cit. on p. 64).

- [ZLW19] Christa Zoufal, Aurélien Lucchi, and Stefan Woerner.
“Quantum Generative Adversarial Networks for learning
and loading random distributions”. en. In: *Npj
Quantum Inf.* 5.1 (Nov. 2019) (cit. on p. 64).