
ENIGMA Cortical QC 2.0

Chris Ching, Faisal Rashid, Sophia Thomopoulos, Lianne Schmaal, Premika Boedhoe, Paul Thompson, and Neda Jahanshad




This guide provides:

- Introduction to ENIGMA visual cortical QC
- Introduction to ENIGMA_Cortical_QC_Template.xlsx (standardizes qc records across sites)
- Common QC issues including examples of pass/fail
- Additional issues for optional QC

FreeSurfer Cortical Quality Check

- If you're new to the cortical QC, we highly recommend spending some time viewing some subjects using the fsqc to get familiarized with the FreeSurfer output (see next slide for directions).
- Use nal QC method for checking cortical segment which is good for spotting under/overe mations.
- Use the ENIGMA External QC for checking cortical labels, anatomical boundaries, and confirming arrors aposted on internal QQ.
- Make sure to QC all subjects, net just these flagged in the sutlierder
- Use the ENIGMA_Cortical_QC_Template.xlsx to record your QC ratings.

Freesurfer regions color-coded



Desikan-Killiany Atlas

https://surfer.nmr.mgh.harvard.edu/fswiki/CorticalParcellation

Use the "ENIGMA_Cortical_QC_Template.xlsx" to record your QC

Download the template here: http://enigma.ini.usc.edu/protocols/imaging-protocols/

									Regions color-coded based on FS labels											
	I A	В	C	D	E	F	G	Н		J	K	L	M	N	0	P	Q	R		
	Subject	Internal_QC	External_QC	QC_code	Notes_Questions	temporalpoli	frontalpole	bankssts	superiortem	middletemp	o precentral	postcentral	supramargin	supertorparte	precuneus	cuneus	pericalcarine	Ingual		
2	subj_1	pass	moderate	central;superparietal								L	L	R	R	R				
3	subj_2	fall	fall	pathology	lesion RH	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L	R/L		
4	subj_3	pess	pass	peracingulate																
5	subj. 4	pass	moderate	meninges	motion artifact						R/L	R	R							
6	subj_5	moderate	moderate	poles;pericalcarine		R											R	R		
7	subj_6	moderate	moderate	polesymeninges		R/L						L								
В	subj_7	moderate	moderate	meninges,bankssts;cer	rtral			R	R			L	L							
9	subj_8	pass	pass	ERC/parahippo																
10	subj. 9	pass	moderate	bankssts				L	L	L										
11	subj_10	moderate	pass	poles			L													

First Tab: QC

Subject: Subject ID's

(miornal/External_QC Columns:

- Pass no issues with internal/external QC
- · Hawarata fall particular regions (menator it right; = totto local indep
- Fail severe pathology, image artifacts, registration problems causing severe mislabeling (list R/L for all regions)


Secor Tab: QC Cod Key


- Provinced codes for the "QC_Code" column on first tab to keep track
- Examples of these annon errors are provided in this guide


Good examples




Labels generally correspond to known anatomical boundaries











External QC




Ideal external segmentation (generally correspond to known anatomical boundaries)

External QC: Fail



FAIL (Processing Error)



Fail (pathology)

Moderate: Banks of superior temporal sulcus overestimation

In about 20-30% of subjects, the BanksSTS appears on gyral surface.


In some cases (≈15%) the size of the mislabeled BanksSTS may influence the surrounding ROIs (e.g. superior temporal/middle temporal gyri).


BanksSTS QC Steps:

- Load index.html into browser.
- Press: "Command and + "which zooms in once match the size of your QC images to the size of image above (**Tip: view the PDF version of the guide at 100% and your QC images should be about the same size as the brain above**)
- 3. IF the mislabeled BanksSTS is larger than hand/cursor, THEN fail the BanksSTS + surrounding affected regions (e.g. superior temporal gyrus and/or middle frontal gyrus see examples on next slides.)


OK examples (do not fail BanksSTS)








Pass




Pass


Bad examples BanksSTS (List affected regions in QC sheet)




Fail BanksSTS + STG



Fail BanksSTS + STG + MTG




Fail BanksSTS + STG + MTG




Fail BanksSTS + MTG

STG: superior temporal gyrus; MTG: middle temporal gyrus


Pre/Postcentral Gyrus Issues (<15% cases)






Pre/Postcentral Gyrus Issues (<15% cases)


Misclassifications




Bad examples

Precentral (blue), postcentral (red) and supramarginal gyrus (green)









Pericalcarine Overestimations (<5% cases)

GOOD



 Segmentation confined to calcarine sulcus BAD



- Segmentation overestimates pericalcarine region
- Note failed regions (above: pericalcarine, lingual, and cuneus regions)

Additional Issues

The following includes regions that have been noted as possibly problematic but to a lesser extent than the previous examples.

Choosing to record instances of the following issues may make it easier to perform follow-up analyses.

Uncertain; delineation ventricles vs parahippocampal (light green) and entorhinal cortex (red)

Questionable examples:

A B

Part of the grey matter of the gyri seem not to be allocated to PHG or ERC (but probably to ventricles instead)


However, example A detected in ~70-80% of cases (?), so only exclude when a large part of the gyrus is clearly missing (entorhinal cortex in example B)


Other issues; Entorhinal cortex


- In a large percentage of subjects this region is only partially correctly labeled
- Therefore, we decided to be less stringent with regard to the segmentation of the entorhinal cortex
- We excluded the entorhinal cortex only if more than 50% of the region was poorly segmented (examples on next slides)
 - So only exclude when a LARGE part of the gyrus is clearly missing
- Importantly, possible findings in the entorhinal cortex will be interpreted with caution


Ok examples:

Entorhinal cortex (red)











Bad examples:


Entorhinal cortex (red)




Other issues: cingulate cortex




A paracingulate sulcus (PCS) is present in 30–60% of cases and is more frequently found in the left hemisphere (A and B). This can cause segmentation problems in the cinqulate and surrounding regions.







- In subjects with prominent paracingulate sulcus (example D, E), portions of the cingulate may be underestimated while superior frontal regions may be overestimated.
- **We tend to be less stringent with the QC in this region because of the variability of the anatomy/segmentation. Findings
 in this region should be interpreted with caution and it may be useful to track this issue for follow-up analysis.**

Other issues: Insula (yellow)







- This Freesurfer atlas does not include a subgenual ACC region and sometimes assigns this to the
 insula (yellow) or medial OFC (red/pink) instead Example A. Example B is probably more anatomically
 correct, but judging the accuracy of the insula or medial OFC boundaries becomes difficult.
- Example C shows other somewhat common issues regarding Insula overestimation into the temporal lobes and midline.
- **We tend to be less stringent with the QC of the insula because of the variability of the anatomy/ segmentation and the difficulty with establishing consistent pass/fail criteria. However, it may be useful for some groups to note how often you observe these issues for follow-up analysis.**


Other issues: Superior Parietal Overestimation

Normal




 Segmentation obeys known anatomical boundaries and does not overestimate superior parietal region







 Segmentation overestimates superior parietal region and impacts precuneus/cuneus regions

Other issues: middle/inferior temporal gyrus








- When the middle temporal gyrus looks as if it covers the inferior temporal gyrus, this is usually due to the rotation angle of the brain and is probably Okie Dokie (Pass).
- It is considered normal when the middle and inferior temporal gyri are somewhat overlapping on each other as in the above examples (see examples of non-continuous inferior temporal sulcus).


Supramarginal gyrus overestimation: Extends into superior temporal gyrus



- In some cases the supramarginal gyrus (green) may appear to invade adjacent regions (in this example the superior temporal gyrus).
- We tend to be less stringent with the QC in this region because there is quite a bit of anatomical variability and the exact boundaries dividing the supramarginal gyrus from surrounding regions can be difficult to assess.


Ok examples:


Superior (blue), middle (brown) and inferior (pink) temporal gyrus




Bad examples:

Superior (blue), middle (brown) and inferior (pink) temporal gyrus









QUIZ TIME



















sub-916049




































GOOD LUCK!