

Sharpe ratio and training sample size; $\rho = 0.5$




Solid: mean; shading/bars: ±1 SD; colored dashes: 95th percentile.
100 replications; benchmark fits: N=600, $\rho = 0.5$, $T_{\text{train}} \geq 1600$.


Sharpe ratio and training sample size; N=600




Solid: mean; shading/bars: ±1 SD; colored dashes: 95th percentile.
100 replications; benchmark fits: N=600, $\rho = 0.5$, $T_{\text{train}} \geq 1600$.




Solid: mean; shading/bars: ±1 SD; colored dashes: 95th percentile.
100 replications; benchmark fits: N=600, $\rho = 0.5$, $T_{\text{train}} \geq 1600$.




Solid: mean; shading/bars: ±1 SD; colored dashes: 95th percentile.
100 replications; benchmark fits: N=600, $\rho = 0.5$, $T_{\text{train}} \geq 1600$.



Solid: mean; shading/bars: ±1 SD; colored dashes: 95th percentile.
100 replications; benchmark fits: N=600, $\rho = 0.5$, $T_{\text{train}} \geq 1600$.




100 independent training replications per cell.



Solid: mean; shading: ±1 SD; colored dashes: 95th percentile.
8 paired trials; N=600, $\rho = 0.5$; same kernel, economy and primitive shocks.

Regularization and numerical evaluation



Solid: mean; shading: ±1 SD; colored dashes: 95th percentile.
100 independent training replications; N=600, $\rho = 0.5$.