# Urchin Foraging Rates on Kelp at Different Urchin Densities

**Exploring mutate and distinct functions in RStudio through investigating urchin laboratory data from the SBC LTER.**

Elise LaFreniere

## Set Up

```
# loading in packages
library(tidyverse) # general use
library(janitor) # cleaning data
library(here) # data organization

# reading in data

# urchin density data from DiFiore et al. 2021
urchin <- read.csv(here("data","mesocosm_density_data.csv"))
```

## Cleaning Data & Using Mutate and Distinct Functions

```
# create new object urchin_clean
urchin_clean <- urchin |>
  # clean column names
  clean_names() |>
   # filter for trials with no mortality
  filter(mortality == 0) |>
  # mutate p_r to show the different species of urchins
  rename(species = p_r) |>
  mutate(species = case_when(
    species == "p" ~ "Purple urchin",
    species == "r" ~ "Red urchin"
```

```
  )) |>
# grouping data for summarize
group_by(trial_id, tank, species, urchin_density, total_time, kelp_in, kelp_out) |>
# using summarize to create urchin_biomass
summarize(urchin_biomass = sum(urchin_mass)/1.587) |>
#ungrouping columns
ungroup() |>
# mutate kelp_in and kelp_out columns to the kelp foraging rate of the urchins
 mutate(kelp_consumption_rate = ((kelp_in - kelp_out) / total_time)*24) |>
mutate(rate_per_urchin = (kelp_consumption_rate) / urchin_density) |>
# select for desired columns of interest
select(urchin_density, rate_per_urchin, species, total_time, kelp_consumption_rate, urchin
```

```
`summarise()` has regrouped the output.
i Summaries were computed grouped by trial_id, tank, species, urchin_density,
  total_time, kelp_in, and kelp_out.
i Output is grouped by trial_id, tank, species, urchin_density, total_time, and
  kelp_in.
i Use `summarise(.groups = "drop_last")` to silence this message.
i Use `summarise(.by = c(trial_id, tank, species, urchin_density, total_time,
  kelp_in, kelp_out))` for per-operation grouping (`?dplyr::dplyr_by`) instead.
```

### Mutate (function in dplyr)

The mutate function allows for new columns to be created that are *functions* of existing variables. It can also be used to modify and delete columns.

In this data set, mutate is used to:

- Modify data in the p_r data (which is renamed to species) column to indicate the specific species being looked at instead of the shorthand notation used within the study.
- Create the rate_per_urchin column which is a function of existing variables (kelp_in, kelp_out, total_time, and urchin_density). This new column allows for the rate of kelp consumption over time per individual urchin to be able to be examined.
- Create the kelp_consumption_rate column which is a function of existing variables ((kelp_in, kelp_out, total_time). This new column calculates the rate of kelp consumption over time.

### Other transformations done to the data frame

- clean_names(): used to simplify the column names from the original data file

- filter(): used to remove rows where mortality was seen within an urchin population in a tank
- re_name(): used to rename "p_r" column to "species" for better clarity
- select(): used to select for desired columns within the data frame
- group_by(): showing that rows sharing specified column values belong together
- summarize(): collapses each group into a row, used to add urchin_mass then divide by 1.587 to give biomass in units of gm^-2

## Visualization

```
# base plot
ggplot(data = urchin_clean,
  # x-axis urchin density, y-axis urchin consumption rate
  # color by species
       mapping = aes(x = urchin_density,
                     y = rate_per_urchin,
                     color = species)) +
  # first layer: geom_point
  geom_point(shape = 20,
             alpha = 0.5,
             size = 4) +
  # separating by species
  facet_wrap(~species) +
  # customizing colors
  scale_color_manual(values = c(
    "Purple urchin" = "mediumorchid4",
    "Red urchin" = "firebrick")) +
  # labeling axes and adding title
  labs(x = "Urchin density (per tank)",
       y = "Kelp consumed (gd^-1) per urchin",
       title = "Per-urchin foragaing rate on kelp \nacross different urchin densities") +
  # adding custom theme
  theme_minimal() +
  # removing legend
  theme(legend.position = "none")
```




This visualization shows the kelp foraging rate (gd^-1) per urchin across different urchin densities, separated by urchin species. The red urchin data had fewer observations and were only tested at lower urchin densities than the purple urchin data. Comparing the two panels, the red urchins appear to have higher foraging rates per urchin at lower densities. Overall, foraging rates per urchin were highly variable, particularly at low urchin densities. For red urchins, the foraging rate per urchin appeared to decrease as density increased, yet for the purple urchins, there was no clear trend.

```
# base plot
ggplot(data = urchin_clean,
  # x-axis urchin density, y-axis urchin consumption rate
  # color by species
       mapping = aes(x = urchin_biomass,
                     y = kelp_consumption_rate,
                     color = species)) +
  # first layer: geom_point
  geom_point(shape = 20,
             alpha = 0.5,
             size = 4) +
    # separating by species
  facet_wrap(~species) +
    # customizing colors
  scale_color_manual(values = c(
```

```
    "Purple urchin" = "mediumorchid4",
    "Red urchin" = "firebrick")) +
  labs(x = "Urchin biomass (gm^-2)",
       y = "Kelp consumption rate (gd^-1)",
       title = "Kelp consumption rate versus urchin biomass") +
  # adding custom theme
  theme_minimal() +
  # removing legend
  theme(legend.position = "none")
```




This visualization shows the kelp consumption rate (gd^-1) across different urchin densities, separated by species. The red urchin data had fewer observations and were tested at lower urchin densities than the purple urchin data. In lower densities, kelp consumption rates appear to be similar across both species, though few red urchins had higher rates. Overall, kelp foraging rates appears to increase with increased urchin densities.

## Data Citation

DiFiore, B, M. Rennick, J. Curtis, D. Reed, A. Stier. 2021. SBC LTER: Sea urchin foraging rates on giant kelp ver 1. Environmental Data Initiative. https://doi.org/10.6073/pasta/6af4cc3b0e63b887baf1ae9201e1c Accessed 2026-10-16.