

IPCC Tier 2 Livestock GHG Uncertainty Calculator

Technical Methodology: Equations, Monte Carlo Procedure, and Uncertainty Reporting

CIAT / CGIAR Alliance — ClimateActionNetZero

May 2026

Version 1.0 — May 2026

Developed under the Global Methane Hub initiative

Contents

1	Introduction	4
1.1	Purpose	4
1.2	Scope	4
1.3	IPCC Tier Classification	4
2	Input Parameters	5
3	IPCC Tier 2 Emission Equations	11
3.1	Net Energy for Maintenance	11
3.2	Net Energy for Activity	11
3.3	Net Energy for Growth	11
3.4	Net Energy for Lactation	12
3.5	Net Energy for Work	12
3.6	Net Energy for Pregnancy	12
3.7	Energy Ratios: REM and REG	12
3.8	Gross Energy Intake	13
3.9	Enteric Fermentation Emission Factor	13
3.10	Nitrogen Excretion	13
3.11	Volatile Solids and Manure CH ₄	14
3.12	Manure Management N ₂ O (Direct)	14
3.13	Manure Management N ₂ O (Indirect)	15
3.14	Pasture/Range/Paddock N ₂ O (Direct)	15
3.15	Pasture N ₂ O (Indirect)	16
3.16	GWP-Based CO ₂ eq Conversion	16
4	Monte Carlo Uncertainty Analysis	17
4.1	IPCC Approach 2 Rationale	17
4.2	Probability Distributions	17
4.3	Correlated Sampling	18
4.4	Simulation Procedure	19
4.5	Recommended Iteration Count and Convergence	19
5	Uncertainty Metrics	21
5.1	95% Confidence Interval	21
5.2	Coefficient of Variation	21
5.3	Margin of Error	21
5.4	IPCC Table 3.3 Format	21
6	Uncertainty Decomposition	23
7	Sensitivity Analysis	24

7.1	Standardised Regression Coefficients (SRC)	24
7.2	Partial Rank Correlation Coefficients (PRCC)	24
7.3	Tornado Chart Interpretation	25
8	Correlation Structure	26
8.1	Activity Data Correlations	26
8.2	Emission Factor Correlations	26
8.3	Structural-Defaults Preset (Expert-Elicited)	26
8.4	Cross-Block Correlations	27
8.5	Time-Series Correlation Matrix	27
8.6	Year-to-Year Coefficient Correlation (Trend Mode)	27
9	Validation and Quality Assurance	29
10	References	31

1 Introduction

1.1 Purpose

The **IPCC Tier 2 Livestock GHG Uncertainty Calculator** is a web-based tool that enables national inventory compilers to quantify the uncertainty in cattle greenhouse gas (GHG) emission estimates produced using the IPCC Tier 2 methodology. The tool implements the Monte Carlo (Approach 2) uncertainty framework described in IPCC (2006) Volume 1, Chapter 3, applied to the full emission equation chain of IPCC (2006) Volume 4, Chapter 10.

Under the Paris Agreement Enhanced Transparency Framework (ETF), national inventory reports are expected to include quantitative uncertainty estimates alongside emission totals. This tool automates that process, from parameter sampling to the formatted IPCC Table 3.3 output, for cattle inventories comprising one or more sub-categories (cattle types).

1.2 Scope

The tool covers six emission pathways for cattle:

1. **Enteric fermentation CH_4** — methane produced during enteric fermentation in the rumen
2. **Manure management CH_4** — methane from anaerobic decomposition of manure in storage systems
3. **Manure management N_2O (direct)** — direct nitrous oxide from nitrogen in managed manure
4. **Manure management N_2O (indirect)** — indirect N_2O via atmospheric deposition and leaching from managed manure
5. **Pasture/range/paddock N_2O (direct)** — direct N_2O from dung and urine deposited on pasture
6. **Pasture/range/paddock N_2O (indirect)** — indirect N_2O from dung and urine on pasture

The tool supports multiple cattle sub-categories (e.g. dairy cows, beef bulls, heifers, growing males, calves) within a single inventory run, with results aggregated to national totals.

1.3 IPCC Tier Classification

The tool implements **Tier 2** methods for enteric fermentation (country-specific feed digestibility and body weight data) and **Tier 1 or Tier 2** methods for manure management, depending on the parameters provided. The uncertainty analysis itself follows the **Approach 2** (Monte Carlo) framework, which is the more accurate of the two IPCC uncertainty methods and is recommended when non-linear equation chains are involved.

2 Input Parameters

All parameters are entered via an Excel template (one row per parameter per cattle sub-category) and loaded into the tool. Each parameter requires a central value, an uncertainty specification (distribution type and bounds or percentage), and optional metadata (data source quality, IPCC reference).

Table 1 lists all parameters used in the calculation chain, their units, the IPCC equation in which they appear, the default value pre-loaded in the tool (sourced from the IPCC Guidelines), and the probability distribution typically applied in uncertainty analysis. Where the 2006 Guidelines and the 2019 Refinement give different defaults, both are shown.

Table 1: Input parameters for the IPCC Tier 2 cattle GHG calculation chain — including IPCC defaults and source citations.

Variable	Definition	Unit	IPCC default	IPCC reference (equation / table)	Typical dist.
N	Number of animals (population)	head	<i>n/a</i>	Eq. 10.1 (Tier 1 totals); national activity data	Normal
BW	Mean live body weight	kg/head	275	Eq. 10.3; Annex 10A.2 (Africa cattle)	Normal
Cfi	Coefficient for NE_m	MJ/kg ^{0.75} /d	0.386 (lact.) / 0.322 (non-lact.) / 0.370 (bulls)	Eq. 10.3; Table 10.4 (Updated)	Constant
Tw	Mean winter temperature	°C	20 (inert)	Eq. 10.2 (cold-climate Cfi adjustment)	Constant
Ca	Activity coefficient	—	0.00 (stall) / 0.17 (pasture flat) / 0.36 (hilly)	Eq. 10.4; Table 10.5 (Updated)	Constant
WG	Mean daily weight gain	kg/head/d	0 (non-growing default)	Eq. 10.6	Normal
MW	Mature body weight	kg/head	300	Eq. 10.6; Annex 10A.2	Normal
C	Sex/condition coefficient for NE_g	—	0.8 (female) / 1.0 (castrate) / 1.2 (bull)	Eq. 10.6	Constant

Continued on next page

Table 1 continued

Variable	Definition	Unit	IPCC default	IPCC reference (equation / table)	Typical dist.
Milk	Daily milk yield	kg/head/d	4.0 (Africa)	Eq. 10.8; Annex 10A.1 / 10A.2	Normal/Lognormal
MilkFat	Milk fat content	%	4.0	Eq. 10.8; Annex 10A.1	Normal
pct_calving	Fraction of females calving / yr	—	0.60	2019 Refinement Vol.4 Ch.10 §10.5	Beta
MilkPR	Milk protein content	%	3.3 ($\approx 1.9 + 0.4 \times \%Fat$)	Eq. 10.33 (uses 6.38 milk-protein-to-N); Annex 10A.1–10A.3	Normal
WorkHours	Daily working hours	h/head/d	0 (no work)	Eq. 10.11	Constant
Cp	Pregnancy coefficient	—	0.10 (cattle)	Eq. 10.13; Table 10.7 (Updated)	Constant
DE_pct	Digestible energy (% of GE)	%	55.0 (Africa)	Eq. 10.14–16; Annex 10A	Normal
Ym	Methane conversion factor	%	6.5 (cattle on forage)	Eq. 10.21; Table 10.12	Normal/PERT
MMS_fractions	Fraction of manure in each MMS	—	country-specific (Worksheet 10.A-9)	Eq. 10.23	Uniform/Dirichlet

Continued on next page

Table 1 continued

Variable	Definition	Unit	IPCC default	IPCC reference (equation / table)	Typical dist.
Bo	Maximum CH ₄ producing capacity	m ³ CH ₄ /kg VS	0.13 (2019R “Other regions, low productivity” cattle); 2006 Africa = 0.10	Eq. 10.23; Table 10.16(a) (Updated)	Normal
MCF	Methane conversion factor (MMS)	—	per-MMS & climate (e.g. 80% lagoon tropical)	Eq. 10.23; Table 10.17 (Updated)	Normal/PERT
EF3	Direct-N ₂ O MMS	EF (per kg N ₂ O-N/kg N)	per-MMS (e.g. 0.005 solid storage)	Eq. 10.25; Table 10.21	Lognormal/PERT
Frac_GasMS	Fraction N volatilised (MMS)	—	per-MMS, e.g. 0.45 solid storage (2019R)	Eq. 10.26; Table 10.22 (Updated)	Normal

Continued on next page

Table 1 continued

Variable	Definition	Unit	IPCC default	IPCC reference (equation / table)	Typical dist.
EF4	EF for atmospheric deposition	kg N ₂ O-N/(kg NH ₃ -N+NO _x -N)	0.010 aggregated 2019R (wet 0.014 / dry 0.005); 2006 = 0.010	Eq. 10.27; Vol.4 Ch.11 Table 11.3 (Updated)	PERT
Frac_LeachMS	Fraction N leached (MMS)	—	per-MMS, e.g. 0.02 solid storage	Eq. 10.28; Table 10.23 (2019R)	PERT
EF5	EF for leaching/runoff	kg N ₂ O-N/kg N	0.011 (2019R); 2006 = 0.0075	Eq. 10.29; Vol.4 Ch.11 Table 11.3 (Updated)	PERT
Frac_PRP	Fraction of time / N on pasture	—	country-specific	Vol.4 Ch.11 Eq. 11.5 ($MS_{(T,PRP)}$)	Uniform
EF3_PRP	Direct N ₂ O EF, pasture (cattle/poultry/pigs)	kg N ₂ O-N/kg N	0.004 aggregated 2019R (wet 0.006 / dry 0.002); 2006 = 0.02	Vol.4 Ch.11 Eq. 11.1; Table 11.1 (Updated)	PERT

Continued on next page

Table 1 continued

Variable	Definition	Unit	IPCC default	IPCC reference (equation / table)	Typical dist.
Frac_GasPRP	Fraction N volatilised (PRP)	—	0.21 (2019R Frac-GASM); 2006 = 0.20	Vol.4 Ch.11 Eq. 11.9; Table 11.3 (Updated)	Normal
Frac_LeachPRP	Fraction N leached (PRP)	—	0.24 (2019R wet); 2006 = 0.30; dry = 0	Vol.4 Ch.11 Eq. 11.10; Table 11.3 (Updated)	PERT

3 IPCC Tier 2 Emission Equations

This section presents the complete equation chain implemented in the tool, following IPCC (2006) Volume 4, Chapter 10. All equations are evaluated for each cattle sub-category independently; results are then aggregated to national totals by summing across sub-categories.

3.1 Net Energy for Maintenance

The daily net energy requirement for maintenance (NE_m , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.3:

$$NE_m = C_{fi,adj} \times BW^{0.75}$$

where BW is the mean live body weight (kg head⁻¹) and C_{fi} is a species/condition coefficient (MJ kg^{-0.75} day⁻¹). For open-lot cattle in colder climates the 2006 Guidelines (Vol.4 Ch.10, retained unchanged in the 2019 Refinement) prescribe a linear adjustment of C_{fi} — IPCC Eq.~10.2, “Coefficient for calculating net energy for maintenance” — based on the mean daily winter temperature:

$$C_{fi,adj} = C_{fi} + 0.0048 \times (20 - T_w) \quad \text{when } T_w < 20^\circ\text{C}$$

otherwise $C_{fi,adj} = C_{fi}$. This adjustment increases the maintenance energy requirement by approximately 0.48% per degree below 20°C. The same linear formula is implemented in the IPCC Inventory Software (v2.95).

3.2 Net Energy for Activity

The net energy for physical activity (NE_a , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.4:

$$NE_a = C_a \times NE_m$$

where C_a is a dimensionless activity coefficient that accounts for the terrain and movement patterns of the animal (e.g. 0.00 for housed animals, 0.17 for animals on flat pasture, 0.36 for hilly terrain).

3.3 Net Energy for Growth

The net energy required for growth (NE_g , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.6:

$$NE_g = 22.02 \times \left(\frac{BW}{C \times MW} \right)^{0.75} \times WG^{1.097}$$

where WG is the mean daily weight gain (kg head⁻¹ day⁻¹), MW is the mature body weight (kg), and C is a sex/body condition coefficient (0.8 for females, 1.0 for castrates, 1.2 for bulls). $NE_g = 0$

when $WG \leq 0$ or $MW \leq 0$.

3.4 Net Energy for Lactation

The daily net energy for milk production (NE_l , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.8:

$$NE_l = \text{Milk} \times (1.47 + 0.40 \times \text{MilkFat})$$

where Milk is the daily milk yield (kg head⁻¹ day⁻¹) and MilkFat is the fat content (%). For non-lactating sub-categories, set Milk = 0.

3.5 Net Energy for Work

The net energy for work (NE_w , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.11 (used when cattle perform physical work, e.g. traction or load-carrying):

$$NE_w = 0.10 \times NE_m \times \text{WorkHours}$$

where WorkHours is the mean number of hours of work per day. Set to 0 when animals do not perform work — typical for dairy and beef cattle inventories.

3.6 Net Energy for Pregnancy

The net energy allocated to fetal growth during pregnancy (NE_p , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.13:

$$NE_p = C_p \times NE_m$$

where C_p is a coefficient for the energy cost of pregnancy (0.10 for cattle). The parameter `pct_calving` can pro-rate C_p when not all females in a sub-category are pregnant in the same year ($0 \leq \text{pct_calving} \leq 1$).

3.7 Energy Ratios: REM and REG

The ratio of net energy for maintenance to digestible energy consumed (REM) is given by IPCC Eq. 10.14:

$$REM = 1.123 - \frac{4.092 \times 10^{-3} \times DE_{\%}}{1} + \frac{1.126 \times 10^{-5} \times DE_{\%}^2}{1} - \frac{25.4}{DE_{\%}}$$

The ratio of net energy for growth to digestible energy consumed (REG) is given by IPCC Eq. 10.15:

$$REG = 1.164 - \frac{5.160 \times 10^{-3} \times DE_{\%}}{1} + \frac{1.308 \times 10^{-5} \times DE_{\%}^2}{1} - \frac{37.4}{DE_{\%}}$$

where $DE_{\%}$ is the digestible energy expressed as a percentage of gross energy. Both equations are undefined at $DE_{\%} = 0$ (division by zero) — the tool validates $DE_{\%} > 0$ before running any simulation.

3.8 Gross Energy Intake

The mean daily gross energy intake (GE , MJ head⁻¹ day⁻¹) is given by IPCC Eq. 10.16:

$$GE = \frac{(NE_m + NE_a + NE_l + NE_w + NE_p)/REM + NE_g/REG}{DE_{\%}/100}$$

3.9 Enteric Fermentation Emission Factor

The per-head per-day methane emission factor for enteric fermentation (EF_1 , kg CH₄ head⁻¹ day⁻¹) is given by IPCC Eq. 10.21:

$$EF_1 = \frac{GE \times Y_m}{55.65}$$

where Y_m is the methane conversion factor (fraction of gross energy in feed converted to methane, expressed as %) and 55.65 MJ kg⁻¹ is the energy content of methane. The annual per-sub-category enteric emission is:

$$E_{\text{enteric}} = EF_1 \times N \times 365/1000 \quad (\text{tonnes CH}_4 \text{ yr}^{-1})$$

3.10 Nitrogen Excretion

The daily nitrogen excretion rate (N_{ex} , kg N head⁻¹ day⁻¹) follows IPCC Eq. 10.31:

$$N_{\text{ex}} = N_{\text{intake}} - N_{\text{retained}}$$

Nitrogen intake from feed is:

$$N_{\text{intake}} = \frac{GE}{18.45} \times \frac{CP\%}{6.25 \times 100}$$

Nitrogen retained in body tissue and milk is calculated separately. For milk, the nitrogen content uses the **Jones factor for casein (6.38)** — distinct from the generic protein factor (6.25) used for feed and body tissue. The 6.38 factor and the 6.25 factor both appear inside IPCC 2006 Vol.4 Ch.10 Eq. 10.33 (N retention rates for cattle); the same equation is retained, with updated form, in the

2019 Refinement (also numbered 10.33):

$$N_{\text{milk}} = \frac{\text{Milk} \times \text{MilkProtein}\%}{6.38 \times 100}$$

The annual nitrogen excreted per sub-category ($N_{\text{ex,annual}}$, kg N yr⁻¹) is:

$$N_{\text{ex,annual}} = N_{\text{ex}} \times N \times 365$$

3.11 Volatile Solids and Manure CH₄

Total manure CH₄ emissions are given by IPCC 2006 Vol.4 Ch.10 Eq. 10.22 (the top-level emissions equation, $EF \times \text{population} \times 10^{-6}$). The per-head per-year emission factor is given by Eq. 10.23, and the daily volatile-solid (VS) excretion rate that feeds Eq. 10.23 is given by Eq. 10.24:

$$EF_{\text{mm,CH}_4} = VS \times 365 \times B_o \times 0.67 \times \sum_s (MS_s \times MCF_s)$$

where B_o is the maximum methane producing capacity (m³ CH₄ kg⁻¹ VS), 0.67 is the conversion factor from m³ CH₄ to kg CH₄ (the density of methane at standard conditions, kg m⁻³), MS_s is the fraction of manure managed in system s , and MCF_s is the methane conversion factor for system s (fraction of B_o). The 2019 Refinement Table 10.17 (Updated) provides MCF defaults across 11 manure management systems — eight inherited from the 2006 Guidelines plus four added in 2019 (anaerobic digester / biogas, aerobic treatment, solid storage – covered/compacted, and burned for fuel). Default B_o values for cattle are given in Table 10.16a. Per-MMS uncertainty in MCF and in the manure fractions MS_s is sampled independently.

3.12 Manure Management N₂O (Direct)

Direct N₂O emissions from manure management are given by IPCC 2006 Vol.4 Ch.10 Eq. 10.25 (the indirect-pathway losses are calculated separately in Eq. 10.26–10.29; see next sub-section):

$$E_{\text{direct,mm}} = N_{\text{ex,annual}} \times EF_3 \times \frac{44}{28} \quad (\text{kg N}_2\text{O yr}^{-1})$$

where EF_3 is the emission factor for direct N₂O (kg N₂O-N kg⁻¹ N) and 44/28 converts N₂O-N to N₂O. For mixed MMS systems, EF_3 is computed as the weighted average of per-MMS factors:

$$EF_3 = \sum_s MS_s \times EF_{3,s}$$

3.13 Manure Management N₂O (Indirect)

Indirect N₂O from managed manure arises from two pathways. Each pathway is split between an N-loss equation (in Vol.4 Ch.10) and an N₂O equation (in Vol.4 Ch.10, drawing EFs from Vol.4 Ch.11 Table 11.3):

Atmospheric deposition of volatilised N — Frac_GasMS from Eq. 10.26 multiplied by EF_4 in Eq. 10.27:

$$E_{\text{vol,mm}} = N_{\text{ex,annual}} \times \text{Frac_GasMS} \times EF_4 \times \frac{44}{28}$$

Leaching and runoff — Frac_LeachMS from Eq. 10.28 multiplied by EF_5 in Eq. 10.29:

$$E_{\text{leach,mm}} = N_{\text{ex,annual}} \times \text{Frac_LeachMS} \times EF_5 \times \frac{44}{28}$$

where EF_4 (kg N₂O-N kg⁻¹ NH₃-N + NO_x-N volatilised) and EF_5 (kg N₂O-N kg⁻¹ N leached/runoff) are from IPCC Vol.4 Ch.11 Table 11.3. In the 2006 Guidelines $EF_4 = 0.010$ (range 0.002–0.05) and $EF_5 = 0.0075$ (range 0.0005–0.025). The 2019 Refinement keeps the aggregated $EF_4 = 0.010$ (range 0.002–0.018) and additionally provides climate-disaggregated values of 0.014 in wet climates (range 0.011–0.017) and 0.005 in dry climates (range 0.000–0.011); EF_5 is updated to 0.011 (range 0.000–0.020) with no climate disaggregation. Frac_GasMS and Frac_LeachMS are MMS- and country-specific fractions (defaults in Vol.4 Ch.10 Tables 10.22 and 10.23 of the 2019 Refinement); on the soil side of Eq.~11.9/11.10 the same role is played by FracGASM and FracLEACH-(H) in Vol.4 Ch.11 Table 11.3.

3.14 Pasture/Range/Paddock N₂O (Direct)

The nitrogen deposited on pasture by grazing animals (F_{PRP}) is the per-sub-category form of IPCC Vol.4 Ch.11 Eq.~11.5 (N in urine and dung deposited by grazing animals on PRP):

$$F_{\text{PRP}} = N_{\text{ex,annual}} \times \text{Frac_PRP}$$

where Frac_PRP is the fraction of total annual N excretion deposited on pasture (denoted $MS_{(T,\text{PRP})}$ in IPCC notation). Direct N₂O from pasture/range/paddock falls under managed soils (IPCC Volume 4, Chapter 11), not Chapter 10. It is given by IPCC Eq.~11.1 (with F_{PRP} as the input N source):

$$E_{\text{direct,PRP}} = F_{\text{PRP}} \times EF_{3,\text{PRP}} \times \frac{44}{28}$$

where $EF_{3,\text{PRP}}$ is the emission factor for direct N₂O from PRP given in IPCC Vol.4 Chapter 11, Table 11.1. The 2006 value is 0.02 kg N₂O-N kg⁻¹ N (cattle, poultry, pigs combined). The 2019 Refinement updates this to $EF_{3,\text{PRP},\text{CPP}}$ for cattle, poultry and pigs with an aggregated default of 0.004 kg N₂O-N kg⁻¹ N (range 0.000–0.014). When the climate is known, the disaggregated values are 0.006 in wet climates (range 0.000–0.027) and 0.002 in dry climates (range 0.000–0.007). For

sheep and other animals ($EF_{3PRP,SO}$) the 2019 Refinement gives 0.003 (range 0.000–0.010), with no climate disaggregation.

3.15 Pasture N₂O (Indirect)

Indirect N₂O from pasture deposition follows IPCC Vol.4 Ch.11 Eq. 11.9 (volatilisation pathway) and Eq. 11.10 (leaching/runoff pathway), using the same pathway structure as managed manure:

$$E_{\text{indirect,PRP}} = F_{\text{PRP}} \times (\text{Frac_GasPRP} \times EF_4 + \text{Frac_LeachPRP} \times EF_5) \times \frac{44}{28}$$

3.16 GWP-Based CO₂eq Conversion

All emission pathways are converted to CO₂ equivalents using Global Warming Potentials (GWP). The tool supports three IPCC assessment report baselines:

Basis	CH ₄ GWP	N ₂ O GWP	Reference
AR4 (100-yr)	25	298	IPCC AR4 WG1, Ch.2 Table 2.14
AR5 (100-yr)	28	265	IPCC AR5 WG1, Ch.8 Table 8.7
AR6 (100-yr, non-fossil CH ₄)	27.0	273	IPCC AR6 WG1, Ch.7 Table 7.15 (and Table 7.SM)

Total CO₂eq = (total CH₄ tonnes × GWP_{CH₄}) + (total N₂O tonnes × GWP_{N₂O}).

4 Monte Carlo Uncertainty Analysis

4.1 IPCC Approach 2 Rationale

The IPCC (2006) Volume 1, Chapter 3 defines two approaches to quantifying inventory uncertainty:

- **Approach 1 (error propagation):** Combines individual parameter uncertainties using simple algebraic rules. Suitable when the emission equation is a product or sum of independent terms. Underestimates uncertainty when equations are non-linear or inputs are correlated.
- **Approach 2 (Monte Carlo simulation):** Propagates uncertainty through the full equation chain by repeatedly sampling all parameters from their distributions. Exact for any equation structure. Recommended for Tier 2 livestock inventories because the GE equation chain is non-linear in BW , WG , and $DE\%$.

This tool implements Approach 2. The simulation samples all input parameters simultaneously for each iteration, evaluates the full equation chain from NE_m through to total CO_{2eq} , and records the output. The resulting distribution of outputs characterises the uncertainty.

4.2 Probability Distributions

Each parameter is assigned a probability distribution reflecting the state of knowledge about its true value. The tool supports:

Distribution	Parameterisation	Typical use
Normal	mean μ , SD σ	Body weight, milk yield, population counts with symmetric survey error
Positive-Normal	Normal truncated at zero	Non-negative symmetric parameters where negative draws are implausible
Lognormal	log-mean, log-SD (fitted from mean and CV%)	Strictly positive, right-skewed; milk yield in heterogeneous herds; some EFs
Beta	rescaled Beta(α, β) matching the supplied mean and bounds	Fractions in $[0, 1]$ with mode strictly in the interior; pct_calving, Cp
Truncated-Normal [0,1]	Normal clipped to $[0, 1]$	Fractions in $[0, 1]$ when a symmetric error around the mean is appropriate
PERT	minimum, most-likely, maximum	Emission factors with expert-elicited bounds; EF3_PRP, EF4, EF5, Bo, MCF
Triangular	minimum, most-likely, maximum	Similar to PERT but with sharper tails; Ca, C_growth
Uniform	lower, upper	Parameters with only the plausible range known; MMS fractions
Constant	fixed value	Parameters treated as known exactly (uncertainty_pct = 0)

When the user supplies **uncertainty_pct** (a symmetric 95% half-width as a percentage of the mean), the tool derives the distribution parameters as:

$$\sigma = \frac{\text{uncertainty_pct} \times \mu}{100 \times 1.96}$$

for a normal distribution, or equivalent transformations for lognormal and PERT.

4.3 Correlated Sampling

When parameters are correlated (e.g. body weight and mature weight are positively correlated; diet quality and methane conversion factor may be negatively correlated), ignoring correlation underestimates or overestimates combined uncertainty. The tool applies the same correlated-sampling procedure to every path that supplies a correlation matrix — the structural-defaults preset, a manually uploaded matrix, and the matrix estimated automatically from a time-series upload all flow through the same algorithm.

Procedure (per IPCC Volume 1 Chapter 3 §3.2.3.2):

1. For each parameter j , draw n_{iter} independent samples from the parameter's own marginal distribution (normal, PERT, lognormal, beta, triangular, uniform, or constant, exactly as

specified in the template).

2. Reorder the columns of the resulting $n_{\text{iter}} \times p$ sample matrix so that the realised Spearman rank correlation between columns matches the target matrix Σ (implemented via `mc2d::cornode`, which performs the restricted-pairing operation IPCC §3.2.3.2 cites under “restricted pairing methods that are included in many software packages”).

This procedure is distribution-free: the marginal shapes are preserved exactly, regardless of whether the user has mixed normal / lognormal / PERT / beta parameters in the same correlation block. The target Σ is interpreted as a Spearman rank correlation matrix throughout — the same convention the time-series matrix is computed under (Spearman rank correlation of the user’s annual values, after detrending; see Section “Correlation Structure”).

If Σ is not positive definite (e.g. a manually entered matrix that turns out to be inconsistent), the tool projects it onto the nearest positive-definite correlation matrix via `Matrix::nearPD(corr = TRUE)` and emits a warning whenever any individual entry is shifted by more than 0.01. The tool does *not* silently fall back to uncorrelated sampling.

The tool supports both within-block and across-block correlations. The default scope is independence; a single unified correlation matrix spanning AD + coefficients is used whenever cross-block pairs are specified.

4.4 Simulation Procedure

For each Monte Carlo iteration $i = 1, \dots, n_{\text{iter}}$:

1. Sample all parameters from their distributions (with correlation structure if supplied).
2. For each cattle sub-category k :
 - a. Compute $NE_m^{(k)}, NE_a^{(k)}, NE_g^{(k)}, NE_l^{(k)}, NE_w^{(k)}, NE_p^{(k)}$.
 - b. Compute $REM^{(k)}$ and $REG^{(k)}$ from $DE_{\%}^{(k)}$.
 - c. Compute $GE^{(k)}, EF_1^{(k)}, N_{\text{ex}}^{(k)}, VS^{(k)}$.
 - d. Compute emission totals for all six pathways.
3. Sum across all sub-categories to obtain national pathway totals and total CO₂eq.
4. Record all outputs for iteration i .

After all iterations, compute summary statistics from the distribution of recorded outputs.

4.5 Recommended Iteration Count and Convergence

Monte Carlo estimates converge as $n_{\text{iter}} \rightarrow \infty$. For practical use:

- **1,000 iterations** — suitable only for a quick test run to verify the model runs without errors. Do not report results from 1,000 iterations.
- **10,000 iterations** — minimum recommended for any result intended for use. Convergence is generally adequate for inventories with up to 5 sub-categories and 30 parameters.
- **25,000–30,000 iterations** — recommended for final reporting, especially when correlations

are enabled or the inventory has many sub-categories or asymmetric distributions.

- **50,000 iterations** — high-precision run; use when the 95% CI bounds need to be stable to two significant figures.

Convergence check: Re-run with a different random seed (e.g. 123 and 456). If the CV% changes by more than 1–2 percentage points, increase the number of iterations.

5 Uncertainty Metrics

The following metrics are reported for each emission pathway and for the total CO₂eq:

5.1 95% Confidence Interval

The 95% CI is defined as the 2.5th and 97.5th percentiles of the Monte Carlo output distribution:

$$CI_{95} = [Q_{0.025}, Q_{0.975}]$$

This is a simulation-based (quantile) interval — it does not assume a normal distribution and is therefore valid for asymmetric emission distributions.

5.2 Coefficient of Variation

The coefficient of variation (CV%) expresses uncertainty relative to the mean:

$$CV\% = \frac{\sigma}{\bar{x}} \times 100$$

where σ is the standard deviation and $\bar{x}$ is the mean of the Monte Carlo output distribution. CV% is the primary metric used in IPCC Table 3.3 uncertainty reporting. It is undefined (reported as NA) when the mean emission is zero (e.g. pasture N₂O for fully housed systems).

5.3 Margin of Error

The margin of error (MoE%) is the half-width of the 95% CI expressed as a percentage of the mean:

$$MoE\% = \frac{Q_{0.975} - Q_{0.025}}{2 \times \bar{x}} \times 100$$

Unlike CV%, MoE% is derived from the quantiles of the actual distribution and correctly captures skewness. For a perfectly normal distribution, $MoE\% \approx 1.96 \times CV\%$.

5.4 IPCC Table 3.3 Format

IPCC (2006) Volume 1, Table 3.3 requires reporting uncertainty for each emission category as follows:

Source / Sink Category	Emission (Gg CO ₂ eq)	AD Uncertainty (%)	EF Uncertainty (%)	Combined Uncertainty (%)
Enteric fermentation CH ₄	...	CV% from AD-only run	CV% from EF-only run	CV% from combined run
...				

The tool generates this table automatically after each simulation run.

6 Uncertainty Decomposition

To separate the contributions of activity data (AD) uncertainty and emission factor (EF) uncertainty to the combined uncertainty, the tool performs three simulation runs:

1. **Combined run** — all parameters varied according to their distributions. This is the primary result.
2. **AD-only run** — emission factors fixed at their central values ($\sigma = 0$); only AD parameters vary. CV% from this run = AD uncertainty contribution.
3. **EF-only run** — AD parameters fixed at their central values; only emission factors vary. CV% from this run = EF uncertainty contribution.

This approach correctly handles multi-group inventories by running all three simulations through the full multi-sub-category equation chain and aggregating to national totals before computing metrics.

The decomposition reveals whether uncertainty reduction efforts should focus on improving animal population surveys (AD) or on refining emission factor estimates (EF).

7 Sensitivity Analysis

Sensitivity analysis identifies which input parameters have the greatest influence on the total emission estimate. The tool implements two complementary methods.

7.1 Standardised Regression Coefficients (SRC)

SRC is based on fitting a multiple linear regression of the standardised output on all standardised inputs:

$$Y^* = \beta_0 + \sum_j \beta_j X_j^* + \varepsilon$$

where $Y^* = (Y - \bar{Y})/\sigma_Y$ and $X_j^* = (X_j - \bar{X}_j)/\sigma_{X_j}$. The SRC for parameter j is the coefficient β_j from this regression.

Properties: - A positive SRC indicates that higher values of the parameter produce higher emissions. - The magnitude $|\beta_j|$ reflects the relative importance of parameter j . - SRC values range from approximately -1 to $+1$ when a single parameter dominates. - The R^2 of the regression indicates what fraction of the total variance in the output is explained by the linear model. High R^2 (> 0.9) indicates SRC is a reliable measure; low R^2 suggests non-linearity. - SRC requires only a single regression fit, making it computationally efficient for any number of parameters.

7.2 Partial Rank Correlation Coefficients (PRCC)

PRCC is a rank-based method that is robust to non-linearity and non-normal distributions:

1. Replace all input columns and the output column with their ranks.
2. For each parameter j :
 - a. Regress $\text{rank}(X_j)$ on all other ranked inputs; compute residuals e_{X_j} .
 - b. Regress $\text{rank}(Y)$ on all other ranked inputs; compute residuals e_Y .
 - c. $\text{PRCC}_j = \text{cor}(e_{X_j}, e_Y)$.

PRCC measures the unique rank-correlation between each input and the output after removing the shared linear effect of all other inputs. It is more appropriate than SRC when the relationship between inputs and output is monotone but non-linear, or when input distributions are strongly skewed.

Computational note: PRCC requires one regression fit per parameter. For large inventories (more than 40 combined parameters across all sub-categories), PRCC computation can take several minutes. In this case, the tool automatically falls back to SRC for the aggregate sensitivity result; PRCC remains available for individual sub-category analyses.

7.3 Tornado Chart Interpretation

The tornado chart displays the top parameters ranked by $|\text{SRC}|$ or $|\text{PRCC}|$:

- **Green bars** (positive coefficient) — increasing this parameter increases total emissions; reduction in uncertainty would directly reduce total emission uncertainty.
- **Red bars** (negative coefficient) — increasing this parameter decreases total emissions.
- **Dark (slate) bars** — IPCC-defined coefficients that cannot be improved by the national inventory compiler (e.g. GWP values, Bo, MCF default values). Uncertainty in these parameters is irreducible without IPCC guidance changes.
- **Bright (green/red) bars** — user-reducible parameters where better national data could meaningfully reduce inventory uncertainty.

8 Correlation Structure

8.1 Activity Data Correlations

Correlations between activity data parameters (e.g. body weight and mature weight, milk yield and milk fat content) are entered in the Tab~4 *Activity-Data Correlations* card. Four modes are available:

- **No correlations** (independence) — the default.
- **Structural-defaults preset** — a small set of expert-elicited pairs drawn from the IPCC equation structure (see Section “Structural-Defaults Preset” below).
- **From template (auto, time-series)** — the matrix is computed automatically from the `Parameter_TimeSeries` sheet of the upload using **Spearman rank correlation** after detrending (first differences by default; linear and raw alternatives available). Spearman is preferred over Pearson because (i) it is robust to skewed marginals (Ym, Bo, milk yield), (ii) it matches the rank-correlation target the sampler reproduces (per IPCC Vol.1 Ch.3 §3.2.3.2), and (iii) it is the natural choice when the user wants the *rank* dependence preserved rather than a linear product-moment relationship that can be distorted by outliers.
- **Manual entry** — the user uploads a square symmetric CSV.

8.2 Emission Factor Correlations

Emission factors that are structurally linked (e.g. EF4 is applied to volatilised N, whose magnitude is itself driven by `Frac_GasMS`) can be assigned correlations in the Tab~4 *Coefficient Correlations* card. The same four modes are offered; **no correlations'' is the default. The same correlated-sampling procedure used for activity data (SectionCorrelated Sampling''** above) is applied to the coefficient block — no special-case sampler.

8.3 Structural-Defaults Preset (Expert-Elicited)

The tool provides a **structural-defaults preset** correlation matrix as a middle ground between full independence and user-specified correlations. This preset assigns non-zero correlations to a small set of structurally linked pairs drawn from the IPCC equation structure and the published livestock literature (e.g. body weight and mature weight, via the growth-curve asymptote relation; milk yield and milk fat content, via the dilution effect). *These values are expert-elicited; the IPCC publishes no numerical correlation coefficients for any livestock parameter pair.* All other pairs are set to zero. Hover any non-zero cell in the heatmap for the per-pair source citation. This is appropriate when no empirical correlation data are available but the inventory compiler wants to avoid the unrealistic assumption of full independence.

May 2026 audit re-review. DE ↔ Ym was strengthened from \$-0.40 to \$-0.50 to better match IPCC 2019 Refinement Eq 10.21 and the meta-analysis evidence (Niu et al. 2018 Global Change Biology; Hristov et al. 2013). Cfi ↔ Ca was lowered from 0.60 to 0.30: the previous 0.60 invoked an

estimation-covariance argument that does not apply because Cfi and Ca are independent inputs in IPCC Eq 10.3 / 10.4, not jointly estimated. The earlier $N \leftrightarrow BW = 0.30$ cross-block pair was removed (no defensible cross-country source). The resulting preset matrix is positive-definite without nearest-PD repair (smallest eigenvalue 0.06).

8.4 Cross-Block Correlations

The preset and the manual / time-series matrices support cross-block correlations (i.e. correlations between an activity-data parameter and a coefficient — for example $BW \leftrightarrow Cfi$). This is implemented via a single unified correlation matrix that spans the union of all parameter names; the sampler draws from one multivariate distribution covering both blocks rather than two independent block-level distributions. Independence within and across blocks is the default.

8.5 Time-Series Correlation Matrix

When the user uploads a `Parameter_TimeSeries` sheet, the correlation matrix is computed automatically as a **Spearman** rank correlation of the annual values, after **detrending** (first differences by default, with a UI option to switch to linear detrend or raw series). Detrending separates shared long-run growth — which is not the year-to-year joint uncertainty Approach 2 is meant to capture — from genuine parameter co-movement. The resulting matrix is then projected onto the nearest positive-definite correlation matrix via `Matrix::nearPD(corr = TRUE)` so it can be supplied directly to the sampler described in the “Correlated Sampling” section above.

8.6 Year-to-Year Coefficient Correlation (Trend Mode)

When the tool is run in **Trend mode** (multi-year inventory), an additional decision arises: within a single Monte Carlo iteration, how should the IPCC coefficients (Cfi, Ca, Ym, Bo, MCF, EF3, EF4, EF5, ...) move from one year to the next? IPCC Volume 1 Chapter 3 Section 3.2.3.2 (Trend Analysis) and 2019 Refinement §3.2.2.4 discuss this explicitly: emission factor uncertainties are typically estimated once for an inventory series and reused across years (correlated), while activity data is normally re-estimated each year and treated as independent across years.

The tool exposes three options on the Simulate tab when Trend mode is selected. Activity data (N , BW , WG , Milk, ...) is re-sampled fresh each year regardless of this setting:

Option	What happens inside one Monte Carlo iteration
Fully correlated (default)	The coefficient draw for year t is the <i>same</i> as for year 0. If Y_m is drawn at 6.2% for year 0, it is 6.2% for every other year in that iteration. The trend then reflects activity-data changes only — coefficient uncertainty cancels when subtracting two years. Matches IPCC 2019 §3.2.2.4 default behaviour and the Ogle et al. (2003) “identical random seed” construction cited in V1 Ch3 Box 3.2.
Partial (AR(1), $\rho = 0.7$)	Coefficients drift slowly between years with a rank correlation of $\rho^{ i-j }$ between year i and year j ($\rho = 0.7$). Independent samples are drawn for each year from the coefficient’s marginal and then reordered across years so the realised Spearman rank correlation matches the AR(1) target — the same restricted-pairing procedure used within a year (per IPCC Vol.1 Ch.3 §3.2.3.2), applied across years. Neighbouring years share most of the same observational basis but the coefficient can move over a decade. Appropriate when the EF is re-estimated periodically (e.g. every 3–5 years).
Independent	Each year gets a fresh, independent coefficient draw. Maximises the EF contribution to trend uncertainty. Realistic only when every emission factor is genuinely re-measured with new field data every year, which is uncommon for national inventories.

The choice is implemented in R/trend_tab.R. The `full''` branch pre-samples a single $n_{\text{iter}} \times n_{\text{coef}}$ matrix and reuses it for every year.

The `partial''` branch builds an $n_{\text{iter}} \times n_{\text{years}}$ rank-correlated draw for each coefficient against an AR(1) target rank correlation matrix, then runs each year’s MC with the corresponding column slice. The “independent” branch re-samples coefficients inside every per-year call.

Direction of effect. For an emission factor whose uncertainty is largely systematic, full correlation produces the *smallest* trend uncertainty (because the EF cancels) and independent sampling the *largest*. IPCC V1 Ch3 p. 25 explicitly notes that positive correlations in uncertainties when comparing two years as part of trend analysis will decrease uncertainty in the trend. Choosing independent when the EFs are in fact reused will over-state trend uncertainty; choosing full when the EFs are in fact re-measured each year will under-state it. The IPCC default (full) is the conservative choice for the typical case where the inventory compiler is using IPCC defaults or a single national estimation programme.

9 Validation and Quality Assurance

The tool includes a QA/QC module (`R/utlis_qaqc.R`) that runs a series of checks against the loaded parameter data before simulation. Each check returns one of five statuses:

Status	Trigger	What the user sees
Pass	Value is within $\pm 50\%$ of the IPCC default for that parameter, and all structural checks pass.	Green tick; no action needed.
Warn	Value deviates 50–200% from the IPCC default; or implausible bounds ($BW > 1500$ kg, $Y_m > 12\%$); or a fractional parameter is using an unbounded distribution.	Amber row; the message either suggests a fix or asks the user to document the country-specific deviation.
Fail	Value deviates more than 200% from the IPCC default; or $DE\% \leq 0$ or ≥ 100 ; or $lower > upper$; or an MMS fraction-sum is more than 1% off 100%; or a Beta is configured with mean outside $(0, 1)$.	Red row; simulation will still run but the result will be flagged in the report.
Missing	A core parameter was not supplied in the upload; auto-filled with the IPCC default.	Grey row; the message names the IPCC table or equation used and, for context-dependent parameters (EF3_PRP, EF4, Frac_LeachPRP, Bo, MCF, Ym, Cfi, Ca), appends an explicit hint of which alternative IPCC table row should be reviewed for the inventory's climate / production system / animal class.
Info	Reserved for future advisory messages.	Blue row.

The benchmark comparison (Pass / Warn / Fail thresholds) is anchored on the `ipcc_default` column of `PARAM_CATALOGUE`, which is sourced from Vol.4 Ch.10 (Tables 10.4, 10.5, 10.7, 10.11, 10.12, 10.16(a), 10.17, 10.21, 10.22, 10.23) and Vol.4 Ch.11 (Tables 11.1, 11.3). Auto-filled parameters are listed in a banner on Tab~1 immediately after data upload; they should be replaced with country-specific values wherever possible, because both the central estimate and the propagated uncertainty

depend on them.

10 References

- Frey, H.C., and Rhodes, D.S. (1998). Characterizing, Simulating, and Analyzing Variability and Uncertainty: An Illustration of Methods Using an Air Toxics Emissions Example. *Human and Ecological Risk Assessment*, 4(4), 762–797.
- IPCC (2006). 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 1: General Guidance and Reporting, Chapter 3: Uncertainties. Eggleston, H.S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K. (Eds). IGES, Japan.
- IPCC (2006). 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4: Agriculture, Forestry and Other Land Use, Chapter 10: Emissions from Livestock and Manure Management. IGES, Japan.
- IPCC (2019). 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4: Agriculture, Forestry and Other Land Use, Chapter 10 (Emissions from Livestock and Manure Management) and Chapter 11 (N₂O Emissions from Managed Soils, and CO₂ Emissions from Lime and Urea Application). Calvo Buendia, E., Tanabe, K., Kranjc, A., et al. (Eds). IPCC, Switzerland.
- IPCC (2019). 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 1: General Guidance and Reporting, Chapter 3: Uncertainties. IPCC, Switzerland.
- Iman, R.L., and Conover, W.J. (1982). A distribution-free approach to inducing rank correlation among input variables. *Communications in Statistics — Simulation and Computation*, 11(3), 311–334. (The restricted-pairing algorithm explicitly cited in IPCC 2006 Vol.1 Ch.3 §3.2.3.2 and used by this tool whenever a Spearman correlation matrix is estimated from a time series.)
- Jones, D.B. (1931). Factors for Converting Percentages of Nitrogen in Foods and Feeds into Percentages of Proteins. *USDA Circular* No. 183. United States Department of Agriculture, Washington, D.C.
- Karimi-Zindashty, Y., MacDonald, J.D., Nitschelm, J., Del Grosso, S.J., and Desjardins, R.L. (2012). Sources of uncertainty in the Canadian national livestock greenhouse gas inventory. *Journal of Agricultural Science*, 150(6), 677–690.
- Milne, E., Sessitsch, A., Wilson, J., et al. (2014). Estimating uncertainty in pasture-based dairy CH₄ emissions using Monte Carlo simulation. *Animal*, 8(2), 313–320.
- Monni, S., Syri, S., and Savolainen, I. (2007). Uncertainties in the Finnish greenhouse gas emission inventory. *Environmental Science and Policy*, 7(2), 87–98.
- Penman, J., Kruger, D., Galbally, I., Hiraishi, T., Nyenzi, B., Emmanuel, S., Buendia, L., Hoppaus, R.,

Martinsen, T., Meijer, J., Miwa, K., and Tanabe, K. (2000). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.

Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M., and Tarantola, S. (2008). *Global Sensitivity Analysis: The Primer*. Wiley, Chichester.

IPCC (2021). Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report (AR6), Chapter 7 Table 7.15 and Table 7.SM.7: 100-year GWP values for CH₄ (non-fossil) and N₂O. Cambridge University Press.

R packages used by the simulation engine.

Pouillot, R., and Delignette-Muller, M.-L. (2010). *Evaluating variability and uncertainty separately in microbial quantitative risk assessment: a primer using R and a freely-available package, mc2d*. International Journal of Food Microbiology, 142(3), 330–340. (`mc2d::cornode` — restricted-pairing reordering used to reproduce the target rank correlation matrix per IPCC Vol.1 Ch.3 §3.2.3.2.)

Bates, D., and Maechler, M. (2024). *Matrix*: Sparse and Dense Matrix Classes and Methods. R package. (`Matrix::nearPD` — projection to the nearest positive-definite correlation matrix when a user-supplied Σ is not positive-definite.)

Pouillot, R., and Delignette-Muller, M.L. (2010). Evaluating variability and uncertainty separately in microbial quantitative risk assessment using two R packages. *International Journal of Food Microbiology*, 142(3), 330–340. (`mc2d` — provides `cornode` for Iman–Conover restricted pairing, and `rtriang` / `rpert` for triangular and PERT marginals.)

This document was prepared by CIAT / CGIAR Alliance — ClimateActionNetZero.

Funded by the Global Methane Hub. Version 1.0, May 2026.
