

Inverse potentials of one-body densities

Louis Garrigue
Cergy's university

December 6th, 2024
Oslo

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

N -body quantum mechanics

- No spin, static, space \mathbb{R}^d , electrons

N -body quantum mechanics

- No spin, static, space \mathbb{R}^d , electrons
- States are $\Psi \in L_a^2 \left((\mathbb{R}^d)^N, \mathbb{C} \right)$, with $\int_{\mathbb{R}^{dN}} |\Psi|^2 = 1$

N -body quantum mechanics

- No spin, static, space \mathbb{R}^d , electrons
- States are $\Psi \in L^2_a \left((\mathbb{R}^d)^N, \mathbb{C} \right)$, with $\int_{\mathbb{R}^{dN}} |\Psi|^2 = 1$
- $\Psi(\dots, x_i, \dots, x_j, \dots) = -\Psi(\dots, x_j, \dots, x_i, \dots)$

N -body quantum mechanics

- No spin, static, space \mathbb{R}^d , electrons
- States are $\Psi \in L_a^2\left((\mathbb{R}^d)^N, \mathbb{C}\right)$, with $\int_{\mathbb{R}^{dN}} |\Psi|^2 = 1$
- $\Psi(\dots, x_i, \dots, x_j, \dots) = -\Psi(\dots, x_j, \dots, x_i, \dots)$
- Hamiltonian : operator of $L_a^2\left((\mathbb{R}^d)^N, \mathbb{C}\right)$

$$H_N(v) = \sum_{i=1}^N -\Delta_{x_i} + \sum_{1 \leq i < j \leq N} w(x_i - x_j) + \sum_{i=1}^N v(x_i)$$

N -body quantum mechanics

- No spin, static, space \mathbb{R}^d , electrons
- States are $\Psi \in L_a^2\left((\mathbb{R}^d)^N, \mathbb{C}\right)$, with $\int_{\mathbb{R}^{dN}} |\Psi|^2 = 1$
- $\Psi(\dots, x_i, \dots, x_j, \dots) = -\Psi(\dots, x_j, \dots, x_i, \dots)$
- Hamiltonian : operator of $L_a^2\left((\mathbb{R}^d)^N, \mathbb{C}\right)$

$$H_N(v) = \sum_{i=1}^N -\Delta_{x_i} + \sum_{1 \leq i < j \leq N} w(x_i - x_j) + \sum_{i=1}^N v(x_i)$$

- Ground states are given by the eigenspace
 $\text{Ker}(H_N(v) - E_N(v))$, found by

$$E_N(v) = \inf_{\substack{\Psi \in H_a^1((\mathbb{R}^d)^N) \\ \int |\Psi|^2 = 1}} \langle \Psi, H_N(v) \Psi \rangle$$

Pure and mixed states

Define $P_\Psi := |\Psi\rangle\langle\Psi|$.

Pure and mixed states

Define $P_\Psi := |\Psi\rangle\langle\Psi|$.

- Pure states are

$$\left\{ P_\Psi, \Psi \in H_a^1(\mathbb{R}^{dN}), \int_{\mathbb{R}^{dN}} |\Psi|^2 = 1 \right\}$$

Pure and mixed states

Define $P_\Psi := |\Psi\rangle \langle \Psi|$.

- Pure states are

$$\left\{ P_\Psi, \Psi \in H_a^1(\mathbb{R}^{dN}), \int_{\mathbb{R}^{dN}} |\Psi|^2 = 1 \right\}$$

- Choose a basis $(\Psi_i)_{i \in \mathbb{N}}$. Mixed states are

$$\begin{aligned} \text{Conv } & \left\{ P_\Psi, \Psi \in H_a^1(\mathbb{R}^{dN}), \int_{\mathbb{R}^{dN}} |\Psi|^2 = 1 \right\} \\ &= \left\{ \sum_{i \in \mathbb{N}} \lambda_i P_{\Psi_i} \mid \sum_{i=1}^{+\infty} \lambda_i = 1, \lambda_i \geq 0 \right\} \\ &= \left\{ \Gamma \text{ op of } H_a^1(\mathbb{R}^{dN}) \mid \Gamma = \Gamma^* \geq 0, \text{Tr } \Gamma = 1 \right\} \end{aligned}$$

ground mixed states : $\text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v))$

The one-body density

- One-body density (much less information than Ψ)

$$\rho_{\Psi}(x) := N \int_{\mathbb{R}^{d(N-1)}} |\Psi|^2(x, x_2, \dots, x_N) dx_2 \cdots dx_N$$

The one-body density

- One-body density (much less information than Ψ)

$$\rho_{\Psi}(x) := N \int_{\mathbb{R}^{d(N-1)}} |\Psi|^2(x, x_2, \dots, x_N) dx_2 \cdots dx_N$$

- $\rho \geq 0$, $\int \rho_{\Psi} = N$, $\sqrt{\rho} \in H^1$

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Inverse potential

- Given $\rho \geq 0$, $\int \rho = N$, $k \in \mathbb{N}$, find v such that $\rho_{\Psi^{(k)}(v)} = \rho$.




Figure: Density ρ and its inverse v , for $N = 3$ and $k = 2$

Inverse potential

- Given $\rho \geq 0$, $\int \rho = N$, $k \in \mathbb{N}$, find v such that $\rho_{\Psi(v)} = \rho$.




Figure: Density ρ and its inverse v , for $N = 3$ and $k = 2$

Inverse potential

- Given $\rho \geq 0$, $\int \rho = N$, $k \in \mathbb{N}$, find v such that $\rho_{\Psi(v)} = \rho$.




Figure: Density ρ and its inverse v , for $N = 3$ and $k = 2$

Existence/uniqueness ?

Why finding inverse potentials ?

- Finding effective models in DFT

Why finding inverse potentials ?


- Finding effective models in DFT

Why finding inverse potentials ?

- Finding effective models in DFT
- Get rid of the complexity of interactions


Why finding inverse potentials ?

- Finding effective models in DFT
- Get rid of the complexity of interactions




Why finding inverse potentials ?

- Finding effective models in DFT
- Get rid of the complexity of interactions



Why finding inverse potentials ?

- Finding effective models in DFT
- Get rid of the complexity of interactions



Questions

DFT map: $\rho : v \mapsto \rho_{\Psi(v)} = \rho^{\text{HK}}(v)$

Given $\rho \geq 0$, $\int \rho = N$, we search v_ρ such that

$$\boxed{\rho^{\text{HK}}(v_\rho) = \rho}$$

Questions

DFT map: $\rho : v \mapsto \rho_{\Psi(v)} = \rho^{\text{HK}}(v)$

Given $\rho \geq 0$, $\int \rho = N$, we search v_ρ such that

$$\boxed{\rho^{\text{HK}}(v_\rho) = \rho}$$

- Is the DFT map injective ?

Questions

DFT map: $\rho : v \mapsto \rho_{\Psi(v)} = \rho^{\text{HK}}(v)$

Given $\rho \geq 0$, $\int \rho = N$, we search v_ρ such that

$$\boxed{\rho^{\text{HK}}(v_\rho) = \rho}$$

- Is the DFT map injective ?
- What is the image ? Is it dense ?

Questions

DFT map: $\rho : v \mapsto \rho_{\Psi(v)} = \rho^{\text{HK}}(v)$

Given $\rho \geq 0$, $\int \rho = N$, we search v_ρ such that

$$\boxed{\rho^{\text{HK}}(v_\rho) = \rho}$$

- Is the DFT map injective ?
- What is the image ? Is it dense ?
- Is the inverse problem well-posed ?

Questions

DFT map: $\rho : v \mapsto \rho_{\Psi(v)} = \rho^{\text{HK}}(v)$

Given $\rho \geq 0$, $\int \rho = N$, we search v_ρ such that

$$\boxed{\rho^{\text{HK}}(v_\rho) = \rho}$$

- Is the DFT map injective ?
- What is the image ? Is it dense ?
- Is the inverse problem well-posed ?
- How to invert it algorithmically ?

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

ρ contains everything

Theorem (Hohenberg-Kohn, 1964)

Let $w, v_1, v_2 \in L^{p > \max(2, 2d/3)}(\mathbb{R}^d) + L^\infty(\mathbb{R}^d)$. If there are two ground states Ψ_1 and Ψ_2 of $H_N(v_1)$ and $H_N(v_2)$, such that

$$\rho_{\Psi_1} = \rho_{\Psi_2},$$

then $v_1 = v_2 + \frac{E_1 - E_2}{N}$.

ρ contains everything

Theorem (Hohenberg-Kohn, 1964)

Let $w, v_1, v_2 \in L^{p > \max(2, 2d/3)}(\mathbb{R}^d) + L^\infty(\mathbb{R}^d)$. If there are two ground states Ψ_1 and Ψ_2 of $H_N(v_1)$ and $H_N(v_2)$, such that

$$\rho_{\Psi_1} = \rho_{\Psi_2},$$

then $v_1 = v_2 + \frac{E_1 - E_2}{N}$.

- Implies existence of functionals for any quantity

ρ contains everything

Theorem (Hohenberg-Kohn, 1964)

Let $w, v_1, v_2 \in L^{p > \max(2, 2d/3)}(\mathbb{R}^d) + L^\infty(\mathbb{R}^d)$. If there are two ground states Ψ_1 and Ψ_2 of $H_N(v_1)$ and $H_N(v_2)$, such that

$$\rho_{\Psi_1} = \rho_{\Psi_2},$$

then $v_1 = v_2 + \frac{E_1 - E_2}{N}$.

- Implies existence of functionals for any quantity
- Lieb (1964) remarked it relies on SUCP. Conjectures $L^{\frac{3}{2}}(\mathbb{R}^3)$, other mathematicians interested

Strong UCP

Theorem (Strong UCP for many-body Schrödinger operators)

Assume that the potentials satisfy $v, w \in L^p_{\text{loc}}(\mathbb{R}^d)$ with $p > \max(\frac{2d}{3}, 2)$. If $\Psi \in H^2_{\text{loc}}(\mathbb{R}^{dN})$ is a non zero solution to $H_N(v)\Psi = E\Psi$, then $|\{\Psi(X) = 0\}| = 0$.

Strong UCP

Theorem (Strong UCP for many-body Schrödinger operators)

Assume that the potentials satisfy $v, w \in L^p_{\text{loc}}(\mathbb{R}^d)$ with $p > \max(\frac{2d}{3}, 2)$. If $\Psi \in H^2_{\text{loc}}(\mathbb{R}^{dN})$ is a non zero solution to $H_N(v)\Psi = E\Psi$, then $|\{\Psi(X) = 0\}| = 0$.

- For $N = 1 : L^{\frac{3}{2}}$ (Jerison Kenig 1985)

Strong UCP

Theorem (Strong UCP for many-body Schrödinger operators)

Assume that the potentials satisfy $v, w \in L^p_{\text{loc}}(\mathbb{R}^d)$ with $p > \max(\frac{2d}{3}, 2)$. If $\Psi \in H^2_{\text{loc}}(\mathbb{R}^{dN})$ is a non zero solution to $H_N(v)\Psi = E\Psi$, then $|\{\Psi(X) = 0\}| = 0$.

- For $N = 1 : L^{\frac{3}{2}}$ (Jerison Kenig 1985)
- For $N \in \mathbb{N}$: already known that $\{\Psi(X) = 0\}$ is not an open set (Georgescu 1980) and (Schechter-Simon 1980). L^∞ with Jerison-Kenig

Strong UCP

Theorem (Strong UCP for many-body Schrödinger operators)

Assume that the potentials satisfy $v, w \in L^p_{\text{loc}}(\mathbb{R}^d)$ with $p > \max(\frac{2d}{3}, 2)$. If $\Psi \in H^2_{\text{loc}}(\mathbb{R}^{dN})$ is a non zero solution to $H_N(v)\Psi = E\Psi$, then $|\{\Psi(X) = 0\}| = 0$.

- For $N = 1 : L^{\frac{3}{2}}$ (Jerison Kenig 1985)
- For $N \in \mathbb{N}$: already known that $\{\Psi(X) = 0\}$ is not an open set (Georgescu 1980) and (Schechter-Simon 1980). L^∞ with Jerison-Kenig
- Lammert (2018) ; Laestadius, Benedicks, Penz (2020)

Strong UCP

Theorem (Strong UCP for many-body Schrödinger operators)

Assume that the potentials satisfy $v, w \in L^p_{\text{loc}}(\mathbb{R}^d)$ with $p > \max(\frac{2d}{3}, 2)$. If $\Psi \in H^2_{\text{loc}}(\mathbb{R}^{dN})$ is a non zero solution to $H_N(v)\Psi = E\Psi$, then $|\{\Psi(X) = 0\}| = 0$.

- For $N = 1 : L^{\frac{3}{2}}$ (Jerison Kenig 1985)
- For $N \in \mathbb{N}$: already known that $\{\Psi(X) = 0\}$ is not an open set (Georgescu 1980) and (Schechter-Simon 1980). L^∞ with Jerison-Kenig
- Lammert (2018) ; Laestadius, Benedicks, Penz (2020)
- This L^p result uses technics developped by Carleman, Hörmander, Koch and Tataru

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Compactness of $v \mapsto \rho^{\text{HK}}(v)$

Theorem (Main properties of Ψ)

- $v \mapsto \Psi(v)$ is \mathcal{C}^∞ from \mathcal{V}_N to H_p^1

Compactness of $v \mapsto \rho^{\text{HK}}(v)$

Theorem (Main properties of Ψ)

- $v \mapsto \Psi(v)$ is \mathcal{C}^∞ from \mathcal{V}_N to H_p^1
- For $v \in \mathcal{V}_N$, $d_v \Psi : L^{d/2} + L^\infty \rightarrow H^1 \cap \{\Psi(v)\}^\perp$

$$(d_v \Psi) u = -(H_N(v) - E_N(v))_\perp^{-1} (\sum_{i=1}^N u(x_i)) \Psi(v),$$

$d_v \Psi$ is compact

Compactness of $v \mapsto \rho^{\text{HK}}(v)$

Theorem (Main properties of Ψ)

- $v \mapsto \Psi(v)$ is \mathcal{C}^∞ from \mathcal{V}_N to H_p^1
- For $v \in \mathcal{V}_N$, $d_v \Psi : L^{d/2} + L^\infty \rightarrow H^1 \cap \{\Psi(v)\}^\perp$

$$(d_v \Psi) u = -(H_N(v) - E_N(v))_\perp^{-1} (\sum_{i=1}^N u(x_i)) \Psi(v),$$

$d_v \Psi$ is compact

- Let $\Lambda \subset \mathbb{R}^d$ be a bounded open set. Assume $v \in \mathcal{V}_N$, $v_n \rightharpoonup v$ and $v_n \mathbb{1}_{\mathbb{R}^d \setminus \Lambda} \rightarrow v \mathbb{1}_{\mathbb{R}^d \setminus \Lambda}$ in $L^{p+\epsilon} + L^\infty$. Then $E_N(v_n) \rightarrow E_N(v)$, $v_n \in \mathcal{V}_N$ for n large enough, and $\Psi(v_n) \rightarrow \Psi(v)$ in H^1

Ill-posedness of the inversion

Theorem (The set of ν -representable densities is very small)

Consider that the system lives in a bounded open set $\Omega \subset \mathbb{R}^d$.

Then $L^{p>d/2} \ni \nu \mapsto \rho^{HK}(\nu) \in W^{1,1}$ is weak-strong continuous, $(\rho^{HK})^{-1}$ is discontinuous, and $\rho^{HK}(L^p(\mathbb{R}^d))$ has empty interior in $W^{1,1} \cap \{\int \cdot = N\}$.

Ill-posedness of the inversion

Theorem (The set of ν -representable densities is very small)

Consider that the system lives in a bounded open set $\Omega \subset \mathbb{R}^d$.

Then $L^{p>d/2} \ni \nu \mapsto \rho^{HK}(\nu) \in W^{1,1}$ is weak-strong continuous, $(\rho^{HK})^{-1}$ is discontinuous, and $\rho^{HK}(L^p(\mathbb{R}^d))$ has empty interior in $W^{1,1} \cap \{\int \cdot = N\}$.

The inverse problem is ill-posed !

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 **The dual problem**
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 **The dual problem**
 - **Literature**
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} & \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ & \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} & \{\rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1\} \\ & \subset \{\rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1\} \end{aligned}$$

Inverse problem solved for

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Inverse problem solved for

- **approximate invertibility with mixed ground states** (Lieb 1983)

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Inverse problem solved for

- approximate invertibility with mixed ground states (Lieb 1983)
- classical systems at $T > 0$ (Chayes Chayes Lieb 1984)

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} & \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ & \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Inverse problem solved for

- approximate invertibility with mixed ground states (Lieb 1983)
- classical systems at $T > 0$ (Chayes Chayes Lieb 1984)
- quantum systems on lattices for mixed ground states (Chayes Chayes Ruskai 1985)

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Inverse problem solved for

- approximate invertibility with mixed ground states (Lieb 1983)
- classical systems at $T > 0$ (Chayes Chayes Lieb 1984)
- quantum systems on lattices for mixed ground states (Chayes Chayes Ruskai 1985)
- Moreau-Yosida regularization (Kvaal, Ekström, Teale, Helgaker, Tellgren, Ruggenthaler, Penz, Lammert, Herbst)

Existing literature

Target ρ : we search v such that

- $\rho_{\Psi(v)} = \rho$ for pure states, $\Psi(v) \in \text{Ker}(H_N(v) - E_N(v))$
- $\rho_{\Gamma(v)} = \rho$ for mixed states, $\text{Ran } \Gamma(v) \subset \text{Ker}(H_N(v) - E_N(v))$

$$\begin{aligned} \{ \rho_{\Psi} \mid \Psi \in \text{Ker}(H_N(v) - E_N(v)), \|\Psi\|_{L^2} = 1 \} \\ \subset \{ \rho_{\Gamma} \mid \text{Ran } \Gamma \subset \text{Ker}(H_N(v) - E_N(v)), \text{Tr } \Gamma = 1 \} \end{aligned}$$

Inverse problem solved for

- approximate invertibility with mixed ground states (Lieb 1983)
- classical systems at $T > 0$ (Chayes Chayes Lieb 1984)
- quantum systems on lattices for mixed ground states (Chayes Chayes Ruskai 1985)
- Moreau-Yosida regularization (Kvaal, Ekström, Teale, Helgaker, Tellgren, Ruggenthaler, Penz, Lammert, Herbst)
- Numerical articles

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 **The dual problem**
 - Literature
 - **Optimality properties**
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Dual optimality

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho, \quad \sup_{v \in L^p(\mathbb{R}^d)} G_\rho(v) < +\infty$$

Dual optimality

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho, \quad \sup_{v \in L^p(\mathbb{R}^d)} G_\rho(v) < +\infty$$

- Gauge invariance $G_\rho(v + c) = G_\rho(v)$

Dual optimality

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho, \quad \sup_{v \in L^p(\mathbb{R}^d)} G_\rho(v) < +\infty$$

- Gauge invariance $G_\rho(v + c) = G_\rho(v)$
- Concave for $k = 0$

Dual optimality

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho, \quad \sup_{v \in L^p(\mathbb{R}^d)} G_\rho(v) < +\infty$$

- Gauge invariance $G_\rho(v + c) = G_\rho(v)$
- Concave for $k = 0$
- On degenerate potentials, $v \mapsto \rho_{\Psi(v)}$ and E_N are **not differentiable**

Dual optimality

$$G_{\rho}^{(k)}(v) := E_N^{(k)}(v) - \int_{\mathbb{R}^d} v \rho$$

Theorem (Optimality in the dual problem)

Dual optimality

$$G_{\rho}^{(k)}(v) := E_N^{(k)}(v) - \int_{\mathbb{R}^d} v \rho$$

Theorem (Optimality in the dual problem)

Take $w \geq 0$, $\rho \in L^1(\mathbb{R}^d)$, $\rho \geq 0$, $\int \rho = N$, $\sqrt{\rho} \in H^1(\mathbb{R}^d)$, $v \in L^{p>2}$ such that $H_N(v)$ has a ground state.

i) The following statements are equivalent

- there is a k^{th} bound mixed state Γ of v such that $\rho_{\Gamma} = \rho$
- v is a **local maximizer** of $G_{\rho}^{(k)}$
- v is a **global maximizer** of $G_{\rho}^{(k)}$

Dual optimality

$$G_{\rho}^{(k)}(v) := E_N^{(k)}(v) - \int_{\mathbb{R}^d} v \rho$$

Theorem (Optimality in the dual problem)

Take $w \geq 0$, $\rho \in L^1(\mathbb{R}^d)$, $\rho \geq 0$, $\int \rho = N$, $\sqrt{\rho} \in H^1(\mathbb{R}^d)$, $v \in L^{p>2}$ such that $H_N(v)$ has a ground state.

i) The following statements are equivalent

- there is a k^{th} bound mixed state Γ of v such that $\rho_{\Gamma} = \rho$
- v is a **local maximizer** of $G_{\rho}^{(k)}$
- v is a global maximizer of $G_{\rho}^{(k)}$

ii) If v maximizes $G_{\rho}^{(k)}$ and

- $\dim \text{Ker} (H_N(v) - E_N(v)) \in \{1, 2\}$,
- or $d = 1$ and $w = 0$,

then v has a k^{th} bound pure state Ψ such that $\rho_{\Psi} = \rho$.

Dual optimality

$$G_\rho^{(k)}(v) := E_N^{(k)}(v) - \int_{\mathbb{R}^d} v \rho$$

Theorem (Optimality in the dual problem)

Take $w \geq 0$, $\rho \in L^1(\mathbb{R}^d)$, $\rho \geq 0$, $\int \rho = N$, $\sqrt{\rho} \in H^1(\mathbb{R}^d)$, $v \in L^{p>2}$ such that $H_N(v)$ has a ground state.

i) The following statements are equivalent

- there is a k^{th} bound mixed state Γ of v such that $\rho_\Gamma = \rho$
- v is a **local maximizer** of $G_\rho^{(k)}$
- v is a global maximizer of $G_\rho^{(k)}$

ii) If v maximizes $G_\rho^{(k)}$ and

- $\dim \text{Ker} (H_N(v) - E_N(v)) \in \{1, 2\}$,
- or $d = 1$ and $w = 0$,

then v has a k^{th} bound pure state Ψ such that $\rho_\Psi = \rho$.

Does a maximum exist ?

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 **The dual problem**
 - Literature
 - Optimality properties
 - **Regularization**
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn


Regularization

- $G_\rho(v) = E_N(v) - \int v\rho$ is **not coercive** in L^p ! Ex :
 $v \in L^1 \cap L^{p>1}$, $v \geq 0$, $v_n(x) := n^d v(nx)$,
 $\|v_n\|_{L^p}^p = n^{d(p-1)} \int v^p \rightarrow +\infty$ but $E_N(v_n) = 0$, and
 $\int v_n \rho \rightarrow \rho(0) \int v$ is bounded

Regularization

- $G_\rho(v) = E_N(v) - \int v\rho$ is **not coercive** in L^p ! Ex :
 $v \in L^1 \cap L^{p>1}$, $v \geq 0$, $v_n(x) := n^d v(nx)$,
 $\|v_n\|_{L^p}^p = n^{d(p-1)} \int v^p \rightarrow +\infty$ but $E_N(v_n) = 0$, and
 $\int v_n \rho \rightarrow \rho(0) \int v$ is bounded
- Dual : restriction to potentials $V = \sum_{i \in I} v_i \alpha_i$,
 $v \in (v_i)_{i \in I} \in \ell^\infty(I, \mathbb{R})$, $\alpha_i \in L^\infty(\Omega)$, $\sum_{i \in I} \alpha_i = \mathbf{1}_\Omega$, $r_i \in \mathbb{R}_+$,
 $r_i = \int \rho \alpha_i$, $\sum_{i \in I} r_i = N$

$$G_{r,\alpha}^{(k)}(v) := E_N \left(\sum_{i \in I} v_i \alpha_i \right) - \sum_{i \in I} v_i r_i,$$



Coercivity

$$G_{r,\alpha}^{(k)}(v) \leq -\min \left(1, \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \right) \|v - c_\Omega\|_{\ell_r^1} + c_R,$$

Coercivity

$$G_{r,\alpha}^{(k)}(v) \leq -\min \left(1, \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \right) \|v - c_\Omega\|_{\ell_r^1} + c_R,$$

Theorem (Existence of the inverse potential)

When I is finite $G_{r,\alpha}^{(k)}$ is coercive and there exists a maximizer v . If $\Omega \subset \mathbb{R}^d$ is bounded, there is a k^{th} excited N -particle ground mixed state Γ_v of H_N ($\sum_{i \in I} v_i \alpha_i$) such that $\int \alpha_i \rho_{\Gamma_v} = r_i$ ($= \int \alpha_i \rho$) $\forall i$.

Coercivity

$$G_{r,\alpha}^{(k)}(v) \leq -\min \left(1, \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \right) \|v - c_\Omega\|_{\ell_r^1} + c_R,$$

Theorem (Existence of the inverse potential)

When I is finite $G_{r,\alpha}^{(k)}$ is coercive and there exists a maximizer v . If $\Omega \subset \mathbb{R}^d$ is bounded, there is a k^{th} excited N -particle ground mixed state Γ_v of $H_N(\sum_{i \in I} v_i \alpha_i)$ such that $\int \alpha_i \rho_{\Gamma_v} = r_i (= \int \alpha_i \rho) \forall i$.

- Constructive inversion with mixed states**

For a given ρ , $\epsilon > 0$, there exists a potential v and Γ_v with $\text{Ran } \Gamma_v \subset \text{Ker}(H_N(v) - E_N(v))$ such that $\|\rho_{\Gamma_v} - \rho\|_{L^1 \cap L^q} \leq \epsilon$.

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

“Gradient” ascent

Minimize $J(v) := \int_{\mathbb{R}^d} (\rho_{\Psi(v)} - \rho)^2$?

“Gradient” ascent

Minimize $J(v) := \int_{\mathbb{R}^d} (\rho_{\Psi(v)} - \rho)^2$?

Second idea, maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

Local dual problem

$${}^+\delta_v G_\rho^{(k)}(u) = \max_{\substack{\Psi_0, \dots, \Psi_{M_k-k} \in \text{Ker} \left(H_N(v) - E_N(v) \right) \\ \|\Psi_i\|=1, \Psi_i \perp \Psi_j \\ 0 \leq i, j \leq M_k - k}} \min_{\substack{\Psi = \sum_{i=0}^{M_k-k} \lambda_i \Psi_i \\ \lambda_i \in \mathbb{C}, \sum_i |\lambda_i|^2 = 1}} \int (\rho_\Psi - \rho) u$$

Local dual problem

$${}^+\delta_v G_\rho^{(k)}(u) = \max_{\substack{\Psi_0, \dots, \Psi_{M_k-k} \in \text{Ker}(H_N(v) - E_N(v)) \\ \|\Psi_i\|=1, \Psi_i \perp \Psi_j \\ 0 \leq i, j \leq M_k-k}} \min_{\substack{\Psi = \sum_{i=0}^{M_k-k} \lambda_i \Psi_i \\ \lambda_i \in \mathbb{C}, \sum_i |\lambda_i|^2 = 1}} \int (\rho_\Psi - \rho) u$$

Proposition (Local dual problem)

Take $w \geq 0$, $v \in \mathcal{V}_{N,\partial}$. We have

$$\sup_{\substack{u \in L^p + L^\infty \\ \|u\|_{L^p + L^\infty} = 1}} {}^+\delta_v G_\rho(u) = \max_{\substack{Q \subset \text{Ker}_{\mathbb{R}}(H_N(v) - E_N(v)) \\ \dim_{\mathbb{R}} Q = M_k - k + 1}} \min_{\substack{\Gamma \in \mathcal{S}(Q) \\ \Gamma \geq 0, \text{Tr } \Gamma = 1}} \|\rho_\Gamma - \rho\|_{L^{p'}},$$

and the supremum is attained by $u^* = \left| \frac{\rho_{\Gamma^*} - \rho}{\|\rho_{\Gamma^*} - \rho\|_{L^{p'}}} \right|^{p'-1} \text{sgn}(\rho_{\Gamma^*} - \rho)$,
where Γ^* is an optimizer of the right hand side.

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization
- Consider a target $\rho \geq 0$, $\int \rho = N$

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization
- Consider a target $\rho \geq 0$, $\int \rho = N$
- Start from Bohm's potential $\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}$ or Thomas-Fermi $c_{\text{TF}} \rho^{\frac{2}{3}}$

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization
- Consider a target $\rho \geq 0$, $\int \rho = N$
- Start from Bohm's potential $\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}$ or Thomas-Fermi $c_{\text{TF}} \rho^{\frac{2}{3}}$
- Iterate $v_{n+1} = v_n + \alpha u^*$, $+\delta_v G_\rho(u^*) = \max_{\|u\|=1} +\delta_v G_\rho(u) > 0$

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization
- Consider a target $\rho \geq 0$, $\int \rho = N$
- Start from Bohm's potential $\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}$ or Thomas-Fermi $c_{\text{TF}} \rho^{\frac{2}{3}}$
- Iterate $v_{n+1} = v_n + \alpha u^*$, $+\delta_v G_\rho(u^*) = \max_{\|u\|=1} +\delta_v G_\rho(u) > 0$
- Line search for α , temperature

“Gradient” ascent

Maximize

$$G_\rho(v) := E_N(v) - \int_{\mathbb{R}^d} v \rho$$

- Fourier discretization
- Consider a target $\rho \geq 0$, $\int \rho = N$
- Start from Bohm's potential $\frac{\Delta \sqrt{\rho}}{\sqrt{\rho}}$ or Thomas-Fermi $c_{\text{TF}} \rho^{\frac{2}{3}}$
- Iterate $v_{n+1} = v_n + \alpha u^*$, $+\delta_v G_\rho(u^*) = \max_{\|u\|=1} +\delta_v G_\rho(u) > 0$
- Line search for α , temperature
- Convergence criterion: $\|\rho^{\text{HK}}(v_n) - \rho\|_{L^1} / N \leq \epsilon$

Goal

What we know

- Approximate inversion with mixed states for any k

Goal

What we know

- Approximate inversion with mixed states for any k
- When $d = 1$, the set of pure state densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is dense for the $L^1 \cap L^q$ norm

Goal

What we know

- Approximate inversion with mixed states for any k
- When $d = 1$, the set of pure state densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is dense for the $L^1 \cap L^q$ norm

- When $d = 3$, it's **not** (uses Lieb 83)

Goal

What we know

- Approximate inversion with mixed states for any k
- When $d = 1$, the set of pure state densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is dense for the $L^1 \cap L^q$ norm

- When $d = 3$, it's **not** (uses Lieb 83)

What we want to know

Goal

What we know

- Approximate inversion with mixed states for any k
- When $d = 1$, the set of pure state densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is dense for the $L^1 \cap L^q$ norm

- When $d = 3$, it's **not** (uses Lieb 83)

What we want to know

- Uniqueness for $k \geq 1$?

Goal

What we know

- Approximate inversion with mixed states for any k
- When $d = 1$, the set of pure state densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is dense for the $L^1 \cap L^q$ norm

- When $d = 3$, it's **not** (uses Lieb 83)

What we want to know

- Uniqueness for $k \geq 1$?
- Inversion with pure states for $d = 2$?

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - **Graphs**
 - Proof of coercivity
 - What we learn

$d = 1$




Figure: Plot for $d = 1$, $N = 5$, $k = 0$ on the left, $k = 3$ on the right,
 $\log_{10} |\rho_n - \rho|$, $\log_{10} |v_n - v|$

Uniqueness




Figure: $d = 1$, $N = 3$, $k = 0$ left, $k = 1$ middle, $k = 5$ right. Densities in blue, inverse potentials in other colors

$d = 2$




Figure: $d = 2$, $N = 5$, $k = 0$; v , $\rho_{\Psi^{(0)}(v)}$, $\log_{10} |v_n - v|$,
 $\log_{10} |\rho_n - \rho_{\Psi^{(0)}(v)}|$

$d = 3$




Figure: $d = 3$, $N = 4$, $k = 1$; ρ , v_n , $\log_{10} |\rho_n - \rho|$

Simulations at high densities




Figure: Convergence of $\rho_N^{-1}(N\rho)/N^{\frac{2}{d}}$, $\int \rho = 1$

Simulations at high densities




Figure: Convergence of $\rho_N^{-1}(N\rho)/N^{\frac{2}{d}}$, $\int \rho = 1$

Conjecture

For any $\rho \geq 0$ such that $\int \rho = 1$ and $\sqrt{\rho} \in H^1$,

$$\frac{\rho_N^{-1}(N\rho)}{N^{\frac{2}{d}}} \xrightarrow{N \rightarrow +\infty}$$

Conjecture

For any $\rho \geq 0$ such that $\int \rho = 1$ and $\sqrt{\rho} \in H^1$,

$$\frac{\rho_N^{-1}(N\rho)}{N^{\frac{2}{d}}} \xrightarrow{N \rightarrow +\infty} v_{\text{TF},\rho} = -\rho^{\frac{2}{d}}$$

The direct statement version is in Founais, Lewin, Solovej (2019)

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

Statement

Theorem

Take $w \in (L^p + L^\infty)(\mathbb{R}^d, \mathbb{R}_+)$. Let α be a partition of unity of Ω , with $\alpha_i \in L^\infty(\Omega, \mathbb{R}_+)$, such that we have $R > 0$ for which

$$(\text{supp } \alpha_i) \setminus \bigcup_{j \in I, j \neq i} \text{supp } \alpha_j$$

contains a ball of radius R , uniformly in $i \in I$. Let $r \in \ell^1(I, \mathbb{R}_+)$ be such that $\sum_{i \in I} r_i = N$ and $r_i > 0$ for all $i \in I$. For any $v \in \ell_r^1(I, \mathbb{R})$ such that $E_N(\sum_{i \in I} v_i \alpha_i) = 0$, we have

$$G_{r, \alpha}^{(k)}(v) \leq - \min \left(1, \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \right) \|v - c_\Omega\|_{\ell_r^1} + c_R$$

Proof

- We assumed that there are points $y_i \in \mathbb{R}^d$ such that for any $i \in I$,

$$B_R(y_i) \subset (\text{supp } \alpha_i) \setminus \cup_{j \in I, j \neq i} \text{supp } \alpha_j.$$

We write $X = (x_1, \dots, x_N)$ and $Y_i := (y_i, \dots, y_i)$. Take normalized $\Phi_0, \dots, \Phi_k \in \wedge^N H_0^1(B_R)$ with disjoint supports. Take some non-empty $Q \subset I$ and for $j \in \{0, \dots, k\}$, form

$$\psi_{j,Q}(X) := \frac{1}{\sqrt{\sum_{i \in Q} r_i}} \sum_{i \in Q} \sqrt{r_i} \Phi_j(X - Y_i).$$

This satisfies $\int_{\Omega^N} |\psi_{j,Q}|^2 = 1$, $T(\Psi_{j,Q}) = T(\Phi_j)$,
 $W(\Psi_{j,Q}) = W(\Phi_j)$ and

$$\rho_{\Psi_{j,Q}}(x) = \left(\sum_{i \in Q} r_i \right)^{-1} \sum_{i \in Q} r_i \rho_{\Phi_j}(x - y_i).$$

We use the expression

$$E_N^{(k)}(V) = \inf_{\dim A = k+1} \max_{\substack{\Psi \in A \\ \int_{\Omega^N} |\Psi|^2 = 1}} \mathcal{E}_V(\Psi)$$

and choose the frame $A := (\Psi_{0,Q}, \dots, \Psi_{k,Q})$ so that

$$G_{r,\alpha}^{(k)}(v) \leq - \sum_{i \in I} v_i r_i + \max_{\substack{\lambda_j \in \mathbb{C} \\ \sum_{j=0}^k |\lambda_j|^2 = 1}} \mathcal{E}_{V(v)} \left(\sum_{j=0}^k \lambda_j \Psi_{j,Q} \right).$$

For any $i \in I$, the only non-vanishing element of α in $B_R(y_i)$ is α_i , so $\alpha_i = 1$ on $B_R(y_i)$ and

$$\int_{\Omega} \alpha_i \rho \psi_{j,Q} = \frac{Nr_i \delta_{i \in Q}}{\sum_{\ell \in Q} r_{\ell}},$$

$$\int_{\Omega} V(v) \rho_{\sum_{j=0}^k \lambda_j \psi_{j,Q}} = \sum_{j=0}^k |\lambda_j|^2 \int_{\Omega} V(v) \rho \psi_{j,Q} = \frac{N \sum_{i \in Q} v_i r_i}{\sum_{\ell \in Q} r_{\ell}}.$$

We see that the external potential energy of the trial state does not depend on the λ_j 's. Defining $c_R := \max_{\substack{\lambda_j \in \mathbb{C} \\ \sum_{j=0}^k |\lambda_j|^2 = 1}} \mathcal{E}_0 \left(\sum_{j=0}^k \lambda_j \psi_{j,Q} \right)$,

we deduce that

$$\begin{aligned} G_{r,\alpha}(v) &\leq c_R + \frac{N}{\sum_{i \in Q} r_i} \sum_{i \in Q} v_i r_i - \sum_{i \in I} v_i r_i \\ &= c_R + \frac{\sum_{i \in I \setminus Q} r_i}{\sum_{i \in Q} r_i} \sum_{i \in Q} v_i r_i - \sum_{i \in I \setminus Q} v_i r_i. \end{aligned}$$

Since G is gauge invariant, for any $\mu \in \mathbb{R}$ and any non-empty $Q \subset I$, we have

$$G_{r,\alpha}(v) = G_{r,\alpha}(v - \mu) \leq c_R + \frac{\sum_{I \setminus Q} r_i}{\sum_{i \in Q} r_i} \sum_{i \in Q} (v_i - \mu) r_i - \sum_{i \in I \setminus Q} (v_i - \mu) r_i.$$

We define the two sets $I_v^\pm := \{i \in I \mid \pm v_i > \pm c_\Omega\}$. In the case $I_v^- \neq \emptyset$, we take $Q = I_v^-$ and $\mu = c_\Omega$ yielding

$$\begin{aligned} G_{r,\alpha}(v) - c_R &\leq \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \sum_{v_i < c_\Omega} (v_i - c_\Omega) r_i - \sum_{v_i \geq c_\Omega} (v_i - c_\Omega) r_i \\ &\leq \min \left(1, \frac{\sum_{v_i \geq c_\Omega} r_i}{\sum_{v_i < c_\Omega} r_i} \right) \left(\sum_{v_i < c_\Omega} (v_i - c_\Omega) r_i - \sum_{v_i \geq c_\Omega} (v_i - c_\Omega) r_i \right) \\ &\leq - \frac{\sum_{v_i \geq c_\Omega} r_i}{N} \|v - c_\Omega\|_{\ell_r^1}. \end{aligned}$$

Table of contents

- 1 Introduction
 - The setting
 - The goal
- 2 The direct problem
 - Uniqueness
 - Ill-posedness
- 3 The dual problem
 - Literature
 - Optimality properties
 - Regularization
- 4 Numerical inversion
 - The local problem
 - Graphs
 - Proof of coercivity
 - What we learn

What we learn from simulations

- Confirms Gaudoin and Burke (2004), no uniqueness for $k \geq 1$

What we learn from simulations

- Confirms Gaudoin and Burke (2004), no uniqueness for $k \geq 1$
- For $d = 2$, the set of pure states densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is **dense** in the set of positive functions

What we learn from simulations

- Confirms Gaudoin and Burke (2004), no uniqueness for $k \geq 1$
- For $d = 2$, the set of pure states densities

$$\left\{ \rho_{\Psi_v} \mid v \in (L^p + L^\infty)(\Omega), \right. \\ \left. \Psi_v \in \text{Ker} (H_N^{w=0}(v) - E_N(v)), \int_{\Omega^N} |\Psi_v|^2 = 1 \right\}$$

is **dense** in the set of positive functions

- Degeneracies are generic, even for $d = 1$. Need to be considered, not in literature

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**
- Extension for current DFT

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**
- Extension for current DFT
- Coercivity proof applicable to Moreau-Yosida ?

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**
- Extension for current DFT
- Coercivity proof applicable to Moreau-Yosida ?
- Compare the different methods

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**
- Extension for current DFT
- Coercivity proof applicable to Moreau-Yosida ?
- Compare the different methods
- Accelerate the convergence, work with functionals developers

Conclusion

Conclusions

- No uniqueness for $k \geq 1$ (simulations)
- Constructive mixed states inversion: for any ρ, k, d, ϵ , we can find v such that $\|\rho_{\Gamma(v)} - \rho\|_{L^1 \cap L^q} \leq \epsilon$
- Pure states inversion:
 - $d = 1$ **yes** (theoretical)
 - $d = 2$ **yes** (simulations)
 - $d = 3$ **no** (theoretical but not rigorous)
- We gave an algorithm

Next questions

- **Exact v -representability**
- Extension for current DFT
- Coercivity proof applicable to Moreau-Yosida ?
- Compare the different methods
- Accelerate the convergence, work with functionals developers
- Symmetry study