

Regularization and density-potential inversion in density-functional theory

A. Laestadius


Mathematical Modelling group, Department of Computer Science, OsloMet

MPI Hamburg seminar, November 2022

Acknowledgements

OSLOMET


OSLO METROPOLITAN UNIVERSITY
STORBYUNIVERSITETET



Mathematical Modeling group OsloMet
Quantum Hub



European Research Council
Established by the European Commission



- Work part of ERC StG *REGAL–Regularized Density-Functional Analysis*
- Collaboration: M. Penz (Austria/Germany) and M.A. Csirik (Norway/Hungary)

- N interacting electrons:

$$\begin{aligned} T &= \text{kinetic energy}, & W &= \text{two-particle interaction} \\ H &= \underbrace{T + W}_{=:H_0} + V, & V(r_1, \dots, r_N) &= \sum_{j=1}^N v(r_j) \\ E(v) &= \inf \{ \langle \psi, H\psi \rangle : \underbrace{\psi \in H^1(\mathbb{R}^{3N}), \|\psi\| = 1}_{=: \mathcal{W}_N} \}, \end{aligned}$$

- One-body (particle) density:

$$\rho_\psi(r) = N \int_{\mathbb{R}^{3(N-1)}} |\psi|^2, \quad \psi \in \mathcal{W}_N \implies \rho_\psi \in L^1 \cap L^3 \quad (1)$$

- Using $H = H_0 + \sum_j v(r_j)$ and $\rho_\psi = N \int |\psi|^2$

$$\langle \psi, H\psi \rangle = \underbrace{\langle \psi, H_0\psi \rangle}_{\text{universal}} + \int_{\mathbb{R}^3} v\rho_\psi$$

- Ground-state energy via “constrained-search”:

$$\begin{aligned} E(v) &= \inf_{\psi} \left\{ \langle \psi, H_0\psi \rangle + \int v\rho_\psi \mathrm{d}r \right\} \\ &= \inf_{\rho} \left\{ \underbrace{\inf \{ \langle \psi, H_0\psi \rangle : \rho_\psi = \rho \}}_{\text{universal } \rho\text{-functional}} + \int v\rho \mathrm{d}r \right\} \end{aligned}$$

Γ density matrix

$$\Gamma = \sum_j f_j |\psi_j\rangle\langle\psi_j|, \quad f_j \in [0, 1], \quad \sum_j f_j = N$$

$$\rho_\Gamma = \sum_j f_j \rho_{\psi_j}$$

- N -rep. density ρ : exists Γ such that $\rho = \rho_\Gamma$
- v -rep. density ρ : exists Γ such that $\rho = \rho_\Gamma$ and Γ is ground state
- In case Γ is a pure state, we may specify pure-state rep. (otherwise we use “ensemble”).

Ensemble v -rep density are dense on the set of N -rep densities

$$E(v) = \inf_{\rho} \left\{ \underbrace{\inf \{ \langle \psi, H_0 \psi \rangle : \rho_{\psi} = \rho \}}_{\text{Levy-Lieb functional}} + \int v \rho dr \right\}$$

Levy-Lieb functional

$$\tilde{F}(\rho) = \inf_{\psi: \rho_{\psi} = \rho} \{ \langle \psi, H_0 \psi \rangle \}$$

is defined for N -rep. ρ , not convex.

$$E(v) = \inf_{\rho} \left\{ \tilde{F}(\rho) + \int v \rho \right\}$$

Lieb functional

$$F(\rho) = \sup_v \left\{ E(v) - \int v \rho dr \right\}$$

is defined for N -rep. ρ , convex but **NOT** differentiable

Lieb's constrained-search functional

$$F_{\text{DM}}(\rho) = \inf_{\Gamma: \rho_{\Gamma} = \rho} \text{tr}(\Gamma H_0)$$

Convex and defined for N -rep. densities

Theorem (Lieb 1983)

$$F_{\text{DM}}(\rho) = F(\rho) \text{ (convex and lsc)}$$

Moreau–Yosida Regularization


Hohenberg–Kohn variational principle:

$$E(v) = \min_{\rho} \left\{ F(\rho) + \int v \rho \, dr \right\} \quad F(\rho) \text{ is highly irregular}$$

Moreau–Yosida regularization:

$$E^{\varepsilon}(v) = E(v) - \frac{\varepsilon}{2} \|v\|^2$$

$$E^{\varepsilon}(v) = \min_{\rho} \left\{ F^{\varepsilon}(\rho) + \int v \rho \, dr \right\} \quad F^{\varepsilon}(\rho) \text{ is differentiable}$$



- Suppose $\rho \in X$, $v \in X^*$
- Let J be the duality mapping
- J maps elements from X to X^*
- Definition

$$J(\rho) = \{\xi \in X^* \mid \langle \xi, \rho \rangle = \|\rho\|_X^2 = \|\xi\|_{X^*}^2\}. \quad (2)$$

- Fixed choice of $\varepsilon > 0$ (reg. parameter)
- Moreau–Yosida regularization of a convex, lower semicontinuous functional $f : X \rightarrow \cup\{+\infty\}$
- Given by the lower envelope of the parabola $\rho \mapsto \frac{1}{2\varepsilon} \|\rho\|_X^2$, tracing along the function

$$f^\varepsilon(\rho) = \inf_{\rho' \in X} \left\{ f(\rho') + \frac{1}{2\varepsilon} \|\rho - \rho'\|_X^2 \right\}. \quad (3)$$

- Infimum is attained at a unique point,
proximal mapping $\Pi_f^\varepsilon : X \rightarrow X$ makes sense:

$$\rho^\varepsilon := \Pi_f^\varepsilon(\rho) = \operatorname{argmin}_{\rho' \in X} \left\{ f(\rho') + \frac{1}{2\varepsilon} \|\rho - \rho'\|_X^2 \right\}. \quad (4)$$

- Proximal mapping Π_f^ε is singleton-valued (in particular nonempty) everywhere

Moreover, $\Pi_f^\varepsilon(\rho) \rightarrow \rho$ as $\varepsilon \rightarrow 0$

- Derivative:

$$\nabla f^\varepsilon(\rho) = \frac{1}{\varepsilon} J(\rho - \Pi_f^\varepsilon(\rho))$$

- Subdifferential: For any $\rho, \rho_\varepsilon \in X$:

$$\rho^\varepsilon = \Pi_f^\varepsilon(\rho) \iff \nabla f^\varepsilon(\rho) \in \partial f(\rho_\varepsilon)$$

- Given ρ_{gs} from $H = T + W + V_{\text{ext}}$, i.e.,

$$(H + V_{\text{ext}})\psi = E(v_{\text{ext}})\psi, \quad \rho_\psi = \rho_{\text{gs}}$$

- Suppose that ρ_{gs} is both interacting and noninteracting v -rep., then by construction of the KS scheme,

$$\left(-\frac{1}{2}\nabla^2 + v_{\text{KS}}^{\rho_{\text{gs}}}\right)\varphi_j = e_j\varphi_j, \quad (T + V_{\text{KS}})\Phi = E_{\text{KS}}\Phi \quad (5)$$

and

$$\underbrace{\sum_j |\varphi_j|^2}_{\text{non-int}} = \rho_{\text{gs}} = N \underbrace{\int_{\mathbb{R}^{3(N-1)}} |\psi|^2}_{\text{int.}}$$

- In practice ρ_{gs} not known, only v_{ext}
- Define the Hartree potential

$$v_{\text{H}}^{\rho}(r) = (\rho \star |\cdot|^{-1})(r) = \int \frac{\rho(r')}{|r - r'|} dr',$$

- Assumption: representability


We can then write

$$v_{\text{KS}}^{\rho} = v_{\text{ext}} + v_{\text{H}}^{\rho} + v_{\text{xc}}^{\rho}$$

where v_{H}^{ρ} and v_{xc}^{ρ} are determined by the density ρ (the latter unknown)

$$\left(-\frac{1}{2} \nabla^2 + v_{\text{ext}} + v_{\text{H}}^{\rho} + v_{\text{xc}}^{\rho} \right) \varphi_j = e_j \varphi_j, \quad \rho = \sum_{j=1}^N |\varphi_j|^2 \quad (6)$$

Exchange-correlation = Nature's glue



How can we determine v_{XC} , i.e., the KS potential given ρ_{gs} ?

- The ZMP method uses the constraint

$$\frac{\lambda}{2} \int \int \frac{(\rho(r) - \rho_{\text{gs}}(r))(\rho(r') - \rho_{\text{gs}}(r'))}{|r - r'|} dr dr'$$

- Lagrange multiplier λ
- Given ρ_{gs} , the ZMP numerically allows to determine v_{xc} in the limit $\lambda \rightarrow \infty$ from KS-like eqs. (“ZMP eqs.”)

ρ_{gs} is a ground-state of $H = T + W + V_{\text{ext}}$

- $\rho^\lambda(r) = \sum_{j=1}^N |\varphi_j^\lambda(r)|^2,$

$$v^\lambda(r) = \lambda \int \frac{\rho^\lambda(r') - \rho_{\text{gs}}(r')}{|r - r'|} r', \quad v_{\text{H}}^\lambda = (\rho^\lambda \star |\cdot|^{-1}),$$

where φ_j^λ satisfy the ZMP equations

$$\left[-\frac{1}{2} \nabla^2 + v_{\text{ext}} + v_{\text{H}}^\lambda + v^\lambda \right] \varphi_j^\lambda = e_j^\lambda \varphi_j^\lambda, \quad (7)$$

- Then, formally,

$$v_{\text{xc}}(r) = \lim_{\lambda \rightarrow \infty} (\lambda + 1) \int \frac{\rho^\lambda(r') - \rho_{\text{gs}}(r')}{|r - r'|} dr'.$$

ZMP III (details: iteration at a given λ)

- The pair $(\rho^\lambda, v_{\text{xc}}^\lambda)$ is determined self-consistently

$$v_{\text{xc}}^\lambda(r) = (\lambda + 1) \lim_{i \rightarrow \infty} \int \frac{\rho_i^\lambda(r') - \rho_{\text{gs}}(r')}{|r - r'|} dr'$$

- ρ_i is obtained from the orbitals being solutions of (7) with

$$v_i^\lambda(r) = \lambda \int \frac{\rho_{i-1}^\lambda(r') - \rho_{\text{gs}}(r')}{|r - r'|} dr',$$

- Procedure is repeated for increasing values of λ : $\lambda \rightarrow \infty$ yields a potential $v_{\text{xc}}^\lambda \rightarrow v_{\text{xc}}$ that has the ground-state density ρ_{gs}

ZMP from MY regularization

- Choose space X, X^*
- Recall duality map $J : X \rightarrow X^*$
- Use “KS density functional”

$$f(\rho) := \mathcal{Q}_{\text{KS}}(\rho) = T(\rho) + \int v_{\text{ext}} \rho \, dr + \frac{1}{2} \iint \frac{\rho(r)\rho(r')}{|r - r'|} dr dr'$$

- f chosen to that only “ v_{xc} missing”

$$E_f(v^\varepsilon) = \underbrace{\inf_{\rho} (f(\rho) + \langle v^\varepsilon, \rho \rangle)}_{\text{“KS equations”}}$$

$$v^\varepsilon = \text{“ZMP } v^\lambda\text{”}$$

- f is not differentiable
- Regularization: $f^\varepsilon(\rho) = \mathcal{Q}_{\text{KS}}^\varepsilon(\rho)$

Also: $\rho_{\text{gs}}^\varepsilon \rightarrow \rho_{\text{gs}}$

- “KS minimization”

$$E_f(v^\varepsilon) = \inf_{\rho} (f(\rho) + \langle v^\varepsilon, \rho \rangle)$$

- Computation of $v^\varepsilon \in -\underline{\partial} f(\rho^\varepsilon)$ is equivalent to $\rho^\varepsilon \in \bar{\partial} E_f(v^\varepsilon)$ by reciprocity relation of convex analysis:

$$-\nabla f^\varepsilon(\rho_{\text{gs}}) = \frac{1}{\varepsilon} J(\rho_{\text{gs}}^\varepsilon - \rho_{\text{gs}}) \in -\underline{\partial} f(\rho^\varepsilon)$$

$$v^\varepsilon(\rho_{\text{gs}}) = \frac{1}{\varepsilon} J(\rho_{\text{gs}}^\varepsilon - \rho_{\text{gs}}) \quad (8)$$

- Use “KS density functional”

$$f(\rho) := \mathcal{Q}_{\text{KS}}(\rho) = T(\rho) + \int v_{\text{ext}}\rho \, \mathrm{d}r + \frac{1}{2} \int \int \frac{\rho(r)\rho(r')}{|r - r'|} \mathrm{d}r \mathrm{d}r'$$

- To obtain ZMP, determine the proximal point ρ^ε and

$$v^\varepsilon := \frac{1}{\varepsilon} J(\rho^\varepsilon - \rho) \in -\underline{\partial} f(\rho^\varepsilon),$$

and let $\varepsilon \rightarrow 0$.

- Solving $\rho^\varepsilon \in \overline{\partial}E_{f=\mathcal{Q}_{\text{KS}}}(v^\varepsilon)$ for ρ^ε amounts to solving Kohn–Sham equations with v^ε in place of v_{xc} .
- Self-consistent scheme:
 - Given an iterate v_i^ε and ρ_i^ε
 - In the next step, we compute ρ_{i+1}^ε from $\rho_{i+1}^\varepsilon \in \overline{\partial}E_f(v_i^\varepsilon)$
 - update $v_{i+1}^\varepsilon := \frac{1}{\varepsilon}J(\rho_{i+1}^\varepsilon - \rho)$.
- For $X = H^{-1}(\Omega)$, Ω bdd, then $\varepsilon^{-1}J$ “almost” gives the ZMP,

$$\frac{1}{\varepsilon}J(\rho^\varepsilon - \rho) = \frac{1}{4\pi\varepsilon} \int_{\Omega} \frac{\rho^\varepsilon(r') - \rho(r')}{|r - r'|} dr' + \text{corrector-term} \quad (9)$$

Thank you