

# **$\nu$ -Representability on Periodic Domains: a Sobolev Space Approach**


---


Sarina Sutter, Markus Penz, Michael Ruggenthaler, Robert van Leeuwen, Klaas Giesbertz

December 03, 2024










Known as  $v$ -representability problem



Known as  $v$ -representability problem

In this talk: Define a setting where we are able to find a (large) set of densities which are (ensemble)  $v$ -representable.

---

<sup>1</sup>E. H. Lieb. "Density Functionals for Coulomb-Systems". In: *Int. J. Quantum Chem.* 24.3 (1983), pp. 243–277. DOI: 10.1002/qua.560240302.

<sup>1</sup>Set of physical densities  $\mathcal{I}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{R}^3), \rho \geq 0, \int \rho = N\}$ .

---

<sup>1</sup>E. H. Lieb. "Density Functionals for Coulomb-Systems". In: *Int. J. Quantum Chem.* 24.3 (1983), pp. 243–277. DOI: 10.1002/qua.560240302.



<sup>1</sup>Set of physical densities  $\mathcal{I}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{R}^3), \rho \geq 0, \int \rho = N\}$ .

Then,  $\mathcal{I}_N \subset L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$  and if  $v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$  then potential energy satisfies

$$\int dx \rho(x) v(x) < \infty.$$

---

<sup>1</sup>E. H. Lieb. "Density Functionals for Coulomb-Systems". In: *Int. J. Quantum Chem.* 24.3 (1983), pp. 243–277. DOI: 10.1002/qua.560240302.

<sup>1</sup>Set of physical densities  $\mathcal{I}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{R}^3), \rho \geq 0, \int \rho = N\}$ .

Then,  $\mathcal{I}_N \subset L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$  and if  $v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$  then potential energy satisfies

$$\int dx \rho(x) v(x) < \infty.$$

The ground state energy is

$$E(v) := \inf \{ \langle \psi, H_v \psi \rangle \mid \psi \in \mathcal{W}_N \}$$

where  $\mathcal{W}_N := \{ \psi \in Q(H_v) \mid \|\psi\|_2 = 1 \}$ .

---

<sup>1</sup>E. H. Lieb. "Density Functionals for Coulomb-Systems". In: *Int. J. Quantum Chem.* 24.3 (1983), pp. 243–277. DOI: 10.1002/qua.560240302.

<sup>1</sup>Set of physical densities  $\mathcal{I}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{R}^3), \rho \geq 0, \int \rho = N\}$ .

Then,  $\mathcal{I}_N \subset L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$  and if  $v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$  then potential energy satisfies

$$\int dx \rho(x) v(x) < \infty.$$

The ground state energy is

$$E(v) := \inf \{ \langle \psi, H_v \psi \rangle \mid \psi \in \mathcal{W}_N \}$$

where  $\mathcal{W}_N := \{ \psi \in Q(H_v) \mid \|\psi\|_2 = 1 \}$ .


On  $\mathcal{I}_N$  we define the functional  $\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \}$  and we get

$$E(v) = \inf_{\rho \in \mathcal{I}_N} \left\{ \tilde{F}(\rho) + \langle \rho, v \rangle \right\}.$$


---

<sup>1</sup>E. H. Lieb. "Density Functionals for Coulomb-Systems". In: *Int. J. Quantum Chem.* 24.3 (1983), pp. 243–277. DOI: 10.1002/qua.560240302.







$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } \mathcal{I}_N$$



$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$




$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$



$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$

$$F(\rho) := \sup \left\{ E(v) - \langle v, \rho \rangle \mid v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3) \right\} \text{ on } L^1(\mathbb{R}) \cap L^3(\mathbb{R}^3)$$






$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$

$$F(\rho) := \sup \left\{ E(v) - \langle v, \rho \rangle \mid v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3) \right\} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$

$F$  is equal to the density matrix approach

$$F_{\text{DM}}(\rho) := \begin{cases} \inf \{ \text{Tr}[\Gamma H_0] \mid \Gamma \rightarrow \rho \} & \text{if } \rho \in \mathcal{I}_N \\ \infty & \text{otherwise.} \end{cases}$$



$$\tilde{F}(\rho) := \inf \{ \langle \psi, H_0 \psi \rangle \mid \psi \rightarrow \rho, \psi \in \mathcal{W}_N \} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$

$$F(\rho) := \sup \left\{ E(v) - \langle v, \rho \rangle \mid v \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3) \right\} \text{ on } L^1(\mathbb{R}^3) \cap L^3(\mathbb{R}^3)$$

$F$  is equal to the density matrix approach

$$F_{\text{DM}}(\rho) := \begin{cases} \min \{ \text{Tr}[\Gamma H_0] \mid \Gamma \rightarrow \rho \} & \text{if } \rho \in \mathcal{I}_N \\ \infty & \text{otherwise.} \end{cases}$$



Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \underline{\partial}F(\rho)$ :

$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \underline{\partial}F(\rho)$ :

$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

$\Rightarrow v$  generates the density  $\rho$ .

Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \underline{\partial}F(\rho)$ :


$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

$\Rightarrow v$  generates the density  $\rho$ .

Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \partial F(\rho)$ :

$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

$\Rightarrow v$  generates the density  $\rho$ .




$F$  locally bounded implies non-empty subdifferential.

$\Rightarrow$  Densities where  $F$  is locally bounded are (ensemble)  $v$ -representable.

Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \partial F(\rho)$ :

$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

$\Rightarrow v$  generates the density  $\rho$ .



$F$  locally bounded implies non-empty subdifferential.


$\Rightarrow$  Densities where  $F$  is locally bounded are (ensemble)  $v$ -representable.



Subgradient  $-v$  of  $F$  at a density  $\rho$ ,  $-v \in \partial F(\rho)$ :

$$F(\rho) - F(\tilde{\rho}) \leq \langle -v, \rho - \tilde{\rho} \rangle \Leftrightarrow F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho}.$$

$\Rightarrow v$  generates the density  $\rho$ .




$F$  locally bounded implies non-empty subdifferential.

$\Rightarrow$  Densities where  $F$  is locally bounded are (ensemble)  $v$ -representable.

Problem: The interior of the domain of  $F$  is empty.




## Empty interior




Element in  $\mathcal{I}_N \Rightarrow F < \infty$


## Empty interior




Element in  $\mathcal{I}_N \Rightarrow F < \infty$



Element not in  $\mathcal{I}_N \Rightarrow F = \infty$




Element not in  $\mathcal{I}_N \Rightarrow F = \infty$




Element not in  $\mathcal{I}_N \Rightarrow F = \infty$


## Empty interior




Element in  $\mathcal{I}_N \Rightarrow F < \infty$



Element not in  $\mathcal{I}_N \Rightarrow F = \infty$



Element not in  $\mathcal{I}_N \Rightarrow F = \infty$



Element not in  $\mathcal{I}_N \Rightarrow F = \infty$

$\Rightarrow$  Include derivative in the norm.

$\Rightarrow$  Use  $H^1$  norm:  $\|f\|_{1,2}^2 = \|f\|_2^2 + \|\nabla f\|_2^2$ .



Recall the set of physical densities  $\mathcal{I}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{T}), \rho \geq 0, \int \rho = N\}$ .

We have

$$\mathcal{I}_N \subset H^1(\mathbb{T}) \hookrightarrow C^0(\mathbb{T}) \hookrightarrow L^3(\mathbb{T}).$$

Recall the set of physical densities  $\mathcal{J}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{T}), \rho \geq 0, \int \rho = N\}$ .

We have

$$\mathcal{J}_N \subset H^1(\mathbb{T}) \hookrightarrow C^0(\mathbb{T}) \hookrightarrow L^3(\mathbb{T}).$$

Potential energy is finite if  $v \in H^{-1}(\mathbb{T})$ . That is,  $v = f + \nabla g$  with  $f, g \in L^2(\mathbb{T})$  and it is paired with  $\rho$  by

$$\langle v, \rho \rangle = \int dx f(x) \rho(x) - \int dx g(x) \nabla \rho(x).$$



Recall the set of physical densities  $\mathcal{J}_N := \{\rho \mid \sqrt{\rho} \in H^1(\mathbb{T}), \rho \geq 0, \int \rho = N\}$ .

We have

$$\mathcal{J}_N \subset H^1(\mathbb{T}) \hookrightarrow C^0(\mathbb{T}) \hookrightarrow L^3(\mathbb{T}).$$

Potential energy is finite if  $v \in H^{-1}(\mathbb{T})$ . That is,  $v = f + \nabla g$  with  $f, g \in L^2(\mathbb{T})$  and it is paired with  $\rho$  by

$$\langle v, \rho \rangle = \int dx f(x) \rho(x) - \int dx g(x) \nabla \rho(x).$$

Example:  $\delta$ -function  $\delta = f + \nabla g$  with  $f(x) = 1$  and  $g(x) = -x$ ,

$$\begin{aligned} \delta(\rho) &= \int_0^1 \rho(x) dx + \int_0^1 x \nabla \rho(x) dx \\ &= \int_0^1 \rho(x) dx - \int_0^1 \rho(x) dx + x \rho(x) \Big|_0^1 = \rho(1). \end{aligned}$$



On  $H^1(\mathbb{T})$  we define  $F(\rho) := \sup \{ E(v) - \langle v, \rho \rangle \mid v \in H^{-1}(\mathbb{T}) \}$ .

On  $H^1(\mathbb{T})$  we define  $F(\rho) := \sup \{ E(v) - \langle v, \rho \rangle \mid v \in H^{-1}(\mathbb{T}) \}$ .

There exists two constants  $C_1, C_2$  such that if  $\rho \in \mathcal{I}_N$  then

$$\|\nabla \sqrt{\rho}\|_2^2 \leq F(\rho) \leq C_1 + C_2 \|\nabla \sqrt{\rho}\|_2^2.$$

On  $H^1(\mathbb{T})$  we define  $F(\rho) := \sup \{ E(v) - \langle v, \rho \rangle \mid v \in H^{-1}(\mathbb{T}) \}$ .

There exists two constants  $C_1, C_2$  such that if  $\rho \in \mathcal{I}_N$  then

$$\|\nabla \sqrt{\rho}\|_2^2 \leq F(\rho) \leq C_1 + C_2 \|\nabla \sqrt{\rho}\|_2^2.$$

For  $\rho \in \mathcal{P}_\eta := \{\rho \in \mathcal{I}_N \mid \rho > \eta > 0\}$  we have the bound

$$\|\nabla \sqrt{\rho}\|_2^2 = \left\| \frac{\nabla \rho}{2\sqrt{\rho}} \right\|_2^2 \leq \frac{1}{4\eta} \|\nabla \rho\|_2^2 \leq \frac{1}{4\eta} \|\rho\|_{H^1}^2.$$

On  $H^1(\mathbb{T})$  we define  $F(\rho) := \sup \{ E(v) - \langle v, \rho \rangle \mid v \in H^{-1}(\mathbb{T}) \}$ .

There exists two constants  $C_1, C_2$  such that if  $\rho \in \mathcal{I}_N$  then


$$\|\nabla \sqrt{\rho}\|_2^2 \leq F(\rho) \leq C_1 + C_2 \|\nabla \sqrt{\rho}\|_2^2.$$

For  $\rho \in \mathcal{P}_\eta := \{\rho \in \mathcal{I}_N \mid \rho > \eta > 0\}$  we have the bound

$$\|\nabla \sqrt{\rho}\|_2^2 = \left\| \frac{\nabla \rho}{2\sqrt{\rho}} \right\|_2^2 \leq \frac{1}{4\eta} \|\nabla \rho\|_2^2 \leq \frac{1}{4\eta} \|\rho\|_{H^1}^2.$$

The set  $\mathcal{P}_\eta$  is open due to  $H^1(\mathbb{T}) \hookrightarrow C^0(\mathbb{T})$ .

$\Rightarrow F$  is locally bounded at  $\rho \in \mathcal{P}_\eta$ .





## Connection to Schrödinger Equation

Problem: Is  $F$  related to the internal energy?

If  $-v \in \underline{\partial} F(\rho)$ , then

$$F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$



## Connection to Schrödinger Equation

Problem: Is  $F$  related to the internal energy?

If  $-v \in \underline{\partial} F(\rho)$ , then

$$F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

We can show that  $F(\rho) = F_{\text{DM}}(\rho)$ . Then,

$$F_{\text{DM}}(\rho) + \langle v, \rho \rangle \leq F_{\text{DM}}(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

## Connection to Schrödinger Equation

Problem: Is  $F$  related to the internal energy?

If  $-v \in \underline{\partial} F(\rho)$ , then

$$F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

We can show that  $F(\rho) = F_{\text{DM}}(\rho)$ . Then,

$$F_{\text{DM}}(\rho) + \langle v, \rho \rangle \leq F_{\text{DM}}(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

Problem: Is the Hamiltonian  $H_v$  with potential  $v$  self-adjoint?

## Connection to Schrödinger Equation

Problem: Is  $F$  related to the internal energy?

If  $-v \in \partial F(\rho)$ , then

$$F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

We can show that  $F(\rho) = F_{\text{DM}}(\rho)$ . Then,

$$F_{\text{DM}}(\rho) + \langle v, \rho \rangle \leq F_{\text{DM}}(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

Problem: Is the Hamiltonian  $H_v$  with potential  $v$  self-adjoint?

### KLMN Theorem

Let  $A$  be a positive self-adjoint operator on a Hilbert space  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}})$  with form domain  $Q(A)$ . Let  $B$  be a symmetric quadratic form on  $Q(A)$  such that there exists  $0 \leq a < 1$  and  $b \in \mathbb{R}$  with

$$|\langle \psi, B\psi \rangle_{\mathcal{H}}| \leq a \langle \psi, A\psi \rangle_{\mathcal{H}} + b \langle \psi, \psi \rangle_{\mathcal{H}}$$

for all  $\psi$  in  $Q(A)$ . Then the operator  $A + B$  is self-adjoint with domain  $Q(A)$  and is bounded from below by  $-b$ .

## Connection to Schrödinger Equation

Problem: Is  $F$  related to the internal energy?

If  $-v \in \partial F(\rho)$ , then

$$F(\rho) + \langle v, \rho \rangle \leq F(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

We can show that  $F(\rho) = F_{\text{DM}}(\rho)$ . Then,

$$F_{\text{DM}}(\rho) + \langle v, \rho \rangle \leq F_{\text{DM}}(\tilde{\rho}) + \langle v, \tilde{\rho} \rangle \quad \text{for all } \tilde{\rho} \in \mathcal{I}_N.$$

Problem: Is the Hamiltonian  $H_v$  with potential  $v$  self-adjoint?

### KLMN Theorem

Let  $A$  be a positive self-adjoint operator on a Hilbert space  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}})$  with form domain  $Q(A)$ . Let  $B$  be a symmetric quadratic form on  $Q(A)$  such that there exists  $0 \leq a < 1$  and  $b \in \mathbb{R}$  with

$$|\langle \psi, B\psi \rangle_{\mathcal{H}}| \leq a \langle \psi, A\psi \rangle_{\mathcal{H}} + b \langle \psi, \psi \rangle_{\mathcal{H}}$$

for all  $\psi$  in  $Q(A)$ . Then the operator  $A + B$  is self-adjoint with domain  $Q(A)$  and is bounded from below by  $-b$ .

Remark: We need to assume that the interaction satisfies the KLMN condition.



## Theorem

Assume that the interaction  $W$  is weakly continuous. Then the minimum of the energy functional  $\mathcal{E}(\psi) := \langle \psi, H_v \psi \rangle$  is attained. Moreover, let  $\psi_0$  be the minimizer. Assume that  $|\langle \psi_0, W\phi \rangle|$  is finite for all  $\phi \in [C^\infty(\mathbb{T}) \otimes \mathbb{C}^2]^{\wedge N}$ . Then  $\psi_0$  satisfies

$$\left( \sum_j -\frac{1}{2} \Delta_j + v(x_j) + W \right) \psi_0 = E_0 \psi_0$$

in a distributional sense, with  $E_0 = \inf_{\psi \in \mathcal{W}_N} \langle \psi, H_v \psi \rangle$ .

## Theorem

Assume that the interaction  $W$  is weakly continuous. Then the minimum of the energy functional  $\mathcal{E}(\psi) := \langle \psi, H_v \psi \rangle$  is attained. Moreover, let  $\psi_0$  be the minimizer. Assume that  $|\langle \psi_0, W\phi \rangle|$  is finite for all  $\phi \in [C^\infty(\mathbb{T}) \otimes \mathbb{C}^2]^{\wedge N}$ . Then  $\psi_0$  satisfies

$$\left( \sum_j -\frac{1}{2} \Delta_j + v(x_j) + W \right) \psi_0 = E_0 \psi_0$$

in a distributional sense, with  $E_0 = \inf_{\psi \in \mathcal{W}_N} \langle \psi, H_v \psi \rangle$ .

*Remark:* An interaction of the form  $W = \sum_{i < j} w(x_i - x_j)$  with  $w \in L^1([-1, 1])$  and a distributional interaction of the form  $W = \sum_{i < j} \nabla_{x_i} g(x_i - x_j)$  with  $g \in L^2([-1, 1])$  satisfy all condition on  $W$ .





## Zero Temperature

$$E(\nu) = \inf_{\psi} \langle \psi, H_{\nu} \psi \rangle$$

## Elevated Temperature

$$\Omega_{\beta}(\nu) = \inf_{\Gamma} \text{Tr}[\Gamma(H_{\nu} + \beta^{-1} \log \Gamma)]$$

## Zero Temperature

$$E(\nu) = \inf_{\psi} \langle \psi, H_{\nu} \psi \rangle$$

$$\tilde{F}(\rho) = \inf_{\psi \rightarrow \rho} \langle \psi, H_0 \psi \rangle$$

## Elevated Temperature

$$\Omega_{\beta}(\nu) = \inf_{\Gamma} \text{Tr}[\Gamma(H_{\nu} + \beta^{-1} \log \Gamma)]$$

## Zero Temperature

$$E(v) = \inf_{\psi} \langle \psi, H_v \psi \rangle$$

$$\tilde{F}(\rho) = \inf_{\psi \rightarrow \rho} \langle \psi, H_0 \psi \rangle$$

$$F_{\text{DM}}(\rho) = \min_{\Gamma \rightarrow \rho} \text{Tr}[\Gamma H_0]$$

## Elevated Temperature

$$\Omega_{\beta}(v) = \inf_{\Gamma} \text{Tr}[\Gamma(H_v + \beta^{-1} \log \Gamma)]$$

$$F_{\text{DM}}^{\beta}(\rho) = \inf_{\Gamma \rightarrow \rho} \text{Tr}\{\Gamma(H_0 + \beta^{-1} \log(\Gamma))\}$$

## Zero Temperature

$$E(v) = \inf_{\psi} \langle \psi, H_v \psi \rangle$$

$$\tilde{F}(\rho) = \inf_{\psi \rightarrow \rho} \langle \psi, H_0 \psi \rangle$$

$$F_{\text{DM}}(\rho) = \min_{\Gamma \rightarrow \rho} \text{Tr}[\Gamma H_0]$$

$$\partial F_{\text{DM}}(\rho) \neq \emptyset \text{ for } \rho > 0$$

## Elevated Temperature

$$\Omega_{\beta}(v) = \inf_{\Gamma} \text{Tr}[\Gamma(H_v + \beta^{-1} \log \Gamma)]$$

$$F_{\text{DM}}^{\beta}(\rho) = \inf_{\Gamma \rightarrow \rho} \text{Tr}\{\Gamma(H_0 + \beta^{-1} \log(\Gamma))\}$$

$$\partial F_{\text{DM}}^{\beta}(\rho) \neq \emptyset \text{ for } \rho > 0$$

## Zero Temperature

$$E(v) = \inf_{\psi} \langle \psi, H_v \psi \rangle$$

$$\tilde{F}(\rho) = \inf_{\psi \rightarrow \rho} \langle \psi, H_0 \psi \rangle$$

$$F_{\text{DM}}(\rho) = \min_{\Gamma \rightarrow \rho} \text{Tr}[\Gamma H_0]$$

$$\partial F_{\text{DM}}(\rho) \neq \emptyset \text{ for } \rho > 0$$

## Elevated Temperature

$$\Omega_{\beta}(v) = \inf_{\Gamma} \text{Tr}[\Gamma(H_v + \beta^{-1} \log \Gamma)]$$

$$F_{\text{DM}}^{\beta}(\rho) = \inf_{\Gamma \rightarrow \rho} \text{Tr}\{\Gamma(H_0 + \beta^{-1} \log(\Gamma))\}$$

$$\partial F_{\text{DM}}^{\beta}(\rho) \neq \emptyset \text{ for } \rho > 0$$

v-rep. densities satisfy  $\rho > 0$ .

## Zero Temperature

$$E(v) = \inf_{\psi} \langle \psi, H_v \psi \rangle$$

$$\tilde{F}(\rho) = \inf_{\psi \rightarrow \rho} \langle \psi, H_0 \psi \rangle$$

$$F_{\text{DM}}(\rho) = \min_{\Gamma \rightarrow \rho} \text{Tr}[\Gamma H_0]$$

$$\partial F_{\text{DM}}(\rho) \neq \emptyset \text{ for } \rho > 0$$

## Elevated Temperature

$$\Omega_{\beta}(v) = \inf_{\Gamma} \text{Tr}[\Gamma(H_v + \beta^{-1} \log \Gamma)]$$

$$F_{\text{DM}}^{\beta}(\rho) = \inf_{\Gamma \rightarrow \rho} \text{Tr}\{\Gamma(H_0 + \beta^{-1} \log(\Gamma))\}$$

$$\partial F_{\text{DM}}^{\beta}(\rho) \neq \emptyset \text{ for } \rho > 0$$

v-rep. densities satisfy  $\rho > 0$ .

$\partial F_{\text{DM}}^{\beta}(\rho)$  is single-valued.



System of  $N$  fermions on the one dimensional torus  $\mathbb{T}$ .



System of  $N$  fermions on the one dimensional torus  $\mathbb{T}$ .

Main result: Densities in  $H^1(\mathbb{T})$  that are gapped away from 0 are (ensemble)  $v$ -representable by a  $H^{-1}$  potential  $v$ .

System of  $N$  fermions on the one dimensional torus  $\mathbb{T}$ .

Main result: Densities in  $H^1(\mathbb{T})$  that are gapped away from 0 are (ensemble)  $v$ -representable by a  $H^{-1}$  potential  $v$ .

Ground state of a Hamiltonian  $H_v$  exists and satisfies the Schrödinger equation in a distributional sense.

System of  $N$  fermions on the one dimensional torus  $\mathbb{T}$ .

Main result: Densities in  $H^1(\mathbb{T})$  that are gapped away from 0 are (ensemble)  $v$ -representable by a  $H^{-1}$  potential  $v$ .

Ground state of a Hamiltonian  $H_v$  exists and satisfies the Schrödinger equation in a distributional sense.

For elevated temperature and fixed number of particles, we have a full characterization of  $H^{-1}$   $v$ -representable densities and  $F_{\text{DM}}^\beta$  is Gâteaux differentiable at densities  $\rho > 0$ .