

# A Practical Guide to Quantum Linear Algebra

Fabian M. Faulstich

Department of Mathematics, Rensselaer Polytechnic Institute,  
NY



# Quantum Computing in a Nutshell

## – Keeping Quantum Computing Honest

- Exciting experiment
- New computational paradigm
- Is it competitive?  
→ No
- Not – and never will be(?) – a silver bullet



# Linear Algebra

$$Ax = b \quad \begin{aligned} Av &= \lambda v \\ (H|\Psi\rangle &= E|\Psi\rangle) \end{aligned}$$

# Linear Algebra

$$Ax = b \quad Av = \lambda v$$
$$(H|\Psi\rangle = E|\Psi\rangle)$$

- Underpins most of scientific computing
- Efficient algorithm development and implementation
  - BLAS
  - LAPACK
  - ATLAS
  - ...

(Turing Award 2021 – Jack Dongarra)

Question: Can we do similar things with a quantum computer?

# Quantum Linear Algebra: Desired Speedup

$\text{poly}(N)$       vs.       $\text{poly log}(N)$

Consider  $N$ -dimensional systems (say  $N = 2^n$ )

- Most classical simulations are  $\text{poly}(N)$ 
  - Storing a vector  $\sim N$
  - Sparse MAT-VEC  $\sim N$
- Quantum simulations we hope(!) for  $\text{poly log}(N)$ 
  - An  $n$ -qubit state can be viewed as a  $2^n$ -dim. unit vector
  - Matrix operations are implemented as quantum operations on these  $n$  qubits

!Careful!

!Caveats apply!

# Rules of Quantum Simulation

In general, not all operations can be performed efficiently on a quantum device!

- Vectors (quantum states) are normalized
- Only a subset of operations can be efficiently implemented:  
unitary matrices
  - Other problems have to be rescaled (Block encoding)
  - Rescaling will introduce greater cost
- No cloning theorem
  - Iterative methods are not in general efficient

# Quantum Algorithms for Scientific Computation

Lecture notes by Lin Lin:



- arXiv:2201.08309
- [math.berkeley.edu/~linlin/](http://math.berkeley.edu/~linlin/)



# Roadmap



Towards linear algebra on quantum devices

1. Basic quantum gates
2. (Real) Hadamard Test
3. Quantum Phase Estimation
4. Numerical performances on emulators  
and a real quantum device

# Some Basic Gates & Conventions

Single qubit states:

$$|0\rangle := \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad |1\rangle := \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Single qubit gates, e.g.,

$$X := \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad H := \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

Two-qubit gates, e.g.,


$$CNOT := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \quad \text{and} \quad SWAP := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# The (real) Hadamard Test

- Simplest way to compute


$$\langle U \rangle = \langle \psi | U | \psi \rangle$$

- Simple diagram



Why does it work?

## The real Hadamard Test – Outcome




$$|\pi_0\rangle = |0\rangle \otimes |\psi\rangle = |0\rangle|\psi\rangle$$

$$|\pi_1\rangle = H|0\rangle I|\psi\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)|\psi\rangle = |+\rangle|\psi\rangle$$

$$|\pi_2\rangle = (|0\rangle\langle 0| \otimes I + |1\rangle\langle 1| \otimes U)|\pi_1\rangle = \frac{1}{\sqrt{2}}|0\rangle I|\psi\rangle + \frac{1}{\sqrt{2}}|1\rangle U|\psi\rangle$$

$$|\pi_3\rangle = |0\rangle \frac{I+U}{2}|\psi\rangle + |1\rangle \frac{I-U}{2}|\psi\rangle$$

## The real Hadamard Test – Outcome



We then perform a (single qubit) measurement and find

$$\mathbb{P}_{\text{Re}|0\rangle} = |\langle 0|\pi_3 \rangle|^2 = \frac{2 + \langle \psi|(U + U^\dagger)|\psi \rangle}{4} = \frac{1}{2}(1 + \text{Re}\langle \psi|U|\psi \rangle)$$

and

$$\mathbb{P}_{\text{Re}|1\rangle} = |\langle 1|\pi_3 \rangle|^2 = \frac{1}{2}(1 - \text{Re}\langle \psi|U|\psi \rangle)$$

## The real Hadamard Test – Experiment

Consider

$$U = \begin{bmatrix} 1 & 0 \\ 0 & e^{i\theta} \end{bmatrix} \quad \text{where} \quad \theta = \frac{1}{2} + \frac{1}{2^4} = 0.5625$$

Then

$$\lambda_1 = 1, v_1 = |0\rangle \quad \text{and} \quad \lambda_2 = e^{i\theta} = \cos(\theta) + i \sin(\theta), v_2 = |1\rangle$$

and

$$\langle 1|U|1\rangle = \cos(\theta) + i \sin(\theta)$$

# The real Hadamard Test – Experiment

Initializing with

$$|\Psi\rangle = |1\rangle$$

yields

$$\text{Re}\langle 1|U|1\rangle = \cos(\theta)$$


We find

$$\mathbb{P}_{\text{Re}|0\rangle} = \frac{1}{2}(1 + \cos(\theta)) \approx 0.9230$$

$$\mathbb{P}_{\text{Re}|1\rangle} = \frac{1}{2}(1 - \cos(\theta)) \approx 0.0770$$

Quantum simulation:

- AerSimulator
- Real Hadamard test
- 100 samples



## The real Hadamard Test – Convergence

- Every sample is a random variable with two outcomes:

$$|0\rangle \quad \text{or} \quad |1\rangle$$

$\Rightarrow$  Bernoulli distribution

$$X_i \sim \mathcal{B}(p) \quad \text{with} \quad p = \mathbb{P}_{\text{Re}|0\rangle}$$

- Given  $N_s$  i.i.d. samples, we estimate

$$\hat{p}_{N_s} = \frac{1}{N_s} \sum_{i=1}^{N_s} \mathbb{1}_{\{|0\rangle\}}(X_i)$$

[empirical distribution function]

- By the weak law of large numbers

$$\hat{p}_{N_s} \xrightarrow{\mathbb{P}} \mathbb{E}(X) \quad \Leftrightarrow \quad \lim_{N_s \rightarrow \infty} \mathbb{P}(|\hat{p}_{N_s} - p| > \varepsilon) = 0 \quad \forall \varepsilon > 0$$

## The real Hadamard Test – Convergence rate


- For the empirical distribution function the STD is

$$\sigma^2 = \mathbb{V}(\hat{p}_{N_s}) = \frac{p(1-p)}{N_s}$$

- Then

$$\epsilon \geq \sigma = \sqrt{\frac{p(1-p)}{N_s}} \quad \Rightarrow \quad N_s \geq \frac{p(1-p)}{\epsilon^2} \in \Omega\left(\frac{1}{\epsilon^2}\right)$$


# The real Hadamard test – Convergence rate



⇒ Converges slowly... very slowly!  
(like molasses in January)

# Quantum Phase Estimation (QPE)

- The Hadamard test performs a single qubit measurement
- Can we achieve higher accuracy measuring multiple qubits?  
⇒ Yes, this is the idea of QPE



- The rate of convergence:

$$n \in \Omega\left(\frac{1}{\epsilon}\right)$$



## QPE – Convergence

- Consider

$$U = \begin{bmatrix} 1 & 0 \\ 0 & e^{i2\pi\theta} \end{bmatrix}$$



- 4 ancilla qubits
- phase angles  $\theta = 1/4$  and  $\theta = 2/3$

QPE Probability Distribution



# QPE – Computing the Phase



QPE Probability Distribution



- Find the maximal state:  
0100 and 1011, respectively

# QPE – Computing the Phase


QPE Probability Distribution



- Find the maximal state:  
0100 and 1011, respectively
- Convert to decimal basis:  
4 and 11, respectively

# QPE – Computing the Phase

QPE Probability Distribution




- Find the maximal state:  
0100 and 1011, respectively
- Convert to decimal basis:  
4 and 11, respectively
- Normalize w.r.t. number ancilla qubits ( $2^n$ ):  
 $4/16 = 1/4$  and  $11/16 = 0.6875 \approx 2/3$ , respectively

# Comparison and quantum computation

- Consider

$$U = \begin{bmatrix} 1 & 0 \\ 0 & e^{i2\pi\frac{1}{3}} \end{bmatrix}$$

- 1 to 20 ancilla qubits for QPE (each at  $10^3$  shots)
- $10^3$  to  $10^5$  shots for HT




Why is QPE on the quantum device so bad?

# Quantum Simulation – Transpilation

Original circuit:




Transpiled circuit:  
(Qiskit 1.1.1, optimization\_level=3)




## 2-qubit gate depth

- The dominant noise contribution comes from two-qubit gates  
e.g. controlled or swap gates



# Quantum Simulation – Runtime



⇒ Quantum simulation of QPE has constant runtime

# High-accuracy electronic structure methods

**SUMMER SCHOOL**  
**MATHEMATICAL INTRODUCTION TO**  
**HIGH-ACCURACY ELECTRONIC**  
**STRUCTURE THEORY**  
JUNE 30 - JULY 10, 2025

**OPEN TO JUNIOR GRADUATE STUDENTS AND  
SENIOR UNDERGRADUATE STUDENTS**

The purpose of the Summer School is to provide students with a self-contained introduction to electronic structure theory, spanning from the basic analytical investigations of the Schrödinger equations to hands-on computational experiences using state-of-the-art techniques e.g. coupled cluster and tensor network methods.

**APPLICATION DEADLINE:**  
**DECEMBER 15, 2024**

Organizers:  
Fabian Faulstich (Rensselaer Polytechnic Institute)  
Yuehaw Khoo (University of Chicago)

FUNDING FOR TRAVEL AND LODGING IS AVAILABLE FOR ACCEPTED APPLICANTS WHO ARE NOT LOCAL TO CHICAGO.

NSF National Science Foundation  
The Institute for Mathematical and Statistical Innovation (IMSI) is funded by NSF grant DMS-1929348

1115 EAST 60TH STREET  
CHICAGO, ILLINOIS 60637  
INFO@IMSI.INSTITUTE

**imsi** Institute for Mathematical and Statistical Innovation

## What:

Introduction to high-accuracy electronic structure methods

## Where:

Institute for Mathematical and Statistical Innovation (IMSI)  
Chicago, Illinois, USA

## When:

June 30 – July 10, 2025

## Funding is available!

Application deadline: Dec. 15



# Team



Alexander Weiss



Paul Bruzzi

Thank you for your attention!