#### ERGA Assembly Report

v24.10.15

Tags: ERGA-BGE

TxID	1172132	
ToLID	cbLewAcum8.1	
Species	Lewinskya acuminata	
Class	Bryopsida	
Order	Orthotrichales	

Genome Traits	Expected	Observed
Haploid size (bp)	273,294,980	257,074,491
Haploid Number	6 (source: ancestor)	6
Ploidy	1 (source: ancestor)	2
Sample Sex	Unknown	Unknown

#### EBP metrics summary and curation notes

Obtained EBP quality metric for collapsed: 7.7.Q54

The following metrics were automatically flagged as below EBP recommended standards or different from expected:

- . Observed Ploidy is different from Expected
- . BUSCO single copy value is less than 90% for collapsed
- . BUSCO duplicated value is more than 5% for collapsed

#### Curator notes


- . Interventions/Gb: 84
- . Contamination notes: ""
- Other observations: "The assembly of LEWINSKYA ACUMINATA (cbLewAcum8) is based on 69X PacBio data and 202X Arima Hi-C data generated as part of the European Reference Genome Atlas (ERGA, https://www.erga-biodiversity.eu/) via the Biodiversity Genomics Europe project (BGE, https://biodiversitygenomics.eu/). The assembly process included the following steps: initial PacBio assembly generation with Nextdenovo, removal of contaminant sequences using Context, removal of haplotypic duplications using purge\_dups, and Hi-C-based scaffolding with YaHS. In total, 32 regions totaling 4 Mb were identified as haplotypic duplications and removed. Additionally, 114 contigs were identified as contaminants (bacterial, archaeal, or viral), totaling 23 Mb (with the largest being 4 Mb). The mitochondrial genome was assembled using OATK. Finally, the primary assembly was analyzed and manually improved using Pretext. During manual curation, 1 haplotypic regions and 5 contaminant sequences were removed, totaling 54,950 pb and 456,681 pb, respectively (with the largest being 54,950 pb and 162,112 pb). Chromosome-scale scaffolds confirmed by Hi-C data were named in order of size and contigs were already mostly chromosome-scale. "

# Quality metrics table

Metrics	Pre-curation collapsed	Curated collapsed
Total bp	257,229,379	257,074,491
GC %	37.5	37.51
Gaps/Gbp	0	182.83
Total gap bp	0	5,900
Scaffolds	63	16
Scaffold N50	10,772,754	40,771,022
Scaffold L50	7	3
Scaffold L90	22	6
Contigs	63	63
Contig N50	10,772,754	10,772,754
Contig L50	7	7
Contig L90	22	23
QV	54.3675	54.4163
Kmer compl.	97.548	97.5146
BUSCO sing.	77.1%	77.0%
BUSCO dupl.	5.7%	5.7%
BUSCO frag.	2.0%	2.0%
BUSCO miss.	15.2%	15.3%


BUSCO: 5.4.3 (euk\_genome\_met, metaeuk) / Lineage: embryophyta\_odb10 (genomes:50, BUSCOs:1614)

HiC contact map of curated assembly




collapsed [LINK]

# K-mer spectra of curated assembly




Distribution of k-mer counts per copy numbers found in asm




Distribution of k-mer counts coloured by their presence in reads/assemblies

## Post-curation contamination screening



## TAPAs summary Graph



collapsed. Bubble plot circles are scaled by sequence length, positioned by coverage and GC proportion, and coloured by taxonomy. Histograms show total assembly length distribution on each axis.

## Data profile

Data	PACBIO Hifi	Arima
Coverage	69	202

#### Assembly pipeline

#### Curation pipeline

Submitter: Lola Demirdjian Affiliation: Genoscope

Date and time: 2025-02-12 11:33:33 CET